

ALABAGRUS ENDERLEIN (HYMENOPTERA, BRACONIDAE, AGATHIDINAE) SPECIES OF COSTA RICA, WITH AN EMPHASIS ON SPECIMENS REARED FROM CATERPILLARS IN ÁREA DE CONSERVACIÓN GUANACASTE¹

MICHAEL J. SHARKEY,^{2–5} SARAH MEIEROTTO,² ERIC CHAPMAN,² DANIEL JANZEN,³ WINNIE HALLWACHS,³ TANYA DAPKEY,³ AND M. ALMA SOLIS⁴

ABSTRACT. Eighty-six species of Costa Rican *Alabagrus* are treated; these include species reared from lepidopteran larvae in Área de Conservación Guanacaste, Costa Rica, over 32 years of caterpillar inventory. Sixty-five new species are described, that is, *A. almasolisae*, *A. andresfreitasi*, *A. andywarreni*, *A. axelhausmanni*, *A. barbsharanowskiae*, *A. bernardospinozai*, *A. bobpoolei*, *A. bobrobbinsi*, *A. bobwhartoni*, *A. brendameierottoae*, *A. brianbrowni*, *A. brianharrisi*, *A. craigevansi*, *A. donharveyi*, *A. donlafontainei*, *A. fernandezi*, *A. fernandodiasi*, *A. genemonroei*, *A. hansonii*, *A. hespenheidei*, *A. iankitchingi*, *A. ilgookangi*, *A. isidrochaconi*, *A. jackiemillerae*, *A. jeanfrancoislandryi*, *A. jeanmariecadioui*, *A. jennyphillipsae*, *A. jimmilleri*, *A. johnbrowni*, *A. johnburnsi*, *A. johnobryckii*, *A. johnsharkeyi*, *A. kaciejoae*, *A. karenscharkeyae*, *A. kaydodgeae*, *A. keithwillmotti*, *A. leedyeri*, *A. lindapitkinae*, *A. longinoi*, *A. malcolmscoblei*, *A. marcepsteini*, *A. mariaheikkilae*, *A. markmetzi*, *A. mattottoi*, *A. nickgrishini*, *A. patsharkeyi*, *A. paulgoldsteini*, *A. paulheberti*, *A. paulsharkeyi*, *A. paulthiaucourti*, *A. quickei*, *A. ramyamanjunathae*, *A. reddyapallii*, *A. rudolfmeieri*, *A. sarahmeierottoae*, *A. sarahsharkeyae*, *A. scottmilleri*, *A. scottshawii*, *A. semihesperheidei*, *A. stiremani*, *A. tanyadapkeyae*, *A. tommyersi*, *A. victoriapookae*, *A. yuanmaofangi*, and *A. yuchinkengae*.

Species limits primarily based on cytochrome *c* oxidase subunit I mitochondrial DNA sequence data greatly differed from previous morphological attempts to delimit species of *Alabagrus*. As a result, 17 species reported to occur in Costa Rica are no longer considered to be present in the country, that is, *A. albispina*, *A. imitatus*, *A. juchuy*, *A. kagaba*, *A. latisoma*, *A. latreillei*, *A. maya*, *A. mojos*, *A. nahuatl*, *A. nigrilitus*, *A. pachamama*, *A. paruyana*, *A. parvifaciatus*, *A. semialbus*, *A. tricarainatus*, *A. tripartitus*, and *A. warrau*. Furthermore, five species have been found to be composed of species complexes, that is, *A. cocto*, *A. englishi*, *A. pecki*, *A. roibasi*, and *A. yaruro*. These difficulties point to the impossible task of delimiting species of Agathidinae solely with morphological evidence.

An illustrated key to species, and a plate of color photos of each species are provided.

INTRODUCTION

This article is the third in a series revising the species of Agathidinae reared from lepidopteran caterpillars as part of the inventory of the caterpillars and their parasitoids of Área de Conservación Guanacaste (ACG), northwestern Costa Rica (<http://janzen.sas.upenn.edu>; <http://www.acguanacaste.ac.cr>; Janzen et al., 2009). It includes a key to 87 species of *Alabagrus* Enderlein, of which 65 are newly described. This is not meant to be a comprehensive treatment of the Costa Rican fauna of *Alabagrus*. A conservative estimate, based on extensive collections at Instituto Nacional de Biodiversidad (INBio; Costa Rica), the Universidad de Costa Rica, and the Hymenoptera Institute (Lexington, Kentucky), is that about 300 species occur in the country.

New World members of *Alabagrus* may be distinguished from other agathidines by the following combination of character states: simple claws with a basal lobe, frons bordered laterally by carinae; first metasomal median tergite distinctly convex and/or with a median ridge, gena not elongate. All are presumed to be koinobiont endoparasitoids of concealed larvae of Crambidae. *Pharpa* Sharkey, 1988, is the sister group to *Alabagrus* (Sharkey, 1988) and is used here

as the outgroup for the phylogenetic analyses. Both genera are restricted to the New World.

We include a molecular data set consisting of 680 *Alabagrus* operational taxonomic units (OTUs) sequenced for cytochrome *c* oxidase subunit I (*COI*) for approximately 60 species. The majority of specimens (624) were reared from caterpillars (those with specimen numbers beginning with DHJPAR).

Alabagrus was first revised by Sharkey (1988), who included 104 species. Leathers and Sharkey (2003) revised the species of *Alabagrus* of La Selva, and included previously described species known to occur in Costa Rica. They treated 39 putative species, including six new species. These two revisions relied solely on morphological evidence. The species concepts in the present treatment were formulated with the assistance of *COI* sequence data and a plethora of host data as the result of rearings in Área de Conservación de Guanacaste. These data have had a profound effect. There are many more cryptic species than previously thought, and many species with presumed wide distributions are more likely to be species complexes whose component species have relatively restricted distributions. As a result, 17 species, reported to occur in Costa Rica by Leathers and Sharkey (2003), are no longer considered to be present in the country, these are *Alabagrus albispina*, *A. imitatus*, *A. juchuy*, *A. kagaba*, *A. latisoma*, *A. latreillei*, *A. maya*, *A. mojos*, *A. nahuatl*, *A. nigrilitus*, *A. pachamama*, *A. paruyana*, *A. parvifaciatus*, *A. semialbus*, *A. tricarainatus*, *A. tripartitus*, and *A. warrau*. Furthermore, five of the species reported by Leathers and Sharkey (2003) have been found to be composed of species complexes, that is, *Alabagrus cocto*, *A. englishi*, *A. pecki*, *A. roibasi*, and *A. yaruro*. These difficulties point to the impossible task of delimiting species of Agathidinae solely with morphological evidence.

¹ URL: www.nhm.org/scholarlypublications

² Department of Entomology, University of Kentucky, S-225 Agricultural Science Center North, Lexington, Kentucky 40546-0091, USA.

³ Department of Biology, University of Pennsylvania, 3740 Hamilton Walk, Philadelphia, Pennsylvania 19104, USA.

⁴ Systematic Entomology Laboratory, Agriculture Research Service, U.S. Department of Agriculture, Smithsonian Institution National Museum of Natural History, E-517, MRC 168, Washington, D.C. 20013-7012, USA.

⁵ Corresponding author: Michael J. Sharkey, E-mail: msharkey@uky.edu

METHODS

SPECIMEN INFORMATION

Most specimens used in this study are deposited in the Entomology Museum of Utah State University and the Hymenoptera Institute Collection, with the exception of specimens with numbers beginning with “BIOUG,” which are deposited at the University of Guelph, Centre for Biodiversity Genomics, Ontario, Canada. Source files for the keys, descriptions, illustrations, DNA sequences and distributional data are all freely available to future researchers who may wish to build on these records. The detailed specimen records are available by search of the individual specimen DHJPARxxxxxxx voucher codes at the Janzen database (<http://janzen.bio.upenn.edu/caterpillars/database.lasso>). Some of the Lepidoptera hosts are incompletely identified; however, they also have unique names, such as *Desmia* Solis19 (which is an interim name for *Desmia* species 19 as determined by M. Alma Solis of the U.S. Department of Agriculture Systematic Entomology Laboratory, Washington, D.C.). These names will be updated in the Janzen database when the species is baptized with a formal scientific name, but the interim name, in this case *Desmia* Solis19, will remain searchable in that database. Host caterpillars are uniquely identified by their own voucher code system, which is recognizable by YY-SRNP-XXXXX, where YY is the two-digit year and XXXXX is a unique number within that year.

DNA trace files and primer information are available through the Barcode of Life Data Systems (BOLD, <http://www.boldsystems.org>; Ratnasingham and Hebert, 2007), as is specimen information for specimen numbers INBxxxxxxxxxx, INBIOCRIxxxxxxxxxx, and BIOUGxxxxx-xxx. Specimen information for specimens with numbers beginning with the letter “H” are stored in the Symbiota database (Gries et al., 2014; Symbiota Collections of Arthropods Network, <http://symbiota4.acis.ufl.edu/scan/portal/>) under the Hymenoptera Institute Collection (HIC). To search for a specimen in the database, do the following: Under the *Search* tab (upper left of the screen), select *Search Collections*. Deselect all collections, and scroll down the Southeast section, put a check in the box next to *Hymenoptera Institute Collection*, scroll back up and hit the *Search* button (right side of screen). H-numbered specimens are stored with a four-letter prefix (HICH) followed by a six-digit number. Therefore, H369, as it appears in this publication, is stored as HICH000369. To search for this specimen, scroll down to the Specimen Criteria section, type this number in the box next to *Catalog Number* and hit the search button. This displays a page with a summary of the specimen information. Clicking on *Full Record Details* opens a new window with the full specimen record, including all available images.

Each species is diagnosed and described or redescribed, and a plate of images with a chronogram of occurrence is included for each.

SPECIES CONCEPTS

The biological species concept is used with evidence drawn from *COI* sequences, morphology, and host use as indirect evidence for species integrity. *COI* was the primary source of evidence with no particular percent divergence employed as a cut-off. The sequence data were checked against morphological and host-use data, and these corroborated or (rarely) refuted the sequence data.

TAXON SAMPLING

For molecular analyses, we used all *Alabagrus* sequences available to us on the BOLD database from Costa Rica. Once we removed a few sequences that were contaminated in some way, this amounted to 636 sequences. To this, we added 44 *Alabagrus* sequences generated by the Sharkey lab from Costa Rica, Honduras, Mexico, and USA.

MORPHOLOGICAL TERMS

Morphological terms are from Sharkey and Wharton (1997). The reader is advised to search for any morphological term that is not understood on the Hymenoptera Anatomy Ontology website (<http://portal.hymao.org/projects/32/public/ontology/>; Yoder et al., 2010). To do this, enter the term in the search box, select the best match from the list that pops up, and click on the *show* button.

MUSEUM ACRONYMS

CNCI Canadian National Collection of Insects, Ottawa, Canada
 EMUS Utah State University, Department of Biology, Entomology Museum, Logan, Utah, USA
 HIC University of Kentucky, Department of Entomology, Hymenoptera Institute Collection, Lexington, Kentucky, USA
 INBio Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, Costa Rica
 MUCR Ciudad Universitaria, Universidad de Costa Rica, Museo de Insectos, Costa Rica

DNA EXTRACTION, PCR, AND SEQUENCING

Specimens sequenced in this study are from a variety of sources. Specimens with numbers beginning with DHJPAR, INB, INBIOCRI, or BIOUG were extracted and sequenced for *COI* at the University of Guelph following the methods outlined in Hebert et al. (2004).

Specimens with numbers beginning with the letter “H” were extracted from individual legs with the QIAGEN DNeasy Blood and Tissue Kit using the animal tissue protocol (QIAGEN Inc., Chatsworth, California, USA). The mitochondrial *COI* (~658 bp) gene was amplified with the primer pair LepF1 and LepR1 (Hebert et al., 2004). Polymerase chain reaction (PCR) was conducted using TaKaRa reagents, with each reaction consisting of 1× buffer, 0.3 mM nucleotides, 0.4 μM of each primer, 0.625 U TaKaRa *Ex Taq*, double-distilled water, and 1–3 μL template DNA in a total reaction volume of 25 μL. The thermal cycling protocol had an initial denaturation period at 95 °C for 2.5 min, followed by 40 cycling steps which denatured at 95 °C for 30 s, annealed at 44 °C for 30 s and extended at 68 °C for 45 s, with a final extension step of 72 °C for 7 min. To determine reaction success, PCR products were electrophoresed in 1% agarose stained with ethidium bromide. PCR products were outsourced for Sanger sequencing either by the Advanced Genetic Technologies Center (University of Kentucky, Lexington, Kentucky, USA) or Beckman Coulter Genomics (Danvers, Massachusetts, USA) using labeled dideoxy-nucleotides with ABI 3730, Big-Dye Terminator mix v. 3.0, or with ABI PRISM 3730xl, BigDye Terminator mix v. 3.1 (Applied Biosystems, Foster City, California, USA).

DNA ASSEMBLY AND PHYLOGENETIC ANALYSIS

Bidirectional sequences were aligned and edited using Geneious Pro (v. 6.1.5; Drummond et al., 2009) and multiple alignments were assembled using MAFFT (v. 5; Katoh et al., 2006) using the default settings and refined by eye using MacClade v. 4.08 (Maddison and Maddison, 2005). Maximum likelihood (ML) phylogenetic analyses were conducted on the *COI* data set using Garli (v. 2.01; Zwickl, 2006). The data set consisted of 680 *Alabagrus* OTUs and one sequence from a species of *Pharpa* Sharkey, which was shown to be sister to *Alabagrus* in the most comprehensive molecular agathidine phylogeny to date (Sharkey and Chapman, 2017) for outgroup rooting. The data were partitioned by codon position (three partitions). We applied the most complex model available (GTR+I+G; Rodriguez et al., 1990) to each partition as per recommendations of Huelsenbeck and Rannala (2004) for likelihood-




Figure 1 Tree of highest posterior probability from a 10 million-generation Bayes analysis of *COI*. Species are collapsed (when possible) into single terminals (terminal triangles), with the number of OTUs for each collapsed species in parentheses. The length of the triangles represents the branch length from the node to the tip of the longest branch.

based analyses. We conducted a 50-replicate ML search for the tree of highest log-likelihood and a 200-replicate ML bootstrap analysis (Felsenstein, 1985). Both analyses used the default settings.

A Bayesian inference (BI) phylogenetic analysis was also conducted on the *COI* data set with MrBayes (v. 3.1.2; Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003). As in the ML analyses, the data were partitioned by codon position. To allow each partition to have its own set of parameter estimates, *revmat*, *tratio*, *statefreq*, *shape*, and *pinvar* were all unlinked during the analyses. To obtain the most accurate branch length estimates possible, the option *prset ratepr=variable* (which assigns a separate branch length parameter for each partition) was employed as per the recommendations of Marshall et al. (2006). A temperature (*t*) setting of 0.01 was used because there was insufficient chain swapping in a pilot run with the default value *t*=0.1. With *t*=0.01, the average chain swapping value was 55%. Two independent, simultaneous BI searches were run for 10 million generations, saving a tree every 1000 generations, with four search chains each. The average standard deviation of split frequencies was 0.0235 at the completion of 10 million generations. The log-likelihood values of the chains in both analyses plateaued just above −10,000 by 3 million generations and did not go any lower by the completion of the run. The likelihood of best state for the “cold” chain of run 1 was −9911.85 and that for the “cold” chain of run 2 was −9912.76. The 5000 post-burn-in trees from each run (10,000 total), using a 50% burn-in were used to calculate the majority rule consensus tree using PAUP* (v. 4.0β10; Swofford, 2003). The data set analyzed herein, including the MrBayes command block and Garli configuration files are available from the authors upon request.

RESULTS

PHYLOGENETIC TREES

The tree of highest posterior probability from the Bayesian analysis with species collapsed into single terminals (when possible) is shown in Figure 1, including nodal support values from the Bayes and ML analyses. We also present four supplemental trees: Bayesian tree of highest posterior probability (Supplemental Figure 1), Bayesian consensus tree (Supplemental Figure 2), tree of highest log-likelihood (Supplemental Figure 3), and ML bootstrap consensus tree (Supplemental Figure 4).


PHYLOGENETIC CONSIDERATIONS

The phylogenetic tree in Figure 1 does not have well-supported basal branches. Its primary purpose was as a source of evidence to delineate species. Nonetheless, several lineages are worth noting. *Alabagrus fernandodiasi*, *A. jackiemillerae*, and *A. jeanmariecadioui* form a clade, and all members are similar in color and morphology (BIPP=81, MLBS<50). None are represented by males, which is unusual considering each is represented by 24 or more specimens. It may represent a parthenogenetic clade. Three species, *Alabagrus donlafontainei*, *A. cuna*, and *A. scottmilleri*, are distinctly sexually dichromatic, and all are found in a clade composed of five species (BIP=84, MLBS<50). One of the other species, *A. axelhausmanni*, is represented by a male only as the holotype, and the other member of the clade, *A. donnai*, have males that are darker (orange rather than yellow) than females (BIPP=99, MLBS=82). Overall it is common to have structural sexual dimorphism in members of *Alabagrus*. In particular, sculpture of the propodeum is often much more pronounced in males than females.

KEY TO COSTA RICAN ALABAGRUS


If you know the species of the host caterpillar it might be best to use the search function on electronic versions of this publication and find the species of *Alabagrus* that is (are) on the host, and check your specimen against the images and description. It is doubtful that this key will be very useful outside of Costa Rica and adjacent countries.

- 1A. Gena right angled or acute posteroventrally 2
- 1B. Gena rounded or with an obtuse angle posteroventrally 78



Couplet 1

2(1)A. Mesoscutum mostly or entirely melanic.....	3
2(1)B. Mesoscutum mostly or entirely pale	63




Couplet 2

3(2)A. Forewing almost entirely infusate, ignore small clear patches posteriad stigma	4
3(2)B. Forewing patterned with large clear, white, or yellow areas.....	18



Couplet 3

4(3)A. First tergum varying from weakly convex to with a rounded longitudinal bulge	5
4(3)B. First tergum with well-defined median longitudinal carina	14



Couplet 4

5(4)A. Propodeum melanic	6
5(4)B. Propodeum pale	8
5(4)C. Propodeum melanic anteriorly, pale posteriorly	<i>A. combos</i>




Couplet 5

- 6(5)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface.....7
- 6(5)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface*A. donharveyi* n. sp.




Couplet 6

- 7(6)A. Median tergite 1 narrower*A. leedyeri* n. sp.
- 7(6)B. Median tergite 1 wider.....*A. johnsharkeyi* n. sp.



Couplet 7

- 8(5)A. Median tergite 3 pale 9
- 8(5)B. Median tergite 3 pale anteriorly, melanic posteriorly.....*A. watsoni*




Couplet 8

- 9(8)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface.....10
- 9(8)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface 12




Couplet 9

- 10(9)A. Hind coxa entirely pale.....11
- 10(9)B. Hind coxa pale medially with a large melanic patch or vertical stripe laterally*A. sarapiqui*




Couplet 10

- 11(10)A. Body black and yellow *A. donnai*
- 11(10)B. Body black and red to black and orange*A. jimmilleri* n. sp.




Couplet 11

- 12(9)A. Penultimate metasomal tergum (often more terga) melanic*A. combos*
- 12(9)B. Penultimate metasomal tergum pale, reddish orange to yellow 13




Couplet 12

- 13(12)A. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly; pale parts of body yellow..... *A. donnai*
- 13(12)B. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron; pale parts of body orange.....*A. arawak*




Couplet 13

- 14(4)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface 15
- 14(4)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface*A. axelhausmanni* n. sp.




Couplet 14

- 15(14)A. Hind femur mostly or entirely pale *A. bernardoespinozai* n. sp.
- 15(14)B. Hind femur mostly or entirely melanic 16




Couplet 15

- 16(15)A. Median tergite 1 wider than long. *A. miqa*
- 16(15)B. Median tergite 1 longer than wide.....17




Couplet 16

- 17(16,31,32)A. Median tergite 2 black*A. bobrobbinsi* n. sp.
- 17(16,31,32)B. Median tergite 2 red/orange.....*A. karensSharkeyae* n. sp.



Couplet 17

- 18(3)A. Propodeum areolate, with at least one closed areola.....19
- 18(3)B. Propodeum lacking complete areolae, usually mostly or completely smooth 41




Couplet 18

- 19(18)A. Head entirely melanic 20
- 19(18)B. Head with at least some traces of pale color.....*A. kaydodgeae* n. sp.



Couplet 19

- 20(19)A. Penultimate metasomal tergum (or more) melanic.....21
- 20(19)B. Penultimate metasomal tergum pale, reddish orange to yellow*A. paulSharkeyi* n. sp.




Couplet 20

- 21(20)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface.....22
- 21(20)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface 38




Couplet 21

- 22(21)A. Precoxal sulcus with one or several foveae posteroventrally, with or without a smooth groove extending anteriorly.....23
- 22(21)B. Precoxal sulcus longer, distinct and foveolate ½ or more length of mesopleuron 33



Couplet 22

- 23(22)A. Metapleuron mostly or entirely melanic 24
- 23(22)B. Metapleuron mostly or entirely pale 30




Couplet 23

- 24(23)A. Propodeum melanic 25
- 24(23)B. Propodeum pale 28
- 24(23)C. Propodeum melanic anteriorly, pale posteriorly*A. almasolisae* n. sp.




Couplet 24

25(24)A. First tergum varying from weakly convex to with a rounded longitudinal bulge	26
25(24)B. First tergum with well-defined median longitudinal carina	27



Couplet 25

26(25)A. Hind femur mostly or entirely pale	<i>A. almasolisae</i> n. sp. males
26(25)B. Hind femur mostly or entirely melanic	<i>A. leedyeri</i> n. sp.




Couplet 26

27(25)A. Median tergite 1 narrower with a stronger median ridge	<i>A. bobpoolei</i> n. sp. males
27(25)B. Median tergite wider with a weak longitudinal ridge or anteromedial bulge	<i>A. almasolisae</i> n. sp. males




Couplet 27

- 28(24)A. Hind coxa entirely pale; third tergum with deep transverse depression.....*A. tommyersi* n. sp.
- 28(24)B. Hind coxa mostly pale with a vertical melanic stripe; third tergum lacking transverse depression or depression barely indicated ...29




Couplet 28

- 29(28)A. Forewing with two distinct pale areas, one basally and one below stigma*A. almasolisae* n. sp. males
- 29(28)B. Forewing lacking two distinct pale areas; base of the wing pale with some melanic color behind parastigma..*A. marcepssteini* n. sp.




Couplet 29

- 30(23)A. Third tergum lacking transverse depression or depression barely indicated.....31
- 30(23)B. Third tergum with deep transverse depression.....32




Couplet 30

- 31(30)A. Forewing banded from base: yellow, melanic, yellow, melanic.....*A. genemonroei* n. sp.
- 31(30)B. Forewing mostly or entirely infusate often with pale patches that do not form discrete bands.....17



Couplet 31

32(30)A. First tergum varying from weakly convex to with a rounded longitudinal bulge	<i>A. tommyersi</i> n. sp.
32(30)B. First tergum with well-defined median longitudinal carina	17



Couplet 32

33(22)A. Median tergite 3 pale	34
33(22)B. Median tergite 3 pale anteriorly, melanic posteriorly	36



Couplet 33

34(33)A. Median tergite 1 narrower with a stronger median ridge	35
34(33)B. Median tergite wider with a weak longitudinal ridge or anteromedial bulge	<i>A. andywarreni</i> n. sp. males



Couplet 34

35(34)A. Third tergum lacking transverse depression or depression barely indicated	<i>A. genemonroei</i> n. sp.
35(34)B. Third tergum with deep transverse depression	<i>A. janzeni</i> males



Couplet 35

- 36(33)A. First tergum with well-defined median longitudinal carina 37
- 36(33)B. First tergum varying from weakly convex to with a rounded longitudinal bulge*A. lindapitkinae* n. sp.




Couplet 36

- 37(36)A. Propodeum mostly or entirely pale.....*A. janzeni* males
- 37(36)B. Propodeum entirely melanic*A. bobpoolei* n. sp. males




Couplet 37

- 38(21)A. Median tergite 1 pale; median tergite 2 pale 39
- 38(21)B. Median tergite 1 partly melanic but pale at base; median tergite 2 melanic*A. donharveyi* n. sp.




Couplet 38

- 39(38)A. Third tergum lacking transverse depression or depression barely indicated.....40
- 39(38)B. Third tergum with deep transverse depression.....*A. janzeni* males




Couplet 39

40(39)A. Hind femur entirely pale.....	<i>A. andywarreni</i> n. sp. males
40(39)B. Hind femur melanic dorsally, pale ventrally	<i>A. paulgoldsteini</i> n. sp.
40(39)C. Hind femur entirely melanic, sometimes with a hint of reddish color ventrally but much darker than <i>A. paulgoldsteini</i>	<i>A. ramyamanjunathae</i> n. sp.




Couplet 40

41(18)A. Forefemur entirely melanic or only with a spot of pale color apically	42
41(18)B. Forefemur entirely pale	57
41(18)C. Forefemur melanic bicolored, pale and melanic	61



Couplet 41

42(41)A. Forewing with two distinct yellow bands, one basal to and the other apical to stigma	43
42(41)B. Forewing with two yellow bands, one distinct basal band and a weaker triangular band directly below (posterior to) stigma	46
42(41)C. Forewing lacking two distinct bands; pale basally to near level of apex of stigma with a small darkening near parastigma	50
42(41)D. Forewing with two distinct yellow bands, one basal to and the other directly below (posterior to) stigma	54




Couplet 42

43(42)A. Third tergum lacking transverse depression or depression barely indicated.....	44
43(42)B. Third tergum with deep transverse depression.....	45




Couplet 43

- 44(43)A. Median tergite 3 partly or entirely melanic*A. bobpoolei* n. sp. females
- 44(43)B. Median tergite 3 entirely pale*A. andywarreni* n. sp. females




Couplet 44

- 45(43)A. Metapleuron and propodeum both entirely black*A. bobpoolei* n. sp. females
- 45(43)B. Metapleuron and propodeum from mostly black to entirely yellow, one or the other always with some yellow/orange color
.....*A. janzeni* males




Couplet 45

- 46(42)A. First tergum varying from weakly convex to with a rounded longitudinal bulge 47
- 46(42)B. First tergum with well-defined median longitudinal carina 49




Couplet 46

- 47(46)A. Hind coxa entirely pale 48
- 47(46)B. Hind coxa mostly pale with a melanic patch laterally or a vertical melanic stripe*A. almasolisae* n. sp. females




Couplet 47

- 48(47)A. Third tergum lacking transverse depression or depression barely indicated; metapleuron entirely pale*A. iankitchingi* n. sp.
- 48(47)B. Third tergum with deep transverse depression; metapleuron partly or entirely melanic.....*A. tommyersi* n. sp.




Couplet 48

- 49(46)A. Metapleuron and propodeum both entirely black*A. bobpoolei* n. sp. females
- 49(46)B. Metapleuron and propodeum from mostly black to entirely yellow, one or the other always with some yellow color
.....*A. janzeni* males




Couplet 49

- 50(42)A. First tergum varying from weakly convex to with a rounded longitudinal bulge 51
- 50(42)B. First tergum with well-defined median longitudinal carina 53




Couplet 50

- 51(50)A. Hind coxa entirely pale 52
- 51(50)B. Hind coxa mostly pale with a vertical melanic stripe*A. marcepsteini* n. sp.




Couplet 51

- 52(51)A. Tegula black.....*A. malcolmscoblei* n. sp.
- 52(51)B. Tegula pale, yellow to orange*A. nickgrishini* n. sp.




Couplet 52

- 53(50)A. Metapleuron and propodeum both entirely black*A. bobpoolei* n. sp. females
- 53(50)B. Metapleuron and propodeum from mostly black to entirely yellow, one or the other always with some yellow color
.....*A. janzeni* males




Couplet 53

- 54(42)A. Third tergum lacking transverse depression or depression barely indicated.....55
- 54(42)B. Third tergum with deep transverse depression.....56




Couplet 54

- 55(54)A. Metapleuron mostly or entirely melanic; propodeum melanic*A. bobpoolei* n. sp. females
- 55(54)B. Metapleuron mostly or entirely pale; propodeum pale*A. genemonroei* n. sp.




Couplet 55

- 56(54)A. Metapleuron and propodeum both entirely black*A. bobpoolei* n. sp. females
- 56(54)B. Metapleuron and propodeum from mostly black to entirely yellow, one or the other always with some yellow color
.....*A. janzeni* males



Couplet 56

- 57(41)A. First tergum varying from weakly convex to with a rounded longitudinal bulge 58
- 57(41)B. First tergum with well-defined median longitudinal carina*A. janzeni*




Couplet 57

- 58(57)A. Metapleuron mostly or entirely pale; hind coxa partly or entirely melanic*A. longinoi* n. sp.
- 58(57)B. Metapleuron mostly or entirely pale; hind coxa entirely pale 59
- 58(57)C. Metapleuron mostly or entirely melanic; hind coxa partly or entirely melanic*A. jeanmariecadioui* n. sp.




Couplet 58

- 59(58)A. Femur of midleg entirely yellow 60
- 59(58)B. Femur of midleg mostly yellow but melanic at midlength, sometimes only ventrally*A. fernandodiasi* n. sp.




Couplet 59

- 60(59)A. Midtibia with 0–3 nonapical spines, precoxal sulcus shorter with fewer crenulae*A. fernandezi* n. sp.
- 60(59)B. Midtibia with 5 or more nonapical spines; precoxal sulcus longer with more crenulae*A. andresfreitasi* n. sp.




Couplet 60

- 61(41)A. First tergum varying from weakly convex to with a rounded longitudinal bulge 62
- 61(41)B. First tergum with well-defined median longitudinal carina*A. janzeni* males



Couplet 61

- 62(61)A. Median tergite 3 pale*A. scottshawi* n. sp.
- 62(61)B. Median tergite 3 pale anteriorly, melanic posteriorly.....*A. jackiemillerae* n. sp.



Couplet 62

- 63(2)A. Midfemur melanic.....64
- 63(2)B. Midfemur pale.....71
- 63(2)C. Midfemur bicolored, melanic and pale.....77



Couplet 63

- 64(63)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface.....65
- 64(63)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface 69



Couplet 64

- 65(64)A. Penultimate metasomal tergum (and often more terga) melanic 66
- 65(64)B. Penultimate metasomal tergum pale, reddish orange to yellow 67




Couplet 65

- 66(65)A. Propodeum areolate, with at least one closed areola; first tergum with well-defined median longitudinal carina.....*A. cuna* male
- 66(65)B. Propodeum lacking complete areolae, smooth; first tergum weakly convex to with a rounded longitudinal bulge*A. varius*



Couplet 66


- 67(65)A. Hind coxa entirely pale.....*A. yuanmaofangi* n. sp.
- 67(65)B. Hind coxa with a black patch laterally (may only be present near apex) 68



Couplet 67


68(67) These three species are very difficult to separate and this couplet has limited accuracy.

- 68(67)A. Forewing distinctly yellow in basal $\frac{2}{3}$ but lacking a distinct margin between pale and dark areas. *A. johnbrowni* n. sp.
- 68(67)B. Forewing distinctly yellow in basal $\frac{2}{3}$ and with a distinct margin between pale and dark areas.....*A. cocto*
- 68(67)C. Forewing with only a hint of yellow basally.....*A. mariaheikkilae* n. sp.




Couplet 68

- 69(64)A. Hind coxa entirely pale*A. arawak*
- 69(64)B. Hind coxa with a black patch laterally (may only be present near apex) 70



Couplet 69

- 70(69)A. Forewing with a well-defined margin between apical infusate area and basal yellow area*A. brendameierottoae* n. sp.
- 70(69)B. Forewing lacking a well-defined margin between apical infusate area and basal pale area.....*A. paulheberti* n. sp.




Couplet 70

- 71(63)A. Penultimate metasomal tergum melanic 72
- 71(63)B. Penultimate metasomal tergum pale, reddish orange to yellow 74




Couplet 71

- 72(71)A. Median tergites 1 and 2 mostly or entirely melanic *A. scottmilleri* n. sp. male
- 72(71)B. Median tergites 1 and 2 pale 73




Couplet 72

- 73(72)A. Propodeum areolate, with at least one closed areola.....*A. kaydodgeae* n. sp.
- 73(72)B. Propodeum lacking complete areolae, mostly or completely smooth*A. varius*




Couplet 73

- 74(71)A. Hind coxa entirely pale.....75
- 74(71)B. Hind coxa with a black patch laterally or near apex.....76




Couplet 74

- 75(74)A. Forewing yellow in basal ⅓, melanic apically *A. cuna* female
- 75(74)B. Forewing mostly melanic with a clear vertical band near stigma.....*A. donlafontainei* female




Couplet 75

- 76(73)A. Forewing banded from base: yellow, infusate, yellow, infusate.....*A. scottmilleri* n. sp. female
- 76(73)B. Forewing yellow basally, infusate apically.....*A. cara*




Couplet 76

- 77(63)A. Propodeum lacking complete areolae, mostly or completely smooth*A. varius*
- 77(63)B. Propodeum areolate, with at least one closed areola.....*A. donlafontainei* n. sp. male




Couplet 77

- 78(1)A. Mesoscutum entirely melanic.....79
- 78(1)B. Mesoscutum entirely pale.....111
- 78(1)C. Mesoscutum mostly pale with some melanic infusions 142
- 78(1)D. Mesoscutum mostly melanic with some pale infusions.....144




Couplet 78

- 79(78)A. Median tergite 3 melanic 80
- 79(78)B. Median tergite 3 pale 83
- 79(78)C. Median tergite 3 pale anteriorly, melanic posteriorly*A. yuchinkengae* n. sp.




Couplet 79

80(79)A. Metapleuron mostly or entirely melanic	81
80(79)B. Metapleuron mostly or entirely pale	<i>A. semihesperheidei</i> n. sp.




Couplet 80

81(80)A. Median tergite 1 entirely melanic	82
81(80)B. Median tergite 1 partly or completely pale (reddish)	<i>A. hespenheidei</i> n. sp.



Couplet 81

82(81)A. Forewing almost entirely infusate, ignore small clear patches posteriad stigma	<i>A. ekchuah</i>
82(81)B. Forewing patterned with a clear or white area apically	<i>A. tanyadapkeyae</i> n. sp.




Couplet 82

83(79)A. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron	96
83(79)B. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly	84
83(79)C. Precoxal sulcus absent or mostly absent, represented at most by small, shallow depression, usually posteriorly, crenulae always lacking	105




Couplet 83

84(83)A. First tergum varying from weakly convex to with a rounded longitudinal bulge.....	85
84(83)B. First tergum with well-defined median longitudinal carina	92
84(83)C. First tergum with two converging longitudinal carinae	95




Couplet 84

85(84)A. Occiput dorsolaterally rounded; clypeus unmodified, lacking protuberances.....	86
85(84)B. Occiput dorsolaterally with squared protuberances; clypeus with lateral protuberances	<i>A. victoriapookae</i> n. sp.




Couplet 85

86(85)A. Forewing almost entirely infusate, ignore small clear patches posteriad stigma	87
86(85)B. Forewing patterned with large clear, white, or yellow areas.....	90




Couplet 86

87(86)A. Propodeum smooth.....	89
87(86)B. Propodeum sculptured	88




Couplet 87

- 88(87)A. Body color black and yellow.....*A. johnburnsi* n. sp.
- 88(87)B. Body color black and orange to black and red*A. markmetzi* n. sp.




Couplet 88

- 89(87)A. Propodeum areolate, with several closed areolae.....*A. markmetzi* n. sp.
- 89(87)B. Propodeum lacking, complete areolae, mostly or completely smooth*A. johnobryckii* n. sp.




Couplet 89

- 90(86)A. Propodeum areolate, with at least one closed areola.....*A. markmetzi* n. sp.
- 90(86)B. Propodeum lacking complete areolae, usually mostly or completely smooth 91




Couplet 90

- 91(90)A. Hind coxa entirely or almost entirely pale; hind femur mostly or entirely melanic.....*A. englishi*
- 91(90)B. Hind coxa pale with a large melanic patch laterally; hind femur mostly or entirely pale*A. johnburnsi* n. sp.




Couplet 91

- 92(84)A. First median tergite longer than wide; third tergum lacking transverse depression or depression barely indicated.....93
- 92(84)B. First median tergite wider than long; third tergum with deep transverse depression *A. miqa*




Couplet 92

- 93(92)A. Propodeum with more defined and complete sculpture; first median tergite mostly orange.....94
- 93(92)B. Propodeum with less defined and complete sculpture; first median tergite yellow.....*A. johnburnsi* n. sp.




Couplet 93

- 94(93)A. Propodeum and metapleuron entirely pale*A. sarahmeierottoae* n. sp. male
- 94(93)B. Propodeum and metapleuron with melanic color.....*A. johnobryckii* n. sp.




Couplet 94

- 95(84)A. Propodeum smooth, melanic*A. barbsharanowskiae* n. sp. female
- 95(84)B. Propodeum sculptured, pale.....*A. kaciejoe* n. sp.




Couplet 95

- 96(83)A. Propodeum melanic 97
- 96(83)B. Propodeum pale 99




Couplet 96

97(95)A. Metasoma mostly or entirely yellow	<i>A. keithwillmotti</i> n. sp.
97(95)B. Metasoma mostly or entirely orange	98




Couplet 97

98(97)A. First tergum with a single rounded longitudinal bulge	<i>A. brianharrisi</i> n. sp.
98(97)B. First tergum with two converging longitudinal carinae	<i>A. kaciejoae</i> n. sp. male




Couplet 98

99(96)A. First tergum varying from weakly convex to with a rounded longitudinal bulge	100
99(96)B. First tergum with well-defined median longitudinal carina	104




Couplet 99

100(99)A. Hind femur mostly or entirely pale	101
100(99)B. Hind femur mostly or entirely melanic	103




Couplet 100

101(100)A. Hind femur narrower	102
101(100)B. Hind femur wider	<i>A. nicoya</i>



Couplet 101

102(101)A. Forewing with yellowish or clear area extending to the second submarginal cell	<i>A. roibasi</i>
102(101)B. Forewing entirely infusate or, if with yellowish or clear area basally, it does not extend to second submarginal cell (we cannot distinguish the following three species morphologically).....	
.....	<i>A. jennyphillipsae</i> n. sp., <i>A. isidrochaconi</i> n. sp., <i>A. jeanfrancoislandryi</i> n. sp.




Couplet 102

103(100)A. Protuberance present lateral to clypeus; gena not elongate	<i>A. victoriapookae</i> n. sp.
103(100)B. Protuberance absent lateral to clypeus; gena elongate	<i>A. arua</i>
103(100)C. Protuberance absent lateral to clypeus; gena not elongate (the common state for <i>Alabagrus</i> species)	
.....	<i>A. jeanfrancoislandryi</i> n. sp. males



Couplet 103

- 104(99)A. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface.....
.....*A. sarahmeierottoae* n. sp. male
- 104(99)B. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface
.....*A. paulthiaucourti* n. sp.




Couplet 104

- 105(82)A. Penultimate metasomal tergum (or more) melanic.....106
- 105(82)B. Penultimate metasomal tergum pale, reddish orange to yellow 107




Couplet 105

- 106(105)A. Metapleuron mostly or entirely melanic; propodeum melanic*A. stiremani* n. sp.
- 106(105)B. Metapleuron mostly or entirely pale; propodeum pale*A. quickei* n. sp.




Couplet 106

- 107(104)A. Pronotum entirely melanic.....108
- 107(104)B. Pronotum partly or completely pale*A. sarahsharkeyae* n. sp.




Couplet 107

- 108(107)A. Third tergum lacking transverse depression or depression barely indicated; first tergum varying from weakly convex to with a rounded longitudinal bulge 109
- 108(107)B. Third tergum with deep transverse depression; first tergum with well-defined median longitudinal carina.....*A. miqa*




Couplet 108

- 109(108)A. Propodeum sculptured, minimally with two diverging carinae anteriorly*A. sarahmeierottoae* n. sp. female
- 109(108)B. Propodeum completely smooth or with a slight bulge anteromedially 110



Couplet 109

- 110(109)A. Body melanic and orange.....*A. englishi*
- 110(109)B. Body melanic and yellow*A. johnburnsi* n. sp.



Couplet 110

111(78)A. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron 125

111(78)B. Precoxal sulcus with one or several distinct foveae posteroventrally with or without a smooth groove extending anteriorly, fovea not extending ½ length of mesopleuron 112


111(78)C. Precoxal sulcus absent or mostly absent, represented at most by small, shallow depression, usually posteriorly, crenulae always lacking.....133



Couplet 111

112(110)A. Propodeum areolate, with at least one closed areola.....113


112(110)B. Propodeum lacking complete areolae, mostly smooth 117



Couplet 112


113(112)A. Forewing with 2 distinct yellow bands*A. voto*

113(112)B. Forewing lacking 2 distinct yellow bands 114




Couplet 113

114(113)A. Head entirely melanic	115
114(113)B. Head partly or entirely pale	<i>A. maculipes</i>




Couplet 114

115(114)A. Males	116
115(114)B. Females (see Figure 62)	<i>A. patsharkeyi</i> n. sp.
116(115)A. Body orange and melanic; hind coxa entirely pale	<i>A. derailersi</i> male
116(115)B. Body yellow and melanic; hind coxa partly melanic	<i>A. masneri</i> male




Couplet 116

117(112)A. Ovipositor distinctly longer than body	120
117(112)B. Ovipositor longer than metasoma but not distinctly longer than body	118



Couplet 117

- 118(117)A. Propodeum sculptured *A. reddypallii* n. sp.
- 118(117)B. Propodeum smooth, at most with a smooth bulge anteromedially 119




Couplet 118

- 119(118)A. Fore- and midcoxae entirely pale *A. derailersi* female
- 119(118)B. Fore- and midcoxae both partly or entirely melanic *A. masneri* female



Couplet 119

- 120(117)A. Forewing almost entirely infusate 121
- 120(117)B. Forewing patterned with two yellow stripes *A. varius*




Couplet 120

- 121(120)A. Fore- and midcoxae entirely pale 122
- 121(120)B. Fore- and midcoxae both partly or entirely melanic 123




Couplet 121

- 122(121)A. Propodeum with irregular sculpture anteromedially*A. mattottoi* n. sp.
- 122(121)B. Propodeum entirely smooth except for a weak, rounded median longitudinal ridge*A. craigevansi* n. sp.




Couplet 122

- 123(121)A. Longer specimens, more than 7 mm, and ovipositor longer, clearly longer than body.....124
- 123(121)B. Shorter specimens, less than 6 mm, and ovipositor relatively shorter, about as long as body.....*A. masneri* female



Couplet 123


- 124(123)A. Hind coxa with a melanic patch laterally; AA. Propodeum entirely smooth except for a weak, rounded median longitudinal ridge*A. craigevansi* n. sp.
- 124(123)B. Hind coxa entirely pale or almost entirely pale with melanic color restricted to apex; BB. Propodeum with more sculpture*A. patsharkeyi* n. sp.



Couplet 124

125(111)A. Head entirely melanic 126


125(111)B. Head partly or entirely pale*A. maculipes*



Couplet 125

126(125)A. Forewing almost entirely infusate, ignore small clear patches posteriad stigma 127


126(125)B. Forewing patterned with large clear, white, or yellow areas.....131



Couplet 126

127(126)A. Propodeum areolate, with at least one closed areola.....128

127(126)B. Propodeum lacking complete areolae, usually mostly or completely smooth 130




Couplet 127

- 128(127)A. Gena longer.....*A. arua*
- 128(127)B. Gena shorter (normal condition) 129




Couplet 128

- 129(128)A. Females (see Figure 62).....*A. patsharkeyi* n. sp.
- 129(128)B. Males (see Figure 57)*A. masneri* male
- 130(127)A. Ovipositor less than 2× length of metasoma*A. pecki*
- 130(127)B. Ovipositor 2× or more length of metasoma*A. patsharkeyi* n. sp.



Couplet 130

- 131(126)A. First tergum varying from weakly convex to with a rounded longitudinal bulge 132
- 131(126)B. First tergum with well-defined median longitudinal carina *A. stigma*




Couplet 131

- 132(131)A. Gena longer.....*A. arua*
- 132(131)B. Gena shorter (normal condition)*A. ilgookangi* n. sp.



Couplet 132

- 133(111)A. Propleuron melanic 134
- 133(111)B. Propleuron pale 138




Couplet 133

- 134(133)A. Penultimate metasomal tergum (or more) melanic.....*A. brianbrowni* n. sp.
- 134(133)B. Penultimate metasomal tergum pale, reddish orange to yellow 135




Couplet 134

- 135(134)A. Pronotum entirely pale.....136
- 135(134)B. Pronotum melanic anteriorly, pale posteriorly 137



Couplet 135

- 136(135)A. Ovipositor less than 2× longer than metasoma*A. masneri* female
- 136(135)B. Ovipositor more than 2× longer than metasoma.....*A. sarahsharkeyae* n. sp.




Couplet 136

- 137(135)A. Hind femur mostly or entirely melanic*A. rudolfmeieri* n. sp.
- 137(135)B. Hind femur mostly or entirely pale*A. sarahsharkeyae* n. sp.




Couplet 137

- 138(133)A. Forewing almost entirely infusate, ignore small clear patches posteriad stigma 139
- 138(133)B. Forewing patterned with large clear, white, or yellow areas.....140




Couplet 138

- 139(138)A. Males (see Figure 67)*A. pecki*
- 139(138)B. Females (see Figure 57)..... *A. masneri*
- 140(138)A. Fore- and midfemora mostly or entirely melanic.....141
- 140(138)B. Fore- and midfemora mostly or entirely pale*A. bobwhartoni*



Couplet 140

- 141(140)A. Margins of mesopleuron without a greenish tinge, concolorous with remainder of pleuron; gena smoothly rounded
.....*A. pecki* female
- 141(140)B. Margins of mesopleuron with a greenish tinge; gena more angulate *A. hansonii*



Couplet 141

- 142(78)A. Gena longer.....*A. arua*
- 142(78)B. Gena shorter (normal condition) 143



Couplet 142

- 143(142)A. Penultimate metasomal tergum (or more) melanic.....*A. brianbrowni* n. sp.
- 143(142)B. Penultimate metasomal tergum pale, reddish orange to yellow*A. sarahsharkeyae* n. sp.




Couplet 143

144(78)A. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron	145
144(78)B. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly.....	<i>A. sarahmeierottoae</i> n. sp. male
144(78)C. Precoxal sulcus absent or mostly absent, represented at most by small, shallow depression, usually posteriorly, crenulae always lacking.....	146




Couplet 144

145(144)A. First tergum varying from weakly convex to with a rounded longitudinal bulge	<i>A. arua</i>
145(144)B. First tergum with well-defined median longitudinal carina	<i>A. sarahmeierottoae</i> n. sp. male




Couplet 145

146(144)A. Forewing mostly or entirely infusate, lacking basal pale area	<i>A. sarahmeierottoae</i> n. sp. female
146(144)B. Forewing pale (clear or yellow) basally, infusate apically	147



Couplet 146

- 147(146)A. Propodeum mostly or entirely melanic; first median tergite yellow.....*A. johnburnsi* n. sp.
- 147(146)B. Propodeum mostly or entirely pale; first median tergite orange*A. sarahmeierottoae* n. sp. female



Couplet 147

SYSTEMATICS

Alabagrus almasolisae Sharkey n. sp.

Figure 2

DIAGNOSIS. Gena acute posteroventrally. Forewing with two distinct yellow bands, one basal and the other directly below (posterior to) stigma, the latter incomplete posteriorly. Metapleuron completely melanic or with posterior and dorsal margins pale. Propodeum completely smooth in female, completely areolate in male. First median tergite with a short anteromedial, rounded, longitudinal bulge.

DESCRIPTION. Body length 9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present,

restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum with more sculpture, always with at least one closed areola; some males with a weak, short, median, longitudinal carina on first median tergite.

Variation. Propodeum from completely smooth to mostly smooth with a weak transverse ridge; propodeum almost all black to half pale in posterior half—posterior to transverse ridge.

HOST INFORMATION. *Neea psychotrioides*.

ETYMOLOGY. Named in honor of Dr. Alma Solis of the Systematic Entomology Laboratory in the U.S. National Museum of




Figure 2 *Alabagrus almasolisae* n. sp., **A–G** holotype; **H** paratype male: **A.** lateral habitus, **B.** dorsal head, **C.** anterior head, **D.** dorsal habitus, **E.** lateral head, **F.** wings, **G.** propodeum and tergum 1, **H.** propodeum and tergum 1.

Natural History, the taxonomist who has, along with Dr. Eugene Munroe (deceased), diligently carried out the taxonomy of the naming and identification of the *Alabagrus* host species of Crambidae and supported the ACG biodiversity inventory in many other ways.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Neea psychotrioides* on *Ceratocilia sixolalis*, 10.94076°N, 85.3177°W, 461 m, 17.iii–8.iv.2011, DHJPAR0042817 (EMUS). PARATYPES: hosts all from *Neea psychotrioides* on *Ceratocilia sixolalis*. 2♀, 1♂, 10.9163°N, 85.37869°W, 460 m, 27.ix–20.x.2008, DHJPAR0028269, DHJPAR0028270, DHJPAR0028271. 2♀, 27.ix–21.x.2008, DHJPAR0028272, DHJPAR0028273. 1♀, 27.ix–22.x.2008, DHJPAR0030384. 1♂, 10.9305°N, 85.37223°W, 527 m, 6–30.xi.2009, DHJPAR0037924. 2♀, 1♂, 6.xi–2.xii.2009, DHJPAR0037932, DHJPAR0037933, DHJPAR0037935. 1♂, 6.xi–3.xii.2009, DHJPAR0037931. 1♂, 10.89666°N, 85.29003°W, 400 m, 1–19.x.2010, DHJPAR0041205. 1♀, 27.x–14.xi.2010, DHJPAR0041578. 1♀, 27.x–16.xi.2010, DHJPAR0041171. 1♀, 1♂, same data as holotype, DHJPAR0042816, DHJPAR0042818. 1♂, 10.9301°N, 85.25205°W, 109 m, 26.viii–14.ix.2013, DHJPAR0053625. 1♀, 10.94076°N, 85.3177°W, 461 m, 4–30.x.2013, DHJPAR0054493. 1♂, 10.96187°N, 85.28045°W, 96 m, 28.xii.2013–17.i.2014, DHJPAR0054507. 1♀, 28.xii.2013–20.i.2014, DHJPAR0054508. 1♀, 28.xii.2013–21.i.2014, DHJPAR0054510. 1♀, 10.93010°N, 85.25205°W, 109 m, 22.i–10.ii.2014, DHJPAR0055101. 1♀, 3♂, 10.94741°N, 85.31501°W, 491 m, 26.xii.2014–21.i.2015, DHJPAR0057420, DHJPAR0057422, DHJPAR0057421, DHJPAR0057423. 1♂, 10.97073°N, 85.31434°W, 326 m, 11–30.xii.2015, DHJPAR0058545. 1♀, 11–31.xii.2015, DHJPAR0058544. 1♀, 11.xii.2015–2.i.2016, DHJPAR0058540. ♀, 11.xii.2015–3.i.2016, DHJPAR0058543 (EMUS, HIC).

Alabagrus andresfreitasi Sharkey n. sp.

Figure 3

DIAGNOSIS. Gena acute posteroventrally. Forefemur pale. Mid-femur entirely yellow. Hind coxa entirely pale. Mesoscutum black. Metapleuron entirely pale. Propodeum completely smooth. First tergum varying from weakly convex to with a rounded longitudinal bulge. Midtibia with five or more nonapical spines; precoxal sulcus crenulate ½ length of mesopleuron.

DESCRIPTION. Body length 10 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Similar to female.

HOST INFORMATION. *Phostria truncatalis*.

ETYMOLOGY. Named in honor of Dr. Andres Freitas of Universidade Estadual de Campinas, Brazil, who has assisted heavily with the taxonomy and natural history of the ACG Nymphalidae inventory and introduced D.J. and W.H. to Brazilian lepidopterology.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phostria truncatalis* on *Alibertia edulis*, 11.03004°N, 85.52699°W, 280 m, 23.ix–18.x.2009, DHJPAR0036714 (EMUS). PARATYPES: all from *Phostria truncatalis* on *Alibertia edulis*. 1♀, 23.ix–26.x.2009, DHJPAR0037959.




Figure 3 *Alabagrus andresfreitasi* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

1♀, 23.ix–1.xi.2009, DHJPAR0037880. 1♀, 1♂, 12.ix–5.x.2004, DHJPAR0015362, DHJPAR0015364. 1♂, 12.ix–19.x.2004, DHJPAR0015365. 1♀, 12.ix–22.x.2004, DHJPAR0015366. 1♂, 28.ix–27.x.2009, DHJPAR0037885. 1♀, 10.77074°N, 85.42874°W, 365 m, 13.viii–19.ix.2012, DHJPAR0050365. 1♂, 10.7709°N, 85.42455°W, 375 m, 4.xii.2013–4.i.2014, DHJPAR0054471. 1♂, 11.03226°N, 85.52776°W, 290 m, 18.ix–16.x.2004, DHJPAR0015352 (EMUS, HIC).

Alabagrus andywarreni Sharkey n. sp.

Figure 4

DIAGNOSIS. Gena acute posteroventrally. Precoxal sulcus crenulate 1/2 or more length of mesopleuron. Forefemur melanic. Distal yellow band in forewing not reaching posterior margin. Propodeum melanic, completely smooth in females, areolate in males. First median tergite as wide posteriorly as long with a weak rounded anteromedial longitudinal ridge. Median tergite 3 yellow.

NOTES. This species has two hosts, its sister species, *A. andresfreitasi*, shares one of the same hosts. The host data suggest that they may be the same species, but *COI* and color data suggest that they are separate species.

DESCRIPTION. Body length 10 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge, or with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.




Figure 4 *Alabagrus andywarreni* n. sp., holotype: A. lateral habitus, B. dorsal head, C. anterior head, D. dorsal habitus, E. propodeum and tergum 1, F. wings, G. lateral head and mesosoma.

Males. Some propodea with more sculpture; some hind femora with weak rugose sculpture ventrally.

HOST INFORMATION. *Phostria metalobalis*, *Phostria truncatalis*.

ETYMOLOGY. Named in honor of Dr. Andy Warren of the McGuire Center for Lepidoptera and Biodiversity, University of Florida, Gainesville, Florida, USA, who has both straightened out the ACG Hesperidae higher taxonomy and aided with ACG Hesperinae identifications.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phostria metalobalis* on *Genipa americana*, 11.03226°N, 85.52776°W, 290 m, 7–29.vii.2011, DHJPAR0044988 (EMUS). PARATYPES: The following are from *Phostria metalobalis* on *Genipa Americana*. 1 ♀, 10.7416°N, 85.42734°W, 420 m, 20.ix–16.x.2010, DHJPAR0041199. 1 ♂, 10.83764°N, 85.61871°W, 295 m, 18.vii–12.viii.2000, DHJPAR0015359. 1 ♂, 18.vii–19.viii.2000, DHJPAR0015360. 1 ♂, 18.vii–21.viii.2000, DHJPAR0015358. 1 ♂, 18.vii–22.viii.2000, DHJPAR0015361. 2 ♀, 10.84389°N, 85.61384°W, 300 m, 27.vi–7.viii.1980, DHJPAR0015474, 80-SRNP-244.2. 1 ♀, 10.85145°N, 85.60801°W, 290 m, 18.vi–21.vii.1981, DHJPAR0015473. 2 ♂, 10.85991°N, 85.65396°W, 340 m, 21.vii–13.viii.2000, DHJPAR0015356, DHJPAR0015357. 1 ♀, 21.vii–19.viii.2000, DHJPAR0015355. 1 ♀, 1 ♂, 22.vii–19.viii.2000, DHJPAR0015354, DHJPAR0015353. 1 ♂, 10.92291°N, 85.31877°W, 410 m, 19.vi–29.vii.2012, DHJPAR0049653. 1 ♂, 10.99616°N, 85.45562°W, 560 m, 17.vi–16.vii.2002, 02-SRNP-16081. 1 ♂, 11.01525°N, 85.39766°W, 415 m, 5.x–13.xi.2009, DHJPAR0037906. 1 ♀, 11.03004°N, 85.52699°W, 280 m, 17.vii–30.viii.2001, 01-SRNP-9963. 1 ♀, 11.03239°N, 85.57545°W, 220 m, 13.ix–9.x.2004, DHJPAR0015367. The following are from *Phostria truncatalis* on *Alibertia edulis*: 1 ♀, 10.94683°N, 85.56702°W, 275 m, 24.vi–14.vii.2000, DHJPAR0015472. 1 ♀, 24.vi–21.vii.2000, DHJPAR0015471. 1 ♂, 24.vi–23.vii.2000, 00-SRNP-15421. 1 ♀, 11.03004°N, 85.52699°W, 280 m, 12.ix–8.x.2004, DHJPAR0015363 (EMUS, HIC).

Alabagrus arawak Sharkey, 1988

Figure 5

Alabagrus arawak Sharkey, 1988:357–8. Holotype ♀, Ecuador (CNCI)

DIAGNOSIS. Gena acute posteroventrally. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Scutellum pale in females, melanic in males. Midfemur entirely or mostly melanic. Hind coxa entirely pale. Body of mesosoma entirely yellow in females, hind coxa either entirely yellow (females) or entirely melanic (males). Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron in males. Penultimate metasomal tergum pale, reddish orange to yellow.

NOTES. *Alabagrus warrau* does not occur in Costa Rica. Specimens formerly identified as such in Leathers and Sharkey (2003) belong in *A. arawak*. *Alabagrus arawak* and *A. warrau* may be synonyms, but morphologically they differ slightly.

DESCRIPTION. Body length 4.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.4\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Scutellum and metacoxa melanic; precoxal sulcus longer with more foveae.

DISTRIBUTION. Widespread from southern Mexico to northern Argentina; common year-round, and widespread in Costa Rica. The




Figure 5 *Alabagrus Arawak*, A–D female; F–H male: A. wings, B. lateral habitus, C. dorsal habitus, D. anterior head, E. anterior head, F. wings, G. propodeum and tergum 1, H. dorsal habitus.

holotype is from Ecuador (Sharkey, 1988). No reared specimens (numerous specimens in HIC, INBio, EMUS).

Alabagrus arua Sharkey, 1988

Figure 6

Alabagrus arua Sharkey, 1988:358–9. Holotype ♀, Ecuador (CNCI)

DIAGNOSIS. Gena with an obtuse angle posteroventrally. Head relatively more elongate than any other species of *Alabagrus*. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Pronotum melanic. Metapleuron mostly or entirely pale. First tergum weakly convex with a rounded longitudinal bulge. Median tergite 3 pale.

NOTES. *Alabagrus juchuy* may be a synonym, but contrary to *A. arua*, the type has a yellow stigma. The type of *A. juchuy* is from Trinidad. This may be the best character to separate them until COI data is obtained over a wider latitudinal range. The specimens referred to as *A. juchuy* in Leathers and Sharkey (2003) are all *A. arua*.

DESCRIPTION. Body length 6 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeal sculpture slightly stronger and often yellow; median ridge of median tergite 1 more acute.

HOST INFORMATION. *Hyalorista exuvialis* DHJ02.

DISTRIBUTION. The holotype is from Ecuador (Sharkey, 1988). Widespread from Costa Rica south to Amazonian Bolivia.

MATERIAL EXAMINED. Reared specimens: all from *Hyalorista exuvialis* DHJ02. 1♀, on *Hyptis capitata*, 10.94404°N, 85.31738°W, 455 m, 30.viii–13.ix.2010, DHJPAR0041194. 1♂, on *Hyptis obtusifolia*, 10.89666°N, 85.29003°W, 400 m, 6–23.v.2006, DHJPAR0015518. 1♀, 10.90661°N, 85.28784°W, 400 m, 17.v–1.vi.2007, DHJPAR0021202. 1♀, 10.88409°N, 85.25728°W, 460 m, 27.ii–17.iii.2009, DHJPAR0030487. 2♀, 10.9163°N, 85.37869°W, 460 m, 21.iv–13.v.2009, DHJPAR0035224, DHJPAR0035296. 1♂, 10.9305°N, 85.37223°W, 527 m, 13–29.v.2009, DHJPAR0035297. 1♀, 10.90528°N, 85.27882°W, 405 m, 6–23.vi.2012, DHJPAR0049054. 1♀, 10.89678°N, 85.27001°W, 420 m, 2–22.i.2013, DHJPAR0051379 (EMUS, HIC).

Alabagrus axelhausmanni Sharkey n. sp.

Figure 7

DIAGNOSIS. Female unknown. Male first median tergite with a sharp but short median longitudinal carina extending $\frac{1}{2}$ length of tergite. Propleuron pale or melanic and pale. Pronotum melanic.

DESCRIPTION. Body length 5.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1× longer than wide; with well-defined median




Figure 6 *Alabagrus arua*, female: A. lateral habitus, B. wings, C. dorsal habitus, D. anterior head, E. dorsal head, F. lateral head, G. lateral head and mesosoma.

longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated.

HOST INFORMATION. musoBioLep01 BioLep448.

ETYMOLOGY. Named in honor of Dr. Axel Hausmann of the Zoological State Collection Munich, Germany, who has both seriously

aided ACG inventory Geometridae identifications and inspired the introduction of DNA barcoding into the European taxasphere.

MATERIAL EXAMINED. HOLOTYPE ♂, host=musoBioLep01 BioLep448 on *Serpocaulon maritimum*, 10.98171°N, 85.42785°W, 740 m, 15–28.iv.2011, DHJPAR0042832 (EMUS).




Figure 7 *Alabagrus axelhausmanni* n. sp., holotype: A. lateral habitus, B. anterior head, C. lateral head, D. dorsal head, E. dorsal habitus, F. wings, G. lateral mesosoma, H. dorsal propodeum and tergum 1.

Alabagrus barbsharanowskiae Sharkey n. sp.

Figure 8

DIAGNOSIS. First tergum with two converging longitudinal carinae. Metapleuron entirely black with a subtle reddish tinge.

NOTES. Formerly considered part of *A. englishi* (Leathers and Sharkey, 2003).

DESCRIPTION. Body length 10 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; with two converging longitudinal carinae. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Barbara Sharanowski, hymenopterist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Costa Rica, Ala(juela), 20 km S Upala (10.84°N, 85.07°W), 14–17.viii.1990, F.D. Parker (EMUS). PARATYPES: 2♀, same data as holotype, except dates 22.i.1991 and 11–21.x.1991 (HIC and EMUS).

Alabagrus bernardoespinozai Sharkey n. sp.

Figure 9

DIAGNOSIS. Gena right angled or acute posteroventrally. First tergum wider than long with well-defined median longitudinal carina. Forewing almost entirely infusate or with some white/clear at apex, ignore small clear patches posterad stigma. Pronotum and propleuron melanic. Hind femur entirely reddish-orange.

NOTES. Formerly considered to belong to *A. latisoma* (Leathers and Sharkey, 2003). Very similar to *A. latisoma*, the holotype of which is from Argentina. Specimens of *A. bernardoespinozai* differ in having about 12 pegs near the apex of the hind tibia, whereas those of the holotype of *A. latisoma* have 25 (Sharkey 1988).

DESCRIPTION. Body length 7.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 0.8× longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor short, not longer than metasoma, or longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Microthyris prolongalis*.

ETYMOLOGY. Named in honor of Sr. Bernardo Espinoza of INBio and the Museo Nacional de Costa Rica, who has long and masterfully taxonomized all of the Erebidae Arctiinae found by the ACG Lepidoptera inventory to date, as well as nearly all for Costa Rica.

DISTRIBUTION. Specimens from Guyana, Ecuador, Colombia, and Venezuela (HIC) appear to fit this species concept, and I have identified them as such but have not included them in the type series.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Microthyris prolongalis* on *Ipomoea batatas*, 10.93332°N, 85.25331°W, 135 m,




Figure 8 *Alabagrus barbsharanowskiae* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

29.vi–16.vii.2009, DHJPAR0036338 (EMUS). PARATYPES: reared specimen: 1♀, same data as holotype, except date 23–29.vii.2012, DHJPAR0049651 (EMUS). Nonreared paratypes: 17♀, from Heredia, Limón (EMUS, HIC, INBio).

Alabagrus bobpoolei Sharkey n. sp.

Figure 10

DIAGNOSIS. Female propodeum completely smooth, male areolate. First median tergite long and narrow with well-defined median longitudinal carina extending most of the length of the tergite. Forefemur melanic. Metapleuron and propodeum entirely black. Median tergite 3 pale anteriorly, melanic posteriorly.

DESCRIPTION. Body length 9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.6× longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or




Figure 9 *Alabagrus bernardoespinozai* n. sp., holotype: A. lateral habitus, B. dorsal head, C. anterior head, D. dorsal habitus, E. wings, F. lateral head and mesosoma, G. propodeum and tergum 1.

depression barely indicated, or with deep transverse depression. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum areolate.

HOST INFORMATION. *Eulepte concordalis*, *Eulepte* Janzen06, *Eulepte* Janzen12, *Eulepte* Solis15.

ETYMOLOGY. Named in honor of Dr. Robert Poole of the U.S. National Museum of Natural History of the Smithsonian Institution, who has long rescued the taxonomic possibility of identification of the ACG inventory of Noctuoidea, in collaboration with other noctuid taxonomists.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Eulepte* Solis15 on *Jacaranda copaia*, 10.91847°N, 85.30338°W, 320 m, 31.iii–20.iv.2011, DHJPAR0042813 (EMUS). PARATYPES: 1♂, from *Eulepte* Janzen06 on *Vitex cooperi*, 11.01087°N, 85.48817°W, 400 m, 9–26.ix.2009, DHJPAR0036694. 1♀, from *Eulepte* Janzen12 on *V. cooperi*, 10.86472°N, 85.41531°W, 540 m, 18.xii.2010–18.i.2011, DHJPAR0041169. 1♂, from *Eulepte* Solis15 on *Hadroanthus guayacan*, 11.01087°N, 85.48817°W, 400 m, 24.viii–6.ix.2009, DHJPAR0037887. 1♀, 1♂, 11.03226°N, 85.52776°W, 290 m, 18.xi–9.xii.2011, DHJPAR0048725, DHJPAR0048727. 1♂, on *Jacaranda copaia*, 10.9867°N, 85.38503°W, 440 m, 9–29.iii.2010, DHJPAR0038924. 1♀, 1♂, 10.91847°N, 85.30338°W, 320 m, 31.iii–18.iv.2011, DHJPAR0042805, DHJPAR0042814. 1♂, 31.iii–19.iv.2011, DHJPAR0042828. 1♀, 8–26.iv.2011, DHJPAR0042815. 1♂, 16.i–6.ii.2013, DHJPAR0051382. 1♂, 16.i–9.ii.2013, DHJPAR0051380. 1♀, 8–30.i.2014, DHJPAR0054514. 1♂, on *Tabebuia chrysantha*, 10.9888°N, 85.42336°W, 680 m, 28.iii–16.iv.2011, DHJPAR0042811. 1♀, 28.iii–22.iv.2011, DHJPAR0045366. 1♂, on *Vitex cooperi*, 10.94741°N, 85.31501°W, 491 m, 27.viii–15.ix.2011, DHJPAR0045785. 1♀, 10.xi–5.xii.2013,

DHJPAR0054495. 1♀, 10.9301°N, 85.25205°W, 109 m, 7–21.viii.2012, DHJPAR0050364 (EMUS, HIC).

***Alabagrus bobrobbinsi* Sharkey n. sp.**

Figure 11

DIAGNOSIS. Similar to *A. karensharkeyae*, distinguished most easily by color. Color almost entirely black except parts of propodeum and metapleuron; forewing infuscate with milky white apex; first median tergite narrow with a strong, long median carina.

NOTES. Formerly considered as part of *A. maya* (Leathers and Sharkey, 2003).

DESCRIPTION. Body length 9.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Hind coxa varying from entirely black to entirely red; propodeum, as in the illustrated female, varying to completely red.

HOST INFORMATION. *Microthyris prolongalis* DHJ02, *Microthyris anormalis* DHJ02, *Psara obscuralis*.

ETYMOLOGY. Named in honor of Dr. Robert Robbins of the U.S. National Museum of Natural History of the Smithsonian Institution,




Figure 10 *Alabagrus bobpoolei* n. sp., A–H holotype; I. paratype male: A. lateral habitus, B. dorsal head, C. anterior head, D. lateral head, E. dorsal habitus, F. wings, G. lateral mesosoma and tergum 1, H. dorsal propodeum and tergum 1, I. dorsal propodeum and tergum 1.

who has long rescued the taxonomic possibility of identification of the ACG inventory of Lycaenidae, in collaboration with other butterfly taxonomists.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Microthyris prolongalis*DHJ02 on *Ipomoea batatas*, 10.76828°N, 85.42567°W, 480 m, 17.x–9.xi.2011, DHJPAR0046733 (EMUS). 1♂, from *Microthyris prolongalis*DHJ02 on *Ipomoea batatas*, 10.89666°N, 85.29003°W, 400 m, 7–26.vi.2012, DHJPAR0049657. 1♂, from *Microthyris prolongalis*DHJ02 on *Ipomoea trifida*, 10.77175°N, 85.434°W, 305 m, 27.v–14.vi.2011, DHJPAR0045007. 1♂, 27.v–15.vi.2011, DHJPAR0045017. 1♀, 30.v–15.vi.2011, DHJPAR0045008. 1♂, from *Microthyris anormalis*DHJ02 on *Merremia umbellata*, 11.00199°N, 85.46166°W, 590 m, 26.viii–7.ix.2009, DHJPAR0037881. 1♀, from *Psara obscuralis* on *Merremia tuberosa*, 11.016020°N, 85.38053°W, 380 m, 6.ii–6.iii.2014, DHJPAR0055103.

Alabagrus bobwhartoni Sharkey n. sp.

Figure 12

DIAGNOSIS. Gena rounded posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Body of meso- and metasoma yellow. Fore- and midfemur entirely pale. Midtibia with more than 15 pegs/spines. Forewing with sharply defined infusate apex.




Figure 11 *Alabagrus bobrobbinsi* n. sp., A, C holotype; B paratype male: A. lateral habitus, B. wings, C. dorsal habitus.




Figure 12 *Alabagrus bobwhartoni* n. sp., holotype: A. dorsal habitus, B. lateral habitus, C. wings, D. midtibia.

DESCRIPTION. Body length 7.1 mm. Gena right angled or acute posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron smooth. Metapleur-

on mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Bob Wharton, braconologist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Limón, Sector Cocorí, 30 km N Cariari, 100 m, ii.1995, Malaise, L_N_286000_567500, H10635, E. Rojas (INBio). PARATYPES: 1♀, Limón, Guacimo, 23.v.1987, P. Hanson (MUCR). 1♀, same data as holotype, except date iii.1995 (HIC).

Alabagrus brendameierottoae Sharkey n. sp.

Figure 13

DIAGNOSIS. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Pronotum mostly yellow. Hind coxa with a black patch laterally. Forewing yellow basally infusate along costal margin and in apical 1/3.

NOTES. Formerly considered part of *A. warrau* (Leathers and Sharkey, 2003).

DESCRIPTION. Body length 6 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually




Figure 13 *Alabagrus brendameierottoae* n. sp., holotype: A. lateral habitus, B. dorsal habitus, C. wings, D. lateral head and anterior mesosoma, E. propodeum.




Figure 14 *Alabagrus brianbrowni* n. sp., holotype: **A.** lateral habitus, **B.** lateral head, **C.** wings, **D.** dorsal habitus.

mostly or completely smooth. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. In appreciation of Mrs. Brenda Meierotto, who ran Malaise traps in Alaska for the Sharkey lab and who is the mother of graduate student extraordinaire Sarah Meierotto.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, Est. Biol. La Selva, swamp edge, 50–100 m, 10.43°N , 84.02°W , 17.ix.1998, H10629 (INBio). PARATYPES: 3♀, same data as holotype, except dates x.1995, v.1996, and iv.1993 (EMUS, INBio, HIC).

Alabagrus brianbrowni Sharkey n. sp.

Figure 14

DIAGNOSIS. Precoxal sulcus absent, crenulae always lacking. Terga 1–4 yellow, 5 to apex melanic. Forefemur mostly melanic. Scutellum yellow.

DESCRIPTION. Body length 8.8 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron smooth. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse

depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Brian Brown, dipterist extraordinaire and dear friend.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, Golfito, Estacion Agujas, 250–300 m, 15.x–15.xi.1999, Malaise, L_S_276750_526550, H10631 (HIC).

Alabagrus brianharrisi Sharkey n. sp.

Figure 15

DIAGNOSIS. Gena rounded posteroventrally. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Forewing pale yellow in basal $\frac{2}{3}$, infusate apically, costal margin melanic. Metapleuron mostly melanic. Propodeum pale.

NOTES. Formerly considered part of *A. roibasi* (Leathers and Sharkey, 2003).

DESCRIPTION. Body length 8.2 mm. Gena right angled or acute posteroventrally. Clypeus unmodified, lacking protrusions. Occiput dorsolaterally rounded. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. First metasomal median tergite wider.

HOST INFORMATION. *Phaedropsis cernalis*, *Microthyris prolongalis*.

ETYMOLOGY. Named in honor of Brian Harris of the U.S. National Museum of Natural History of the Smithsonian Institution, who has long curated and accessioned the ACG inventory of many families of butterflies and moths, in collaboration with other Lepidoptera taxonomists.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phaedropsis cernalis* on *Triplaris melaenodendron*, 10.77175°N , 85.434°W , 305 m, 8.viii–5.ix.2010, DHJPAR0041187 (EMUS). PARATYPES: 1♂, from *Microthyris prolongalis* on *Ipomoea alba*, 10.77175°N , 85.434°W , 305 m, 28.ix–19.x.2011, DHJPAR0046738. 1♂, 10.78506°N , 85.6637°W , 10 m, 17.vii–6.viii.1990, DHJPAR0015477. 1♀, 10.78938°N , 85.55098°W , 85 m, 26.viii–10.ix.2000, DHJPAR0015369. 2♀, 26.viii–20.ix.2000, DHJPAR0015373, DHJPAR0015374. 1♀, 26.viii–27.ix.2000, DHJPAR0015372. 1♀, 26.viii–29.ix.2000, DHJPAR0015371. 1♀, 26.viii–7.ix.2000, DHJPAR0015370. 1♀, 10.81224°N , 85.54438°W , 95 m, 21.viii–5.ix.2000, DHJPAR0015375 (EMUS, HIC).

Alabagrus cara Sharkey, 1988

Figure 16

Alabagrus cara Sharkey, 1988:362. Holotype ♀, Ecuador (CNCI)

DIAGNOSIS. Gena acute posteroventrally. Forefemur and propleuron yellow. Metasoma entirely yellow. Hind coxa with a black patch laterally near apex. Forewing yellow basally, infusate apicad stigma; costal vein mostly yellow.

DESCRIPTION. Body length 6.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae




Figure 15 *Alabagrus brianharrisi* n. sp., holotype: A. anterior head, B. lateral mesosoma, C. lateral habitus, D. dorsal habitus, E. dorsal head, F. lateral head, G. wings.




Figure 16 *Alabagrus cara*, A–D, F–H female; E, I male: A. anterior head, B. dorsal head, C. lateral habitus, D. dorsal habitus, E. forewing, F. wings, G. lateral head and mesosoma, H. propodeum and tergum 1, I. propodeum and tergum 1.




Figure 17 *Alabagrus cocto*, A–C female; D–F male: A. lateral habitus, B. wings, C. dorsal habitus, D. lateral habitus, E. wings, F. dorsal habitus.

posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.6\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeal sculpture stronger; forewing band not as well defined.

HOST INFORMATION. *Desmia benealis*, *Desmia ufeus*.

DISTRIBUTION. The holotype is from lowland costal Ecuador. Specimens are widespread in Costa Rica through Panama and western lowland Ecuador. Undoubtedly present in western Colombia, as well, but not yet recorded.

MATERIAL EXAMINED. Reared specimens: 1♀, from *Desmia benealis*DHJ03 on *Hamelia patens*, 10.93548°N, 85.25314°W, 123 m, 6–19.vii.2013, DHJPAR0052681. 1♀, from *Desmia ufeus* on *Cissus alata*, 10.90037°N, 85.37254°W, 500 m, 8–24.vi.2009, DHJPAR0035232. 1♀, from *Desmia* BioLep22DHJ02 on *Begonia plebeja*, 10.97073°N, 85.31434°W, 326 m, 9–30.i.20161, DHJPAR0058550. 1♀, from *Diacme* BioLep02 on *Thelypteris nicaraguensis*, 10.89666°N, 85.29003°W, 400 m, 7.x–7.xi.2010, DHJPAR0041579 (EMUS, HIC).

Alabagrus cocto Sharkey, 1988

Figure 17

Alabagrus cocto Sharkey, 1988:362. Holotype ♀, Panama (USNM)

DIAGNOSIS. Gena acute posteroventrally. Propodeum areolate. Pronotum melanic anteriorly, pale posteriorly. Hind coxa with a black patch laterally. Ovipositor slightly longer than metasoma.

DESCRIPTION. Body length 8.1 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing with more extensive yellow color; propodeum with more sculpture.

HOST INFORMATION. *Conchylodes arcifera*, *Conchylodes grammaphora*, *Desmia ploralis*, *Desmia ploralis*DHJ01, *Desmia ploralis*DHJ03, *Herpetogramma salbialis*, *Phostria vitrifer*, *Salbia cassidalis*, *Salbia haemorrhoidalis*, *Syllepte* Solis22.

DISTRIBUTION. The holotype is from Panama, and specimens are widespread in Costa Rica and Panama.




Figure 18 *Alabagrus combos*, A–G female; H, I male: A. lateral habitus, B. wings, C. anterior head, D. dorsal head, E. dorsal habitus, F. lateral head and mesosoma, G. propodeum and tergum 1, H. lateral head and mesosoma, I. propodeum and tergum 1.

MATERIAL EXAMINED. Reared specimens: 1 ♂, from *Conchylodes arcifera* on *Verbesina gigantea*, 11.03306°N, 85.42876°W, 400 m, 1–16.viii.2007, DHJPAR0021151. 1 ♂, from *Conchylodes grammaphora* on *Zexmenia virgulta*, 10.90093°N, 85.28915°W, 400 m, 5–22.i.2007, DHJPAR0016927. 1 ♀, from *Desmia ploralis* DHJ01 on *Psychotria grandis*, 10.93548°N, 85.25314°W, 123 m, 5–18.x.2013, DHJPAR0053603. 1 ♀, from *D. ploralis* DHJ03 on *Psychotria chagrensis*, 10.9305°N, 85.37223°W, 527 m, 10.v–5.vi.2012, DHJPAR0048715. 1 ♀, from *Herpetogramma salbialis* on *Zexmenia virgulta*, 11.01926°N, 85.40997°W, 440 m, 28.x–17.xi.2008, DHJPAR0030383. 1 ♀, from *Phostria vitrifera* on *Psychotria grandis*, 11.01618°N, 85.35902°W, 340 m, 22.viii–9.ix.2013, DHJPAR0053599. 1 ♂, from *Salbia haemorrhoidalis* on *Lantana camara*, 10.94076°N, 85.3177°W, 461 m, 26.ix–25.x.2011, DHJPAR0045784. 1 ♀, 10.95991°N, 85.28298°W, 160 m, 6–23.viii.2013, DHJPAR0053598. 1 ♂, 6–24.viii.2013, DHJPAR0053601. 1 ♀, 10.9301°N, 85.25205°W, 109 m, 20.x–4.xi.2012, DHJPAR0050945. 1 ♀, 20.x–6.xi.2012, DHJPAR0050940. 1 ♂, from *Syllepte Solis22* on *Psychotria panamensis*, 10.86472°N, 85.41531°W, 540 m, 2–21.vii.2010, DHJPAR0040345. 1 ♂, 18.vii–5.viii.2010, DHJPAR0041594. 1 ♂, 10.94076°N, 85.3177°W, 461 m, 19.ii–10.iii.2011, DHJPAR0042356. 1 ♀, 15.ix–2.x.2012, DHJPAR0050361 (EMUS, HIC).

Alabagrus combos Leathers and Sharkey, 2003
Figure 18

Alabagrus combos Leathers and Sharkey, 2003:23–24. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Gena acute posteroventrally. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First six terga reddish orange,

remaining terga black. Forewing almost entirely infusate, ignore small clear patches posterad stigma.

DESCRIPTION. Body length 6 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. More sculpture on propodeum and first metasomal median tergite; median ridge of median tergite 1 more acute; most notably the ventral surfaces of all coxae are yellow, whereas those of the female are entirely melanic.

HOST INFORMATION. *Ategumia lotanalis*, *Ategumia lotanalis* DHJ08, *Ategumia lotanalis* DHJ09, *Ategumia matutinalis*, *Ategumia matutinalis* DHJ03, *Conchylodes platinalis*, *Herpetogramma salbialis*, *Herpetogramma Solis10*, *Rhctocraspeda periusalis*, *Pleuroptya Solis03*, *Pycnarmon BioLep66*, *spiloJanzen01* Janzen14DHJ02.

DISTRIBUTION. Widespread in Costa Rica; not yet recorded from Panama.

MATERIAL EXAMINED. Reared specimens: 1 ♀, from *Ategumia lotanalis* on *Conostegia xalapensis*, 10.92291°N, 85.31877°W, 410 m, 25.xii.2014–12.i.2015, DHJPAR0057427. 1 ♀, 10.94076°N, 85.3177°W, 461 m, 6–21.x.2009, DHJPAR0037897. 1 ♀, 10.94741°N, 85.31501°W, 491 m, 5–20.x.2009, DHJPAR0037898.

1♂, 10.96331°N, 85.32243°W, 434 m, 13.xii.2015–9.i.2016, DHJPAR0058538. 1♀, 11.01983°N, 85.41342°W, 445 m, 12.ii–3.iii.2004, DHJPAR0015340. 1♀, on *Miconia trinervia*, 10.90424°N, 85.27120°W, 410 m, 6–27.viii.2014, DHJPAR0056358. 1♀, on *Triolena hirsuta*, 10.90528°N, 85.27882°W, 405 m, 19.xi–10.xii.2013, DHJPAR0054498. 1♂, from *Ategumia lotanalis* DHJ08 on *Arthrostemma ciliatum*, 10.94404°N, 85.31738°W, 455 m, 29.iv–15.v.2011, DHJPAR0042819. 1♂, from *Ategumia lotanalis* DHJ09 on *Conostegia xalapensis*, 10.93548°N, 85.25314°W, 123 m, 23.vii–12.viii.2009, DHJPAR0036376. 1♀, 10.9555°N, 85.28381°W, 221 m, 1–21.viii.2012, DHJPAR0050372. 1♀, on *Miconia lacera*, 10.93332°N, 85.25331°W, 135 m, 27.vi–11.vii.2009, DHJPAR0036341. 1♀, 3–18.vi.2009, DHJPAR0035295. 1♀, 10.93548°N, 85.25314°W, 123 m, 23.vii–14.viii.2009, DHJPAR0036377. 1♂, 10.94404°N, 85.31738°W, 455 m, 30.xii.2011–14.i.2012, DHJPAR0046735. 2♀, from *Ategumia matutinalis* on *Miconia lacera*, 10.89678°N, 85.270010°W, 420 m, 16.vii–4.viii.2014, DHJPAR0055974, DHJPAR0055975. 1♂, from *Ategumia matutinalis* DHJ03 on *Miconia lacera*, 10.93332°N, 85.25331°W, 135 m, 9–25.x.2009, DHJPAR0037955. 1♂, 10.94741°N, 85.31501°W, 491 m, 15–27.ix.2009, DHJPAR0037896. 1♀, 10.98931°N, 85.42581°W, 675 m, 9.x–9.xi.2009, DHJPAR0037914. 1♀, from *Conchylodes platinalis* on *Verbesina turbacensis*, 11.00891°N, 85.40977°W, 500 m, 7–24.ii.2007, DHJPAR0017276. 1♀, from *Herpetogramma salbialis* on *Baltimora recta*, 11.00199°N, 85.46166°W, 590 m, 31.viii–14.ix.2009, DHJPAR0037891. 1♂, on *Clibadium leiocarpum*, 10.94404°N, 85.31738°W, 455 m, 27.viii–8.ix.2010, DHJPAR0041185. 1♀, 10.94741°N, 85.31501°W, 491 m, 20.x–10.xi.2012, DHJPAR0053607. 1♂, 11.0006°N, 85.438°W, 620 m, 26.x–11.xi.2009, DHJPAR0037879. 2♂, 26.x–13.xi.2009, DHJPAR0037884, DHJPAR0037958. 1♂, 26.x–7.xi.2009, DHJPAR0037878. 1♀, 26.x–9.xi.2009, DHJPAR0037877. 1♀, on *Clibadium pittieri*, 10.91589°N, 85.26631°W, 420 m, 14.x–4.xi.2009, DHJPAR0037929. 1♀, on *Melanthera aspera*, 10.96187°N, 85.28045°W, 96 m, 16.ix–1.x.2013, DHJPAR0053635. 1♀, 29.ii–11.iii.2012, DHJPAR0050368. 2♂, *Melanthera nivea*, 10.961870°N, 85.28045°W, 96 m, 4–18.xi.2014, DHJPAR0057437, DHJPAR0057438. 1♂, on *Verbesina turbacensis*, 11.01355°N, 85.42406°W, 510 m, 16.i–5.ii.2013, DHJPAR0057419. 1♀, 16.i–9.ii.2013, DHJPAR0050929. 1♀, on *Zexmenia virgulta*, 10.87741°N, 85.32363°W, 560 m, 10.xi–2.xii.2010, DHJPAR0041562. 1♂, 10.90661°N, 85.28784°W, 400 m, 17.xi–6.xii.2010, DHJPAR0041583. 1♂, 10.91847°N, 85.30338°W, 320 m, 3–20.viii.2010, DHJPAR0041598. 1♂, 3–22.viii.2010, DHJPAR0041596. 1♂, from *Herpetogramma Solis10* on *Solanum hazenii*, 10.89678°N, 85.27001°W, 420 m, 22.xii.2011–6.i.2012, DHJPAR0046739. 1♀, 22.xii.2011–8.i.2012, DHJPAR0046734. 1♀, from *Pleuroptya Solis03* on *Laportea aestuans*, 10.96187°N, 85.28045°W, 96 m, 14.iii–1.iv.2012, DHJPAR0049475. 1♀, 11.02681°N, 85.49547°W, 290 m, 23.x–8.xi.2004, DHJPAR0009361. 1♂, on *Urera caracasana*, 10.87766°N, 85.39343°W, 645 m, 5–19.xii.2012, DHJPAR0051354. 1♂, 5–21.xii.2012, DHJPAR0051350. 1♂, on *Urera elata*, 10.87097°N, 85.39144°W, 640 m, 22.ix–11.x.1998, DHJPAR0015339. 1♀, 22.ix–12.x.1998, DHJPAR0015338. 1♀, 22.ix–13.x.1998, DHJPAR0015337. 1♂, 10.87766°N, 85.39343°W, 645 m, 23.i–11.ii.2013, DHJPAR0051383. 1♂, 23.i–9.ii.2013, DHJPAR0051384. 1♀, 10.90661°N, 85.28784°W, 400 m, 22.ii–8.iii.2006, DHJPAR0015530. 1♀, 10.9305°N, 85.37223°W, 527 m, 14.i–18.ii.2009, DHJPAR0030609. 1♀, from *Pycnarmon BioLep66* on *Lepidaploa tortuosa*, 11.01983°N, 85.41342°W, 445 m, 3–31.iii.2009, DHJPAR0035529. 1♂, from *Rhectocraspeda periusalis* on *Piper auritum*,




Figure 19 *Alabagrus craigevansi* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

10.94076°N, 85.3177°W, 461 m, 17–31.i.2010, DHJPAR0038812. 1♀, on *Piper umbellatum*, 10.9301°N, 85.25205°W, 109 m, 7–26.iii.2010, DHJPAR0038807. 1♀, 10.93548°N, 85.25314°W, 123 m, 22.xi–5.xii.2013, DHJPAR0054506. 1♀, 10.94404°N, 85.31738°W, 455 m, 24.viii–7.ix.2010, DHJPAR0041186. 1♀, from *spilo* Janzen01 Janzen14 DHJ02 on *Piper* 21153, 10.89666°N, 85.29003°W, 400 m, 12.iv–8.v.2011, DHJPAR0042826 (EMUS, HIC).

***Alabagrus craigevansi* Sharkey n. sp.**

Figure 19

DIAGNOSIS. Gena rounded posteroventrally. Propleuron melanic and yellow. Hind coxa with a melanic patch laterally. Propodeum entirely smooth except for a weak, rounded median longitudinal bulge. Scutellum entirely yellow.

NOTES. Included under *A. imitatus* in Leathers and Sharkey (2003).



Figure 20 *Alabagrus cuna*: A, C–G female; B male: A. lateral habitus, B. dorsal habitus, C. dorsal habitus, D. lateral head and mesosoma, E. propodeum and tergum 1, F. wings, G. anterior head.

DESCRIPTION. Body length 7.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Mr. Craig Evans, extraordinary friend.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, 300 m, deciduous woods (25–30 yr), open site, 24.viii–14.ix.1985, Gauld and Janzen, SE50, H10963 (HIC).

Alabagrus cuna Sharkey, 1988

Figure 20

Alabagrus cuna Sharkey, 1988:362. Holotype ♀, Panama (AMNH)

DIAGNOSIS. Female: Propleuron pale. Propodeum areolate, with at least one closed areola. Midfemur pale. Head melanic. Hind coxa entirely pale. Forewing yellow in basal ⅓, melanic apically. Midfemur entirely or mostly pale. Ovipositor slightly longer than metasoma. Male: metasoma entirely melanic, contrasting with a yellow mesosoma. Forefemur melanic, or pale apically and basally, melanic at midlength.

DESCRIPTION. Body length 5.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply

crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.4× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor short, not longer than metasoma, or longer than metasoma but not longer than body.

Males. First median tergite with well-defined median carina; forewing entirely infuscate, lacking yellow color; metasomal terga mostly melanic; legs more melanic, e.g., hind leg completely black.

HOST INFORMATION. *Neurophyseta* BioLep226.

DISTRIBUTION. Costa Rica and Panama (Leathers and Sharkey, 2003).

MATERIAL EXAMINED. Reared specimens: all from *Neurophyseta* BioLep226 on *Hypolepis repens*. 1♂, 10.89666°N, 85.29003°W, 400 m, 21.ii–6.iii.2011, DHJPAR0043025 (EMUS). 1♀, 21.ii–7.iii.2011, DHJPAR0042807. 1♂, 21.ii–8.iii.2011, DHJPAR0043033. 1♀, 4–20.iii.2012, DHJPAR0049045. 1♀, 4–21.iii.2012, DHJPAR0049047. 1♀, 1♂, 10.90661°N, 85.28784°W, 400 m, 27.xii.2013–10.i.2014, DHJPAR0054521, DHJPAR0054535. 1♀, 27.xii.2013–14.i.2014, DHJPAR0054525 (EMUS, HIC).

Alabagrus derailersi Leathers and Sharkey, 2003

Figure 21

Alabagrus derailersi Leathers and Sharkey, 2003:25–26. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Female: Gena rounded posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with a smooth shallow groove extending anteriorly. Propodeum mostly smooth with a few crenulae anteromedially but lacking areolae. Mesoscutum yellow. Propleuron melanic. Fore- and midcoxae entirely yellow. Ovipositor




Figure 21 *Alabagrus derailersi*, A–D paratype; E, F male: A. lateral habitus, B. lateral head, C. wings, D. dorsal habitus, E. lateral habitus, F. dorsal mesosoma and tergum 1.

slightly shorter than body. Male: Mesoscutum mostly reddish orange with some melanic infusions, or mostly melanic with some reddish orange infusions. Scutellum reddish orange.

DESCRIPTION. Body length 5.9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Mesoscutum mostly reddish orange with some melanic infusions, or mostly melanic with some reddish orange infusions; mesoscutellum reddish orange.

DISTRIBUTION. Costa Rica (Leathers and Sharkey, 2003).

Alabagrus donharveyi Sharkey n. sp.

Figure 22

DIAGNOSIS. Median tergite 1 mostly melanic but red at base. Hind coxa entirely red. Metapleuron mostly or entirely melanic.

NOTES. Specimens of this species were included under *A. semialbus* in Leathers and Sharkey (2003). The holotype of *A. semialbus* is from Brazil, and the metapleuron is more areolated than those of *A. donharveyi*.

DESCRIPTION. Body length 8.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral half or more. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities,

distinctly more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Similar to female.

HOST INFORMATION. *Coelorhynchidia pandarali*, *Coenostolopsis apicalis*.

ETYMOLOGY. Named in honor of Dr. Donald Harvey of the U.S. National Museum of Natural History of the Smithsonian Institution, who has long curated and accessioned the ACG inventory of many families of butterflies and moths, in collaboration with other Lepidoptera taxonomists.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Coenostolopsis apicalis* on *Dalechampia websteri* 10.93332°N, 85.25331°W, 135 m, 14.xii.2009–4.i.2010, DHJPAR0037930 (EMUS). PARATYPES: reared specimens: 1♀, from *Coelorhynchidia pandaralis* on *Dichapetalum grayumii*, 10.9305°N, 85.37223°W, 527 m, 4–21.iv.2007, DHJPAR0021165. 1♀, 1♂, DHJPAR0021158, DHJPAR0021163. 1♂, from *Coenostolopsis apicalis* on *Dalechampia scandens*, 10.992840°N, 85.42936°W, 660 m, 25.i–19.ii.2014, DHJPAR0054512. 1♀, 25.i–22.ii.2014, DHJPAR0054513. 1♀, on *Dalechampia websteri*, 10.93332°N, 85.25331°W, 135 m, 20.ix–3.x.2009, DHJPAR0036711. 1♀, 20.ix–8.x.2009, DHJPAR0036688. 1♀, 10.93548°N, 85.25314°W, 123 m, 4–15.ix.2011, DHJPAR0045810. 1♀, 11.01926°N, 85.40997°W, 440 m, 21.ii–16.iii.2012, DHJPAR0048734 (EMUS, HIC). Nonreared paratypes: 1♀, Guanacaste, Estación Pitilla, P.N. Guanacaste, 700 m, 9 km S Santa Cecilia, 19.v–3.vi.1993, C. Moranga, L-N-330200, 380200 (INBio). 4♀, 2♂, Alajuela, 20 km S Upala (10.84°N, 85.07°W), 16–24.vii.1995, 1–10.iv.1991, 20–26.iii.1991, 21–30.iv.1991, 12.iii.1991, 10–19.iii.1991, F.D. Parker (EMUS, HIC). 2♀, Guanacaste, 3 km SE R. Narango, 20–31.i.1992 and 11.i.1993, F.D. Parker (EMUS, HIC). 1♀, PANAMA, Canal Zone, Barro Colorado Is., 9.17°N, 79.83°W, 14.viii.1978, H. Hespeneheide (HIC). 1♀, ECUADOR, Sucumbios, Rio Napo, Sacha Lodge, 0.5°S, 76.5°W, 290 m, 31.x–10.xi.1994, Hibbs (EMUS).




Figure 22 *Alabagrus donharveyi* n. sp., paratype: **A.** anterior head, **B.** lateral habitus, **C.** dorsal head, **D.** dorsal habitus, **E.** lateral head, **F.** wings, **G.** lateral mesosoma, **H.** lateral head and mesosoma.

Alabagrus donlafontainei Sharkey n. sp.

Figure 23

DIAGNOSIS. Female: Propodeum areolate, with at least one closed areola. Ovipositor slightly shorter than body. Head melanic. Propleuron. Midfemur yellow. Hind coxa entirely yellow. Forewing mostly melanic with a clear vertical band near stigma. Male: Terga 1 and 2 mostly or entirely yellow, remaining terga mostly or entirely black. Midfemur melanic apically and basally, pale elsewhere.

DESCRIPTION. Body length 4.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge, or with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing evenly infusate, lacking clear band; hind leg and metasomal terga mostly melanic.

HOST INFORMATION. musoBioLep01 BioLep448.

ETYMOLOGY. Named in honor of Dr. Don Lafontaine of the Canadian National Collection, Ottawa, who has long labored at helping the taxonomic identification of the ACG inventory of Noctuoidea, in collaboration with other moth taxonomists.

MATERIAL EXAMINED. HOLOTYPE ♀, host=musoBioLep01 BioLep448 on *Polypodium fraxinifolium*, 10.98816°N, 85.39581°W,

485 m, 4–21.iii.2010, DHJPAR0038929 (EMUS). PARATYPES: all from musoBioLep01 BioLep448 on *Polypodium fraxinifolium*: 1♂, 10.98816°N, 85.39581°W, 485 m, 4–21.iii.2010, DHJPAR0038927. 1♂, 10.98816°N, 85.39581°W, 485 m, 4–23.iii.2010, DHJPAR0038923. 2♂, 4–23.iii.2010, DHJPAR0038928, DHJPAR0050137 (EMUS, HIC).

Alabagrus donnai Leathers and Sharkey, 2003

Figure 24

Alabagrus donnai Leathers and Sharkey, 2003:26–27. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Gena acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with a smooth groove extending anteriorly. First tergum weakly convex. Body black and yellow. Mesoscutum melanic. Forefemur melanic. Metasoma entirely yellow.

DESCRIPTION. Body length 5.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Pale parts of body darker than females, pale orange rather than dark or pale yellow.

DISTRIBUTION. Widespread in Costa Rica and also known from Yucatán, Mexico (Leathers and Sharkey, 2003).




Figure 23 *Alabagrus donlafontainei* n. sp., **A–D, F–H** holotype; **E** male paratype: **A.** anterior head, **B.** lateral habitus, **C.** dorsal head, **D.** lateral head and mesosoma, **E.** dorsal wing, propodeum and metasoma, **F.** wings, **G.** propodeum and tergum 1, **H.** dorsal habitus.




Figure 24 *Alabagrus donnai*, **A–C** paratype female; **D** male: **A.** lateral habitus, **B.** wings, **C.** dorsal habitus, **D.** dorsal habitus.




Figure 25 *Alabagrus ekchuah*: A. lateral habitus, B. dorsal habitus, C. wings, D. lateral mesosoma, E. propodeum and tergum 1, F. anterior head, G. dorsal head, H. lateral head.

Alabagrus ekchuah Sharkey, 1988

Figure 25

Alabagrus ekchuah Sharkey, 1988:366–7. Holotype ♀, Costa Rica (EMUS)

DIAGNOSIS. Body entirely black. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Forewing almost entirely infusate, ignore small clear patches posterad stigma.

DESCRIPTION. Body length 10.7 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.7\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Fore- and midtarsi yellow; propodeum with slightly more sculpture.

HOST INFORMATION. *Pantographa suffusalis*, *Syllepte amando*DHJ02.

DISTRIBUTION. Known from Costa Rica, El Salvador, and Mexico.

MATERIAL EXAMINED. Reared specimens: 1♂, from *Pantographa suffusalis* on *Triumfetta bogotensis*, 10.88919°N, 85.47609°W, 600 m, 16–28.viii.2004, DHJPAR0015405 (EMUS). 1♀, 16.viii–7.ix.2004, DHJPAR0015404. 1♀, from *Syllepte amando*DHJ02 on *Prockia costaricensis*, 11.01454°N, 85.47492°W, 380 m, 26.viii–

23.ix.2002, DHJPAR0015403. 1♀, 11.01087°N, 85.48817°W, 400 m, 27.v–17.viii.2007, DHJPAR0021168 (EMUS, HIC).

Alabagrus englishi Leathers and Sharkey, 2003

Figure 26

Alabagrus englishi Leathers and Sharkey, 2003:27–28. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Precoxal sulcus with one or several distinct foveae posteroventrally or absent. Propodeum smooth. Ovipositor longer than body. Forewing weakly infusate in basal $\frac{2}{3}$, more infusate apically. Pronotum melanic. Hind coxa, metapleuron, and metasoma all entirely orange.

NOTES. The concept of *A. englishi* is much changed from that of Leathers and Sharkey (2003) and has a much more restricted morphological definition. Many paratypes of *A. englishi*, sensu Leathers and Sharkey (2003), will now key to the following species: *Alabagrus johnobryckii*, *A. sarahmeierottoae*, *A. sarahsharkeyae*, and *A. barbsharanowskiae*.

DESCRIPTION. Body length 7.6 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.3\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking




Figure 26 *Alabagrus englishi*, A–D paratype; E, F male: A. lateral habitus, B. wings, C. dorsal habitus, D. lateral head and mesosoma, E. propodeum and tergum 1, F. lateral habitus.

transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown. The concept of *A. englishi* in Leathers and Sharkey (2003) conflates many species, and without *COI* sequence data, it is not possible at this time to discern which males are conspecific with the holotype female.

DISTRIBUTION. Costa Rica.

Alabagrus fernandezi Sharkey n. sp.

Figure 27

DIAGNOSIS. Gena acute posteroventrally. Midtibia with zero to three nonapical spines. Precoxal sulcus short with a few crenulae restricted to posterior margin. Propodeum smooth with a subtle hint of smooth shallow areolae. First tergum weakly convex. All femora, metapleuron, and hind coxa all entirely yellow. Mesoscutum melanic. Forewing with two yellow bands, apical band complete to posterior margin, costal vein yellow.

NOTES. Specimens of this species were included under *A. pachamama* in Leathers and Sharkey (2003). *Alabagrus pachamama* does not occur in Costa Rica. Members are very similar to *A. andresfreitasi* but have fewer spines on midtibia and more sculpture on propodeum. *Alabagrus fernandezi* is also similar to some small specimens of *A. fernandodiasi*. The color of the midfemur, with at least a hint of melanic color and a smoother propodeum in the latter, will distinguish these species.

DESCRIPTION. Body length 7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal

surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Fernando Fernandez, in thanks for his collaboration during years of research in Colombia.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, Puerto Viejo, OTS, La Selva, 100 m, i.1993, P. Hanson, H17270 (HIC). PARATYPES: 23 ♀, from the following provinces: Limón, Puntarenas, Heredia (HIC, EMUS, INBio, MUCR).

Alabagrus fernandodiasi Sharkey n. sp.

Figure 28

DIAGNOSIS. Gena acute posteroventrally. First tergum weakly convex. Forefemur, hind coxa, and metapleuron all entirely pale. Mesoscutum melanic. Propodeum lacking complete areolae, usually mostly or completely smooth. Midfemur yellow at base and apex but melanic at midlength, sometimes only ventrally.

NOTES. Specimens of this species were included under *A. pachamama* in Leathers and Sharkey (2003). *Alabagrus pachamama* as it is circumscribed here does not occur in Costa Rica.

DESCRIPTION. Body length 8.1 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking




Figure 27 *Alabagrus fernandezi* n. sp., A–D, F holotype; E paratype: A. lateral habitus, B. dorsal habitus, C. lateral head and mesosoma, D. propodeum and tergum 1, E. propodeum, F. wings.

transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Rhectocraspeda* Solis05, *Rhectocraspeda* Janzen347, *Desmia benealis* DHJ02, *Desmia* Janzen19, *Herpetogramma* Solis10, *Rhectocraspeda periusalis*, spiloBioLep01 BioLep243, spiloBioLep01 BioLep311, spiloBioLep01 BioLep402, spiloBioLep01 BioLep403, spiloBioLep01 BioLep414, spiloBioLep01 BioLep415.

ETYMOLOGY. Named in honor of Dr. Fernando Dias of the Universidade Federal do Paraná, Brazil, in recognition of his careful and precise identification and description of ACG Nymphalidae.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Rhectocraspeda* Solis05 on *Cestrum rugulosum*, 11.016020°N, 85.380530°W, 380 m, 8.viii–1.ix.2014, DHJPAR0056299 (EMUS). PARATYPES: 1 ♀, from *Desmia benealis* DHJ02 on *Besleria columnneoides*, 10.8962°N, 85.27769°W, 430 m, 10–25.ix.2012, DHJPAR0050360. 1 ♀, from




Figure 28 *Alabagrus fernandodiasi* n. sp., holotype: A. lateral habitus, B. dorsal habitus, C. lateral head and mesosoma, D. propodeum and tergum 1, E. wings.




Figure 29 *Alabagrus genemonroei* n. sp., holotype: **A.** dorsal habitus, **B.** lateral habitus, **C.** propodeum and tergum 1, **D.** lateral head and mesosoma, **E.** wings.

Desmia Janzen19 on *Cissus verticillata*, 10.87766°N, 85.39343°W, 645 m, 31.v–22.vi.2013, DHJPAR0052090. 1♀, from *Herpetogramma* Solis10 on *Justicia aurea*, 10.86472°N, 85.41531°W, 540 m, 23.x–7.xi.2010, DHJPAR0041163. 1♀, on *Justicia macrantha*, 11.02364°N, 85.49139°W, 290 m, 24.i–12.ii.2005, DHJPAR0009377. 1♀, from *Herpetogramma* Solis11 on *Solanum aturense*, 10.89666°N, 85.29003°W, 400 m, 26.ii–15.iii.2014, DHJPAR0055088. 1♀, 26.ii–17.iii.2014, DHJPAR0055236. 1♀, on *Solanum rovirosanum*, 10.90037°N, 85.37254°W, 500 m, 12–30.iii.2012, DHJPAR0048731. 1♀, 12–31.iii.2012, DHJPAR0048732. 1♀, 10.87766°N, 85.39343°W, 645 m, 4–24.iv.2013, DHJPAR0052085. 1♀, 10.88009°N, 85.38887°W, 575 m, 1–25.v.2014, DHJPAR0055818. 1♀, on *Solanum schlechtendalianum*, 11.02367°N, 85.41884°W, 375 m, 16.i–5.ii.2007, DHJPAR0021198. 1♀, 16.i–6.ii.2007, DHJPAR0017273. 1♀, 10.86472°N, 85.41531°W, 540 m, 2–22.vii.2010, DHJPAR0040215. 1♀, 27.viii–13.ix.2011, DHJPAR0045801. 1♀, on *Witheringia solanacea*, 10.94076°N, 85.3177°W, 461 m, 29.x–13.xi.2013, DHJPAR0054488. 2♀, from *Rhctocraspeda* Janzen347 on *Solanum lanceifolium*, 11.01087°N, 85.48817°W, 400 m, 25.viii–11.ix.2009, DHJPAR0037883, DHJPAR0037890. 1♀, from *Rhctocraspeda periusalis* on *Piper aequale*, 10.90445°N, 85.28412°W, 400 m, 7.xii.2011–1.i.2012, DHJPAR0046750. 1♀, on *Piper sancti-felices*, 10.86472°N, 85.41531°W, 540 m, 20.iii–10.iv.2011, DHJPAR0042804. 1♀, on *Piper urostachyum*, 10.98705°N, 85.42816°W, 700 m, 7–27.iii.2014, DHJPAR0055099. 1♀, 7–29.iii.2014, DHJPAR0055098. 1♀, on *Piper auritum*, 10.93548°N, 85.25314°W, 123 m, 12–27.x.2012, DHJPAR0050924. 1♀, from spiloBioLep01 BioLep243 on *Solanum schlechtendalianum*, 10.9305°N, 85.37223°W, 527 m, 10–27.xi.2013, DHJPAR0054490. 1♀, from spiloBioLep01 BioLep311 on *Columnea purpurata*, 10.8962°N, 85.27769°W, 430 m, 10–25.vi.2013,

DHJPAR0052089. 1♀, from spiloBioLep01 BioLep402 on *Mikania cordifolia*, 10.86472°N, 85.41531°W, 540 m, 11–27.viii.2011, DHJPAR0045800. 1♀, 11–29.viii.2011, DHJPAR0045799. 1♀, from spiloBioLep01 BioLep403 on *M. cordifolia*, 11.01983°N, 85.41342°W, 445 m, 12–26.viii.2011, DHJPAR0044991. 1♀, 12–29.viii.2011, DHJPAR0045804. 1♀, from spiloBioLep01 BioLep414 on *Mikania guaco*, 10.87766°N, 85.39343°W, 645 m, 1–20.iv.2013, DHJPAR0052081. 1♀, 1–21.iv.2013, DHJPAR0052080. 1♀, from spiloBioLep01 BioLep415 on *Tournefortia bicolor*, 10.8702°N, 85.39153°W, 640 m, 8–22.xi.2014, DHJPAR0057002 (EMUS, HIC).

***Alabagrus genemonroei* Sharkey n. sp.**

Figure 29

DIAGNOSIS. Gena acute posteroventrally. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Third tergum lacking transverse depression, or depression barely indicated. Forewing with two distinct yellow bands, one basal and the other directly below (posterior to) stigma; distal band complete to hind margin; costal vein mostly yellow. Forefemur and mesoscutum black. Metapleuron entirely yellow. Terga 1–3 yellow, remaining terga black.

NOTES. Specimens of this species will key to *A. pachamama* in Leathers and Sharkey (2003). *Alabagrus pachamama* does not occur in Costa Rica.

DESCRIPTION. Body length 8.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present,

restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge, or with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor short, not longer than metasoma, or longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Lamprosema* Dapkey23.

ETYMOLOGY. Named in honor of Dr. Gene Munroe (deceased) of the Canada Department of Agriculture, Ottawa, in recognition of his enormous contributions to the taxonomy and systematics of Pyraloidea of ACG and the world.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Lamprosema* Dapkey23 on *Serjania valerii*, 10.9163°N, 85.37869°W, 460 m, 13.i–1.ii.2006, DHJPAR0015510 (EMUS). PARATYPES: 16♀, from the following provinces: Limón, Guanacaste, Heredia (HIC, EMUS, INBio, MUCR). 1♀, PANAMA, Canal Zone, 20.vii.1978 (HIC).

Alabagrus hansonii Sharkey n. sp.

Figure 30

DIAGNOSIS. Body predominantly yellow. Margins of mesopleuron with a greenish tinge.

DESCRIPTION. Body length 9 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron smooth. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.6× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Paul Hanson, hymenopterist extraordinaire, for the many occasions on which he facilitated research in Costa Rica.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, RF Golgo Dulce el 200 m, 24 km W Piedras Blancas, P. Hanson, ix.1992, H10966 (HIC). PARATYPE: 1♀, same data as holotype, except date vi.1993 (MUCR).

Alabagrus hespenheidei Sharkey n. sp.

Figure 31

DIAGNOSIS. Gena rounded posteroventrally. Hind coxa entirely or almost entirely melanic. Median tergite 1 red at base and melanic apically. Median tergite 2 melanic.

NOTES. Referred to as *A. semialbus* by Leathers and Sharkey (2003). The holotype of *A. semialbus* is from Brazil, and the species does not occur in Costa Rica.

DESCRIPTION. Body length 9.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present,




Figure 30 *Alabagrus hansonii* n. sp., holotype: A. wings, B. lateral habitus, C. dorsal habitus.

restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing color variable, from similar to the female to almost completely infuscate; propodeum from almost completely black to completely black; propodeum with more sculpture.

ETYMOLOGY. Named in honor of Dr. Henry Hespeneide, coleopterist extraordinaire, for his many gifts of reared and otherwise collected braconids.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, La Selva Biol. Sta., 3 km S Pto. Viejo, 10.43°N, 84.02°W, 25.vii.1992, H. Hespeneide, H17275 (HIC). PARATYPES: 1♀, 7♂, Heredia Prov. (HIC, EMUS, INBio).

Alabagrus iankitchingi Sharkey n. sp.

Figure 32

DIAGNOSIS. Gena acute posteroventrally. Propodeum smooth with a subtle hint of smooth shallow areolae. Third tergum lacking transverse depression, or depression barely indicated. Forewing with two yellow bands, one distinct basal band and a weaker triangular band directly below (posterior to) stigma that is not complete to the posterior wing margin. Mesoscutum melanic. Median terga 1–4 orange, remaining terga black.

DESCRIPTION. Body length 7.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly.




Figure 31 *Alabagrus hespenheidei* n. sp., A–C, E, F holotype; D paratype male: A. lateral habitus, B. dorsal habitus, C. wings, D. forewing, E. lateral head and mesosoma, F. propodeum and tergum 1.

Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Phaenodropsis* Solis350DHJ02.

ETYMOLOGY. Named in honor of Dr. Ian Kitching of The Natural History Museum, London, in recognition of his very fruitful labors and web site for the taxonomy and systematics of Sphingidae of ACG and the world.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phaenodropsis* Solis350DHJ02 on *Guazuma ulmifolia*, 10.76261°N, 85.42979°W, 330 m, 24.x–5.xii.2013, DHJPAR0054545 (EMUS). PARATYPE: 1 ♀, same host data as holotype, 10.7648°N, 85.38445°W, 550 m, 29.vi–21.vii.2011, DHJPAR0045001 (HIC).




Figure 32 *Alabagrus iankitchingi* n. sp., paratype: A. anterior head, B. dorsal head, C. lateral habitus, D. dorsal habitus, E. wings, F. lateral head and mesosoma.

Alabagrus ilgookangi Sharkey n. sp.

Figure 33

DIAGNOSIS. Gena rounded posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Forewing weakly infusate in basal $\frac{2}{3}$ with a yellowish tinge, more deeply infusate apically. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Pronotum melanic anteriorly, pale (yellow to orange) posteriorly.

DESCRIPTION. Body length 7.6 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Mr. Ilgoo Kang, former graduate student in my (M.J.S.) program.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, San Vito, Estac. Biol. Los Altures, 1500 m, iv.1992, P. Hanson, H10695 (HIC). PARATYPES: 4 ♀, from San Jose and Puntarenas Provinces, all between 1200 and 1500 m (HIC, EMUS, MUCR, INBio).

Alabagrus isidrochaconi Sharkey n. sp.

Figure 34

DIAGNOSIS. Difficult or impossible to separate morphologically from *Alabagrus jeanfrancoislandryi* and *A. jennyphillipsae*. Gena rounded posteroventrally. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Head less elongate than *A. arua*. Hind femur narrower (lateral view) than *A. nicoya*. First metasomal median tergite slightly longer than wide; weakly convex in females and with a rounded longitudinal bulge in males. Pronotum entirely melanic. Metapleuron entirely orange. Terga all orange, except apical two terga slightly to distinctly melanic.

NOTES. Specimens of this species were included under *A. albispina* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 7.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth, sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Medial ridge of median tergite 1 more acute; forewing color variable, from as in the female to completely infusate; hind coxa color variable, from as in the female to completely orange.

HOST INFORMATION. *Eulepte* Janzen03, *Glyphodes sibillalis* DHJ01, *Phaedropsis cernalis*, *Phaedropsis* Solis348.

ETYMOLOGY. Named in honor of Sr. Isidro Chacon of INBio and the Museo Nacional de Costa Rica, who has long and masterfully




Figure 33 *Alabagrus ilgookangi* n. sp., A, C–E holotype; B paratype male: A. lateral habitus, B. propodeum, C. wings, D. dorsal head and mesosoma, E. dorsal metasoma.

taxonomized all the ACG macrolepidoptera inventory to date, as well as for all of Costa Rica.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Glyphodes sibillalis* DHJ01 on *Trophis racemosa*, 11.00217°N, 85.46337°W, 560 m, 15.vii–1.viii.2005, DHJPAR0015520 (EMUS). PARATYPES: 1 ♂, from *Eulepte* Janzen03 on *Amphilophium crucigerum*, 10.88996°N, 85.47966°W, 550 m, 24.ix–13.x.2004, DHJPAR0015525. 1 ♀, 1 ♂, from *Glyphodes sibillalis* DHJ01 on *Trophis racemosa*, 10.72750°N, 85.41212°W, 315 m, 14.xi–4.xii.2014, DHJPAR0056985, DHJPAR0057426. 1 ♀, 10.72750°N, 85.41212°W, 315 m, 14.xi–5.xii.2014, DHJPAR0056987. 1 ♀, 10.83013°N, 85.61372°W, 285 m, 23.vii–5.viii.1990, DHJPAR0015466. 1 ♀, 10.84389°N, 85.61384°W, 300 m, 19.vii–4.viii.1981, DHJPAR0015469. 1 ♀, 6–18.vii.1981, DHJPAR0015467. 1 ♀, 6–19.vii.1981, DHJPAR0015468. 1 ♂, 15–31.vii.2005, DHJPAR0009405. 1 ♀, 11.00217°N, 85.46337°W, 560 m, 5–22.viii.2005, DHJPAR0015523. 2 ♂, 11.03226°N, 85.52776°W, 290 m, 8–19.viii.2013, DHJPAR0052674, DHJPAR0052684. 2 ♂, 8–20.viii.2013, DHJPAR0052675, DHJPAR0052676,




Figure 34 *Alabagrus isidrochaconi* n. sp., A, D–G, I–K holotype; B, C, H paratype males: A. lateral habitus, B. forewing, C. hind leg, D. wings, E. dorsal habitus, F. lateral mesosoma, G. propodeum and tergum 1, H. propodeum and tergum 1, I. dorsal head, J. anterior head, K. lateral head.

DHJPAR0052683. 2 ♀, 1 ♂, 8–21.viii.2013, DHJPAR0053609, DHJPAR0053624, DHJPAR0053631. 2 ♂, 8–22.viii.2013, DHJPAR0053626, DHJPAR0053629. 4 ♀, 8–23.viii.2013, DHJPAR0053630, DHJPAR0053632, DHJPAR0053633, DHJPAR0053634. 3 ♀, 8–26.viii.2013, DHJPAR0053610, DHJPAR0053627, DHJPAR0053628. 1 ♀, 8–28.viii.2013, DHJPAR0053608. 1 ♀, from *Phaenodropsis cernalis* on *Triplaris melaenodendron*, 10.81224°N, 85.54438°W, 95 m, 21.viii–7.ix.2000, DHJPAR0015325. 1 ♀, from *Phaenodropsis* Solis348 on *Triplaris melaenodendron*, 10.8269°N, 85.60413°W, 240 m, 24.xi–29.xii.1983, DHJPAR0015470. 1 ♀, from unknown on unknown, 10.83575°N, 85.61253°W, 290 m, 8–20.vi.1995, DHJPAR0015323 (EMUS, HIC).

Alabagrus jackiemillerae Sharkey n. sp.

Figure 35

DIAGNOSIS. Median tergite 3 pale anteriorly, melanic posteriorly. Midfemur pale basally and apically, otherwise melanic. Third tergum lacking transverse depression, or depression barely indicated. Apical yellow band of forewing reaching posterior wing margin.

NOTES. Specimens of this species were included under *A. pachamama* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 8.6 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae,

usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. spiloBioLep01 BioLep403, *Pleuroptya* Solis191, *Desmia* Janzen07, *Herpetogramma salbialis*, *Herpetogramma* Solis11, *Phostria* Janzen03, *Phostria* Janzen05, *Rhectocraspeda* Janzen347, *Rhectocraspeda* Solis05, *Pilocrocis purpurascens*, *Pleuroptya* Solis03.

ETYMOLOGY. Named in honor of Dr. Jackie Miller of the McGuire Center for Lepidoptera and Biodiversity of the University of Florida in recognition of her contributions to the taxonomy and systematics of Neotropical butterflies and their biodiversity.

MATERIAL EXAMINED. HOLOTYPE ♀, host=spiloBioLep01 BioLep403 on *Mikania cordifolia*, 11.01983°N, 85.41342°W, 445 m, 12.viii–5.ix.2011, DHJPAR0045802 (EMUS). PARATYPES: 1 ♀, from *Pleuroptya* Solis191 on *Justicia aurea*, 11.01926°N, 85.40997°W, 440 m, 4–29.ix.2003, DHJPAR0015350. 1 ♀, from *Desmia* Janzen07 on *Psychotria microbotrys*, 10.9959°N, 85.39842°W, 470 m, 27.viii–17.ix.2013, DHJPAR0053593. 1 ♀, from *Desmia* Solis19 on *Psychotria berteriana*, 10.94076°N, 85.3177°W, 461 m, 9–22.ix.2009, DHJPAR0037895. 1 ♀, 11.01373°N, 85.42531°W, 525 m, 16.ix–3.x.2010, DHJPAR0041165. 1 ♀, 10.94741°N, 85.31501°W, 491 m, 19.v–14.vi.2012, DHJPAR0048729. 1 ♀, 10.90037°N, 85.37254°W, 500 m, 12.ix–3.x.2014, DHJPAR0057448. 1 ♀, 10.944040°N, 85.31738°W, 455 m, 22.xi–11.xii.2014,




Figure 35 *Alabagrus jackiemilleriae* n. sp., holotype: A. anterior head, B. lateral head, C. dorsal head, D. lateral habitus, E. dorsal habitus, F. wings, G. lateral mesosoma, H. propodeum and tergum 1.

DHJPAR0057441. 2♀, 10.940760°N, 85.3177°W, 461 m, 16.xi–13.xii.2014, DHJPAR0057439, DHJPAR0057440. 1♀, from *Herpetogramma salbialis* on *Clibadium pittieri*, 10.89666°N, 85.29003°W, 400 m, 13.ix–1.x.2011, DHJPAR0045798. 1♀, from *Herpetogramma Solis11* on *Solanum aturense*, 10.89666°N, 85.29003°W, 400 m, 21.v–13.vi.2013, DHJPAR0052078. 1♀, on *Solanum rovirosanum*, 10.9305°N, 85.37223°W, 527 m, 10–31.i.2015, DHJPAR0057449. 1♀, 10.i–1.ii.2015, DHJPAR0057451. 1♀, on *Solanum schlechtendallianum*, 11.01087°N, 85.48817°W, 400 m, 14.viii–4.ix.2009, DHJPAR0037886. 1♀, 10.91589°N, 85.26631°W, 420 m, 29.i–18.ii.2010, DHJPAR0038805. 1♀, on *Witheringia solanacea*, 10.94076°N, 85.3177°W, 461 m, 10–27.viii.2013, DHJPAR0053596. 1♀, from *Phostria* Janzen03 on *Psychotria berteriana*, 10.90425°N, 85.28651°W, 410 m, 10.x–1.xi.2003, DHJPAR0015351. 1♀, from *Phostria* Janzen05 on *Palicourea guianensis*, 11.01823°N, 85.45024°W, 410 m, 8.xi.2005–25.i.2006, DHJPAR0029183. 1♀, 10.93548°N, 85.25314°W, 123 m, 12–28.ii.2013, DHJPAR0051668. 1♀, 10.91847°N, 85.30338°W, 320 m, 10.xi–6.xii.2014, DHJPAR0057446. 1♀, 10.96543°N, 85.32043°W, 383 m, 8–26.IX.2015, DHJPAR0058044. 2♀, from *Rhectocraspeda* Janzen347 on *Solanum rovirosanum*, 10.91847°N, 85.30338°W, 320 m, 11–28.xi.2013, DHJPAR0054491, DHJPAR0054492. 1♀, from *Rhectocraspeda* Solis05 on *Solanum hayesii*, 10.89678°N, 85.270010°W, 420 m, 12.i–9.ii.2015, DHJPAR0057450. 1♀, on *Solanum rovirosanum*, 10.944040°N, 85.31738°W, 455 m, 13.x–4.xi.2014, DHJPAR0057445. 1♀, from *Pilocrocis purpurascens* on *Pentagonia donnell-smithii*, 10.94076°N, 85.3177°W, 461 m, 5–19.xi.2011, DHJPAR0046753. 1♀, from *Pleuroptya* Solis03 on *Laportea aestuans*, 10.96187°N, 85.28045°W, 96 m, 28.vi–16.vii.2013, DHJPAR0052682. 1♀, from spiloBioLep01 BioLep403 on *Mikania cordifolia*, 11.01983°N, 85.41342°W, 445 m, 12–31.viii.2011, DHJPAR0045803. 1♀, 12.viii–1.ix.2011, DHJPAR0045805 (EMUS, HIC).

Alabagrus janzeni Sharkey, 1988

Figure 36

Alabagrus janzeni Sharkey, 1988:373. Holotype ♀, Costa Rica (EMUS)

DIAGNOSIS. Gena right angled or acute posteroventrally. Third tergum with deep transverse depression. Forefemur pale. First tergum long and narrow with well-defined median longitudinal carina extending most of length.

DESCRIPTION. Body length 8.9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; with well-defined median longitudinal carina. Third tergum with deep transverse depression. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum variable but usually areolate; hind femur sometimes with some weak rugose sculpture.

HOST INFORMATION. *Eulepte concordalis*, *Eulepte* Janzen06, *Eulepte* Janzen02, *Eulepte* Solis15.

MATERIAL EXAMINED. Reared specimens: 1♀, from *Eulepte* Janzen06 on *Handroanthus impetiginosus*, 10.83764°N, 85.61871°W, 295 m, 20.vii–3.viii.1994, DHJPAR0015482. 1♀, 3–19.x.2000, 00-SRNP-18596. 1♂, 3–25.x.2000, 00-SRNP-18607. 2♂, 10.76828°N, 85.42567°W, 480 m, 8–25.x.2010, DHJPAR0041197, DHJPAR0041198. 1♂, 10.76828°N, 85.42567°W, 480 m, 8–26.x.2010, DHJPAR0041195. 1♂, 10.76737°N, 85.43313°W, 325 m, 27.xi–12.xii.2010, DHJPAR0041166. 2♀, 1♂, 27.xi–30.xii.2010, DHJPAR0041576. 3♀, 10.76828°N, 85.42567°W, 480 m, 28.vii–




Figure 36 *Alabagrus janzeni*, A–H females; I, J males: A. lateral habitus, B. dorsal habitus, C. lateral habitus, D. wings, E. lateral mesosoma, F. lateral head, G. dorsal head, H. anterior head, I. lateral habitus, J. propodeum and terga 1–3.

11.viii.2011, DHJPAR0044992, DHJPAR0044993, DHJPAR0044994. 1♀, 1♂, 28.vii–12.viii.2011, DHJPAR0044995, DHJPAR0044996. 1♀, 28.vii–13.viii.2011, DHJPAR0044997. 1♂, 10.76261°N, 85.42979°W, 330 m, 5–23.xi.2012, DHJPAR0050927. 1♀, 4–20.x.2013, DHJPAR0053588. 1♂, 4–23.x.2013, DHJPAR0053589. 1♀, 4–24.x.2013, DHJPAR0053622. 2♀, 4–26.x.2013, DHJPAR0053590, DHJPAR0054487. 1♂, from *Eulepte concordalis* on *Spathodea campanulata*, 10.9163°N, 85.37869°W, 460 m, 10–23.iii.2014, DHJPAR0055089. 2♀, 10–26.iii.2014, DHJPAR0055090, DHJPAR0055096. 1♀, 1♂, 10–27.iii.2014, DHJPAR0055093, DHJPAR0055095. 2♀, 10–28.iii.2014, DHJPAR0055092, DHJPAR0055094. 1♂, 10–30.iii.2014, DHJPAR0055091. 1♀, 10–31.iii.2014, DHJPAR0055097. 1♂, from *E. concordalis* on *Tabebuia rosea*, 10.74769°N, 85.58577°W, 90 m, 3–19.viii.2000, DHJPAR0015499. 1♂, 3–20.viii.2000, DHJPAR0015498. 1♂, 10.75725°N, 85.55354°W, 80 m, 10.viii–1.ix.2006, DHJPAR0010510. 1♂, 10.viii–26.ix.2006, 06-SRNP-18511. 1♂, 10–28.viii.2006, DHJPAR0010509. 1♀, 13.v–1.vi.2006, DHJPAR0010083. 1♀, 10.76261°N, 85.42979°W, 330 m, 22.ix–10.x.2013, DHJPAR0053587. 1♂, 22.ix–11.x.2013, DHJPAR0053623. 1♀, 10.76271°N, 85.56004°W, 95 m, 12.viii–1.ix.2009, DHJPAR0036363. 1♂, 12–27.viii.2008, DHJPAR0036365. 1♀, 12–29.viii.2009, DHJPAR0036366. 1♀, 10.76737°N, 85.43313°W, 325 m, 21.ii–19.iii.2005, DHJPAR0009400. 2♀, 24.ii–14.iii.2005, DHJPAR0009374, DHJPAR0009375. 1♀, 24.ii–17.iii.2005, DHJPAR0009403. 1♂, 25.i–19.ii.2005, DHJPAR0009376. 1♀, 10.76828°N, 85.42567°W, 480 m, 7–26.viii.2011, DHJPAR0044998. 1♂, 10.77074°N, 85.42874°W, 365 m, 25.vii–11.viii.2012, DHJPAR0049940. 1♂, 10.77175°N, 85.434°W, 305 m, 29.xi–17.xii.2012, DHJPAR0050926. 1♀, 10.80274°N, 85.67423°W, 10 m, 2–18.viii.2000, DHJPAR0015496. 1♂, 2–19.viii.2000, DHJPAR0015497. 1♂, 10.84389°N, 85.61384°W, 300 m, 5–20.viii.1982, DHJPAR0015481.

1♀, 10.85827°N, 85.61089°W, 280 m, 21.i–9.ii.2007, DHJPAR0016930. 1♀, 10.86333°N, 85.57443°W, 205 m, 25.viii–12.ix.2006, DHJPAR0016415. 1♂, 25.viii–8.ix.2006, DHJPAR0016417. 1♂, 25.viii–9.ix.2006, DHJPAR0016416. 2♀, 10.87124°N, 85.38749°W, 700 m, 10–27.ix.2006, DHJPAR0016429, DHJPAR0016432. 1♂, 10–29.ix.2006, DHJPAR0016428. 1♂, 10.8962°N, 85.27769°W, 430 m, 7–31.xii.2004, DHJPAR0009348. 1♂, 10.89678°N, 85.27001°W, 420 m, 26.ix–12.x.2012, DHJPAR0050363. 1♀, 10.90187°N, 85.27495°W, 415 m, 11.xi–2.xii.2005, DHJPAR0009434. 1♀, 11.xi–4.xii.2005, DHJPAR0009433. 1♂, 11–28.xi.2005, DHJPAR0015513. 1♂, 11–29.xi.2005, DHJPAR0015511. 2♀, 19.ix–6.x.2006, DHJPAR0016430, DHJPAR0016431. 1♀, 19.ix–8.x.2006, DHJPAR0021208. 1♀, 20.iv–7.v.2013, DHJPAR0052083. 1♂, 10.90383°N, 85.25964°W, 383 m, 31.x–20.xi.2005, DHJPAR0015515. 1♂, 31.x–21.xi.2005, DHJPAR0015514. 1♂, 31.x–23.xi.2005, DHJPAR0015512. 1♂, 31.x–26.xi.2005, DHJPAR0029179. 2♂, 10.9305°N, 85.37223°W, 527 m, 26.viii–10.ix.2009, DHJPAR0036712, DHJPAR0037190. 1♀, 26.viii–11.ix.2009, DHJPAR0036697. 1♂, 26.viii–18.ix.2009, DHJPAR0036698. 1♂, 10.93319°N, 85.25335°W, 95 m, 16.xi–4.xii.2006, DHJPAR0016926. 1♂, 21.x–7.xi.2006, DHJPAR0016928. 1♀, 10.93548°N, 85.25314°W, 123 m, 2–24.ix.2009, DHJPAR0036699. 1♀, 10.95991°N, 85.28298°W, 160 m, 25.viii–10.ix.2009, DHJPAR0037922. 1♀, 25.viii–12.ix.2009, DHJPAR0037902. 1♂, 10.96331°N, 85.32243°W, 434 m, 13–27.xii.2015, DHJPAR0058537. 1♀, 13–29.xii.2015, DHJPAR0058542. 1♂, 13–30.xii.2015, DHJPAR0058539. 2♂, 10.97853°N, 85.54628°W, 280 m, 13.viii–1.ix.1994, DHJPAR0015488, DHJPAR0015490. 2♀, 13.viii–4.ix.1994, DHJPAR0015487, DHJPAR0015489. 2♂, 13–28.viii.1994, DHJPAR0015485, DHJPAR0015486. 1♀, 13–29.viii.1994, DHJPAR0015484. 1♂, 13–31.viii.1994, DHJPAR0015483. 1♂,




Figure 37 *Alabagrus jeanfrancoislandryi* n. sp., A–J paratypes; K holotype: A. anterior head, B. lateral habitus, C. dorsal head, D. dorsal habitus, E. lateral mesosoma, F. lateral head and mesosoma, G. propodeum and tergum 1, H. wings, I. hind leg, J. hind leg, K. forewing.

11.00217°N, 85.46337°W, 560 m, 13–30.ix.2004, DHJPAR0015381. 1 ♀, 11.02681°N, 85.49547°W, 290 m, 10–23.ix.2004, DHJPAR0015383. 1 ♂, 10–25.ix.2004, DHJPAR0015384. 2 ♀, 10–28.ix.2004, DHJPAR0015380, DHJPAR0015382. 1 ♀, 10–29.ix.2004, DHJPAR0015379. 1 ♀, 1–20.xi.2004, DHJPAR0009362. 1 ♀, 1–23.xi.2004, DHJPAR0015315. 1 ♀, 25.x–10.xi.2004, DHJPAR0009354. 1 ♀, 25.x–12.xi.2004, DHJPAR0009355. 1 ♀, 25.x–9.xi.2004, DHJPAR0009356. 1 ♀, 1 ♂, 11.03376°N, 85.47715°W, 290 m, 22.vi–8.vii.2009, DHJPAR0036716, DHJPAR0036718. 1 ♂, 11.04562°N, 85.45742°W, 280 m, 6–18.vii.2007, DHJPAR0021211. 1 ♀, 6–19.vii.2007, DHJPAR0021149. 1 ♀, 10.94076°N, 85.3177°W, 461 m, 8–23.iii.2012, DHJPAR0050371. 1 ♀, from *Eulepte* Janzen02 on *Cydista heterophylla*, 10.77594°N, 85.65799°W, 7 m, 27.ix–13.x.2000, DHJPAR0015385. 1 ♀, 2–21.x.2000, DHJPAR0015386. 1 ♂, 10.77175°N, 85.434°W, 305 m, 25.x–18.xi.2010, DHJPAR0041167. 1 ♂, from *Eulepte* Janzen06 on *Hadroanthus guayacan*, 10.93548°N, 85.25314°W, 123 m, 18.x–4.xi.2012, DHJPAR0050923. 1 ♀, 23.x–10.xi.2012, DHJPAR0050928. 1 ♂, from *Eulepte* Solis15, 11.03226°N, 85.52776°W, 290 m, 23.xi–10.xii.2011, DHJPAR0048724. 1 ♀, 1 ♂, from *Eulepte* Solis15 on *Hadroanthus ochraceus*, 10.85145°N, 85.60801°W, 290 m, 23.x–12.xi.1998, DHJPAR0015492, DHJPAR0015493. 1 ♂, 23.x–20.xi.1998, DHJPAR0015491. 1 ♀, 23.x–15.xii.1998, DHJPAR0015494. 1 ♂, 23.x–17.xii.1998, DHJPAR0015495 (EMUS, HIC).

***Alabagrus jeanfrancoislandryi* Sharkey n. sp.**

Figure 37

DIAGNOSIS. Difficult or impossible to distinguish morphologically from *A. isidrochaconi* and *A. jennyphillipsae*. Gena rounded or with an obtuse angle posteroventrally. Hind femur heavily sculptured ventrally

with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum weakly convex or with a short, rounded longitudinal bulge. Hind coxa mostly or entirely black in lateral view. Pronotum melanic. Metapleuron entirely orange. Metasomal terga all orange.

DESCRIPTION. Body length 6.5 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral half or more. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. As in females.

HOST INFORMATION. *Ategumia lotanalis*, *Ategumia lotanalis*DHJ07, *Ategumia lotanalis*DHJ09, *Ategumia matutinalis*, *Bicilia iarchasalis*, *Conchylodes arcifera*, *Conchylodes ovulalis*, *Conchylodes platinalis*, *Desmia* BioLep22DHJ02, *Eulepte alialis*, *Eulepte concordalis*, *Eulepte* Janzen02, *Herpetogramma phaeopteralis*, *Herpetogramma salbialis*, *Herpetogramma semilaniata*, *Herpetogramma* Solis11, *Hyalorista exuvialis*DHJ02, *Microthyris anormalis*, *Palpusia* Janzen02, *Phaedropsis cernalis*, *Phostria samealis*, *Rhectocraspeda* Solis05, *Pilocrocis ramentalis*, *Pleuroptya* Solis03, *Pleuroptya* Solis04, *Salbia cassidalis*, *Salbia haemorrhoidalis*.

ETYMOLOGY. Named in honor of Dr. Jean-Francois Landry of the Canada Department of Agriculture, Ottawa, in recognition of his contributions to the taxonomy and systematics of ACG microlepidoptera.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Herpetogramma salbialis* on *Zexmenia virgulta*, 10.99637°N, 85.40195°W, 510 m, 7–20.ix.2009, DHJPAR0036683 (EMUS). PARATYPES: 1♂, from *Ategumia lotanalis* on *Conostegia xalapensis*, 10.99697°N, 85.39666°W, 470 m, 22.x–17.xi.2008, DHJPAR0030388. 1♀, on *Conostegia subcrustulata*, 10.95991°N, 85.28298°W, 160 m, 25.ii–19.iii.2013, DHJPAR0051670. 1♂, from *Ategumia lotanalis* DHJ07 on *Aciotis caulialata*, 10.94404°N, 85.31738°W, 455 m, 11–18.i.2013, DHJPAR0050933. 1♀, 11–27.i.2013, DHJPAR0050934. 1♀, from *A. lotanalis* DHJ07 on *Arthrostemma ciliatum*, 11.01983°N, 85.41342°W, 445 m, 21.ix–15.x.2006, DHJPAR0015387. 1♀, from *A. lotanalis* DHJ09 on *Conostegia xalapensis*, 10.93548°N, 85.25314°W, 123 m, 23.vii–7.viii.2009, DHJPAR0036356. 1♂, from *Ategumia matutinalis* on *Tibouchina longifolia*, 10.95991°N, 85.28298°W, 160 m, 21.ii–10.iii.2013, DHJPAR0051671. 1♀, from *Bicilia iarchasalis* on *Petiveria alliacea*, 10.78506°N, 85.6637°W, 10 m, 7–28.vii.1990, DHJPAR0015465. 1♀, from *Conchylodes arcifera* on *Verbesina gigantea*, 11.03004°N, 85.52699°W, 280 m, 8–22.ix.2000, DHJPAR0015332. 1♂, from *Conchylodes arcifera*, 11.03306°N, 85.42876°W, 400 m, 30.vii–6.viii.2005, DHJPAR0009406. 1♂, 30.vii–7.viii.2005, DHJPAR0015524. 1♀, 30.vii–9.viii.2005, DHJPAR0015521. 1♂, from *Conchylodes ovulalis* on *Baltimora recta*, 10.84867°N, 85.63046°W, 285 m, 12.viii–3.ix.1992, DHJPAR0015322. 1♀, 12.viii–6.ix.1992, DHJPAR0015320. 1♂, 12–28.viii.1992, DHJPAR0015321. 1♂, from *Conchylodes platinalis* on *Varronia guanacastensis*, 10.83764°N, 85.61871°W, 295 m, 19.vi–5.vii.2009, DHJPAR0036353. 1♂, on *Verbesina turbacensis*, 11.00891°N, 85.40977°W, 500 m, 7.ii–26.v.2007, DHJPAR0021157. 1♀, from *Desmia* BioLep22DHJ02 on *Begonia plebeja*, 10.94404°N, 85.31738°W, 455 m, 11–24.xi.2014, DHJPAR0057436. 1♀, from *Eulepte alialis* on *Clibadium pittieri*, 10.89928°N, 85.27486°W, 420 m, 10.xi–1.xii.2009, DHJPAR0037950. 1♀, on *Elephantopus angustifolius*, 10.77074°N, 85.42874°W, 365 m, 10–26.vii.2012, DHJPAR0049939. 1♀, on *Elephantopus mollis*, 10.77079°N, 85.37422°W, 733 m, 14.vi–3.vii.2011, DHJPAR0045003. 1♂, 21.vii–11.viii.2011, DHJPAR0044999. 1♂, 21.vii–7.viii.2011, DHJPAR0045000. 1♀, 22.vii–11.viii.2009, DHJPAR0036349. 1♂, 10.89678°N, 85.27001°W, 420 m, 24.vi–18.vii.2013, DHJPAR0052693. 1♀, 6.ix–1.x.2012, DHJPAR0050065. 1♂, 10.89928°N, 85.27486°W, 420 m, 10.xi–7.xii.2009, DHJPAR0037948. 1♂, 10.90427°N, 85.3073°W, 373 m, 5.vi–2.vii.2013, DHJPAR0052689. 1♂, 10.90528°N, 85.27882°W, 405 m, 17.vii–10.viii.2014, DHJPAR0057738. 1♂, 17.vii–7.viii.2014, DHJPAR0057745. 2♂, 27.vi–24.vii.2014, DHJPAR0055989, DHJPAR0055990. 1♂, 10.9061°N, 85.28281°W, 400 m, 31.vii–16.viii.2009, DHJPAR0036374. 1♀, 10.91847°N, 85.30338°W, 320 m, 7.x–7.xi.2009, DHJPAR0037917. 1♂, 10.94076°N, 85.3177°W, 461 m, 1–25.xi.2011, DHJPAR0046741. 1♀, 16.x–3.xi.2014, DHJPAR0057435. 1♂, 25.vii–7.viii.2014, DHJPAR0057005. 1♀, 25.vii–8.viii.2014, DHJPAR0056298. 1♀, 10.94404°N, 85.31738°W, 455 m, 3–21.x.2011, DHJPAR0045786. 1♀, 25.ix–13.x.2014, DHJPAR0057432. 1♀, 10.96952°N, 85.38258°W, 440 m, 4–23.x.2012, DHJPAR0050932. 1♂, 10.99102°N, 85.39539°W, 475 m, 20.ix–17.x.2010, DHJPAR0041196. 1♀, 10.99616°N, 85.45562°W, 560 m, 16.x–6.xi.2004, DHJPAR0009357. 1♂, 10.99697°N, 85.39666°W, 470 m, 18.ix–7.x.2009, DHJPAR0037187. 1♀, 29.x–20.xi.2009, DHJPAR0037956. 1♀, 11.0006°N, 85.438°W, 620 m, 8.x–1.xi.2004, DHJPAR0009373. 1♂, 8–28.x.2004, DHJPAR0009353. 1♀, on *Elephantopus spicatus*, 10.90353°N, 85.30916°W, 360 m, 12.ix–4.x.2009, DHJPAR0036703. 1♀, 12–25.ix.2009, DHJPAR0036717. 1♂, 10.9305°N, 85.37223°W, 527 m, 8.x–4.xi.2014, DHJPAR0057001. 1♀, 8–25.x.2014,

DHJPAR0056990. 1♂, on *Spiracantha cornifolia*, 10.90187°N, 85.27495°W, 415 m, 29.ix–14.x.2009, DHJPAR0036685. 1♂, 10.93548°N, 85.25314°W, 123 m, 23.viii–3.ix.2014, DHJPAR0056296. 1♂, from *Eulepte concordalis* on *Tabebuia rosea*, 11.02681°N, 85.49547°W, 290 m, 1–20.xi.2004, DHJPAR0015316. 1♀, from *Eulepte* Janzen02 on *Cydista heterophylla*, 10.85827°N, 85.61089°W, 280 m, 16.viii–1.ix.1990, 90-SRNP-1985. 1♂, from *Herpetogramma phaeopteralis* on *Axonopus fissifolius*, 10.76737°N, 85.43313°W, 325 m, 18–30.v.2011, DHJPAR0045006. 1♂, from *Herpetogramma salbialis* on *Clibadium leiocarpum*, 10.77175°N, 85.434°W, 305 m, 22.v–6.vi.2011, DHJPAR0045015. 1♀, 2♂, 22.v–7.vi.2011, DHJPAR0045004, DHJPAR0045010, DHJPAR0045012. 1♀, on *Zexmenia virgulta*, 10.90661°N, 85.28784°W, 400 m, 30.iv–12.v.2014, DHJPAR0055349. 1♀, on *Eleutheranthera ruderalis*, 10.90528°N, 85.27882°W, 405 m, 25.xi–19.xii.2014, DHJPAR0056999. 1♀, 25.xi–26.xii.2014, DHJPAR0056994. 1♀, on *Melanthera aspera*, 10.93548°N, 85.25314°W, 123 m, 26.viii–6.ix.2014, DHJPAR0056295. 1♀, on *Melanthera nivea*, 10.77175°N, 85.434°W, 305 m, 11–25.v.2011, DHJPAR0042824. 1♀, 27.v–10.vi.2011, DHJPAR0045011. 2♂, 27.v–7.vi.2011, DHJPAR0045005, DHJPAR0045014. 1♀, 30.v–10.vi.2011, DHJPAR0045013. 1♂, 10.96187°N, 85.28045°W, 96 m, 11–20.x.2014, DHJPAR0057430. 1♀, on *Zexmenia virgulta*, 10.99637°N, 85.40195°W, 510 m, 7–18.ix.2009, DHJPAR0037951. 1♀, 1♂, from *Herpetogramma semilaniata* on *Justicia* 14522, 10.77594°N, 85.65799°W, 7 m, 24.ix–6.x.2000, DHJPAR0015329, DHJPAR0015330. 1♀, 27.ix–21.x.2000, DHJPAR0015331. 1♀, from *Herpetogramma* Solis11 on *Solanum hayesii*, 10.86472°N, 85.41531°W, 540 m, 27.viii–12.ix.2011, DHJPAR0045783. 1♀, 11.04118°N, 85.4417°W, 320 m, 8–30.ix.2010, DHJPAR0041603. 1♀, from *Hyalorista exuvialis* DHJ02 on *Hyptis obtusiflora*, 10.94076°N, 85.3177°W, 461 m, 16.ix–9.x.2009, DHJPAR0037900. 1♀, from *Microthyris anormalis* on *Merremia umbellata*, 10.88996°N, 85.47966°W, 550 m, 22.vi–10.vii.2005, DHJPAR0015522. 1♀, from *Palpusia* Janzen02 on *Ipomoea batatas*, 10.93332°N, 85.25331°W, 135 m, 21.vii–6.viii.2013, DHJPAR0052680. 1♀, from *Phaedropsis cernalis* on *Triplaris melaenodendron*, 10.78938°N, 85.55098°W, 85 m, 26.viii–10.ix.2000, DHJPAR0015327. 1♀, 26.viii–14.ix.2000, DHJPAR0015326. 1♂, 26.viii–7.ix.2000, DHJPAR0015328. 1♂, from *Phostria samealis* on *Tetracera hydrophila*, 10.93332°N, 85.25331°W, 135 m, 8–24.ix.2009, DHJPAR0036700. 1♀, from *Rhectocraspeda* Solis05 on *Solanum hayesii*, 10.93548°N, 85.25314°W, 123 m, 15–25.vii.2014, DHJPAR0055979. 1♂, on *Solanum torvum*, 10.9301°N, 85.25205°W, 109 m, 2–14.x.2012, DHJPAR0050931. 1♀, from *Pilocrocis ramentalis* on *Dyschoriste quadrangularis*, 10.78626°N, 85.55835°W, 105 m, 2–17.ix.1994, DHJPAR0015324. 1♀, 10.83764°N, 85.61871°W, 295 m, 29.vi–15.vii.1993, DHJPAR0015318. 1♂, on *Ruellia inundata*, 10.7416°N, 85.42734°W, 420 m, 14–31.viii.2013, DHJPAR0053650. 1♀, from *Pleuroptya* Solis03 on *Laportea aestuans*, 10.96187°N, 85.28045°W, 96 m, 26.iii–9.iv.2012, DHJPAR0050370. 1♂, 11.01823°N, 85.45024°W, 410 m, 2–23.iii.2009, DHJPAR0036696. 1♀, 2–21.iii.2009, DHJPAR0036695. 1♀, 11.00519°N, 85.47398°W, 490 m, 16–31.viii.2002, DHJPAR0015333. 1♂, 11.02865°N, 85.48669°W, 280 m, 20.viii–5.ix.2004, DHJPAR0015334. 1♀, from *Pleuroptya* Solis04 on *Cecropia peltata*, 10.83525°N, 85.62485°W, 285 m, 18.ix–5.x.2008, DHJPAR0028307. 1♀, on *Myriocarpa longipes*, 11.02865°N, 85.48669°W, 280 m, 25.ix–15.x.2004, DHJPAR0015335. 1♂, from *Salbia cassidalis* on *Lasiacis sorghoidea*, 10.83764°N, 85.61871°W, 295 m, 3–27.xi.1995, DHJPAR0015319. 1♂, from *Salbia haemorrhoidalis* on *Lantana camara*, 10.92291°N, 85.31877°W, 410 m, 9–20.x.2014, DHJPAR0057431. 1♀,




Figure 38 *Alabagrus jeanmariecadioui* n. sp., holotype: A. anterior head, B. lateral habitus, C. dorsal head, D. dorsal habitus, E. wings, F. lateral head and mesosoma, G. propodeum and tergum 1.

10.93332°N, 85.25331°W, 135 m, 6–21.x.2009, DHJPAR0037953. 1 ♀, 10.95991°N, 85.28298°W, 160 m, 6–21.viii.2013, DHJPAR0053613. 1 ♀, 10.918470°N, 85.30338°W, 320 m, 8–28.x.2014, DHJPAR0056995.

***Alabagrus jeanmariecadioui* Sharkey n. sp.**

Figure 38

DIAGNOSIS. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Third tergum lacking transverse depression, or depression barely indicated. Propodeum completely smooth. Forefemur yellow. Metapleuron entirely melanic. Hind coxa mostly melanic laterally.

NOTES. Remarkably consistent in color; no males. May be *Alabagrus fuscovittatus* Enderlein, which was collected in “Central America.”

DESCRIPTION. Body length 8.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Desmia ploralis*DHJ03, *Desmia ploralis*DHJ01, *Desmia ploralis*DHJ02, *Desmia* Janzen09, *Desmia octomaculalis*, *Desmia* Solis19, *Desmia ploralis*DHJ04, *Desmia ploralis*DHJ06, *Desmia* Solis100.

ETYMOLOGY. Named in honor of Dr. Jean-Marie Cadiou (deceased) of Belgium in recognition of his inspirational contributions to the taxonomy and systematics of ACG Sphingidae.

MATERIAL EXAMINED. HOLOTYPE ♀, from *D. ploralis*DHJ03 on *Psychotria chagrensis*, 10.90037°N, 85.37254°W, 500 m, 12.xi–2.xii.2009, DHJPAR0037934 (EMUS). PARATYPES: 1 ♀, from *Desmia* Janzen09 on *Psychotria chagrensis*, 10.87621°N, 85.38632°W, 670 m, 13–30.vi.2005, DHJPAR0009424. 1 ♀, 10.90014°N, 85.38284°W, 495 m, 19.vii–12.viii.2005, DHJPAR0015508. 1 ♀, on *Psychotria jimenezii*, 10.94076°N, 85.3177°W, 461 m, 20.ix–14.x.2012, DHJPAR0050925. 1 ♀, from *Desmia octomaculalis* on *Psychotria chagrensis*, 10.87766°N, 85.39343°W, 645 m, 21.ii–29.iii.2013, DHJPAR0053592. 1 ♀, from *Desmia ploralis*DHJ01 on *Coccocypselum herbaceum*, 10.94076°N, 85.3177°W, 461 m, 27.viii–10.ix.2011, DHJPAR0045806. 1 ♀, 11.00311°N, 85.42041°W, 580 m, 10–29.ix.2011, DHJPAR0045796. 1 ♀, 10–29.ix.2011, DHJPAR0045807. 1 ♀, 13.x–14.xi.2011, DHJPAR0046752. 1 ♀, 13.x–25.xi.2011, DHJPAR0051346. 1 ♀, 13.x–8.xi.2011, DHJPAR0046751. 1 ♀, from *D. ploralis*DHJ02 on *Psychotria horizontalis*, 11.02681°N, 85.49547°W, 290 m, 24.vii–18.viii.2005, DHJPAR0015507. 1 ♀, from *D. ploralis*DHJ03 on *Psychotria chagrensis*, 10.89021°N, 85.38803°W, 520 m, 10.v–3.vi.2012, DHJPAR0048716. 1 ♀, 22.ix–15.x.2009, DHJPAR0036687. 1 ♀, 10.89296°N, 85.3788°W, 520 m, 30.ix–22.x.2009, DHJPAR0036710. 1 ♀, 10.89678°N, 85.27001°W, 420 m, 26.v–17.vi.2005, DHJPAR0009427. 1 ♀, 10.90037°N, 85.37254°W, 500 m, 11.viii–2.ix.2009, DHJPAR0036704. 1 ♀, 22.vi–14.vii.2014, DHJPAR0055969. 2 ♀, 10.90445°N, 85.28412°W, 400 m, 10.vi–2.vii.2005, DHJPAR0009416, DHJPAR0009429. 2 ♀, 10.vi–4.vii.2005, DHJPAR0009428, DHJPAR0009430. 1 ♀, 10.90528°N, 85.27882°W, 405 m, 29.viii–26.ix.2011, DHJPAR0045797. 1 ♀, 10.9305°N, 85.37223°W, 527 m, 10.v–2.vi.2012, DHJPAR0048717. 1 ♀, 10.94076°N, 85.3177°W, 461 m, 1–23.vii.2011, DHJPAR0044989. 1 ♀, 10.940760°N, 85.3177°W, 461 m, 4–24.xi.2014, DHJPAR0057444. 1 ♀, 10.98705°N,




Figure 39 *Alabagrus jennyphillipsae* n. sp., A–G holotype; H, I paratypes: A. anterior head, B. lateral habitus, C. dorsal head, D. dorsal habitus, E. wings, F. lateral head and mesosoma, G. propodeum and tergum 1, H. hind leg, I. hind leg.

85.42816°W, 700 m, 22.vi–15.vii.2006, DHJPAR0015509. 1♀, 10.9888°N, 85.42336°W, 680 m, 10.xii.2012–5.i.2013, DHJPAR0050922. 1♀, 15.vii–4.viii.2006, DHJPAR0010513. 1♀, 10.98931°N, 85.42581°W, 675 m, 11.vii–11.viii.2003, DHJPAR0015377. 2♀, 11.vii–8.viii.2003, DHJPAR0015376, DHJPAR0015378. 2♀, on *Psychotria graciliflora*, 10.88068°N, 85.26968°W, 540 m, 26.viii–14.ix.2010, DHJPAR0041193, DHJPAR0041206. 1♀, 10.88111°N, 85.38889°W, 570 m, 11–28.vi.2013, DHJPAR0052690. 1♀, 23.vi–13.vii.2013, DHJPAR0052691. 1♀, 10.89666°N, 85.29003°W, 400 m, 10–30.vi.2008, DHJPAR0028030. 1♀, 16.v–6.vi.2012, DHJPAR0048730. 1♀, 7–30.v.2013, DHJPAR0052084. 1♀, 10.89678°N, 85.27001°W, 420 m, 13.vi–2.vii.2012, DHJPAR0049656. 1♀, 10.89928°N, 85.27486°W, 420 m, 24.v–17.vi.2005, DHJPAR0009425. 1♀, 10.90187°N, 85.27495°W, 415 m, 1–20.vii.2011, DHJPAR0044985. 1♀, 1–25.vii.2011, DHJPAR0044987. 1♀, 10.9305°N, 85.37223°W, 527 m, 17.vi–9.viii.2013, DHJPAR0052695. 1♀, from *D. ploralis* DHJ04 on *Coccocypselum herbaceum*, 10.89928°N, 85.27486°W, 420 m, 16.x–5.xi.2009, DHJPAR0037918. 1♀, 10.91589°N, 85.26631°W, 420 m, 17.ii–12.iii.2010, DHJPAR0038804. 1♀, 10.9163°N, 85.37869°W, 460 m, 3–22.xi.2009, DHJPAR0037954. 1♀, from *Desmia ploralis* DHJ06 on *Psychotria lamarinensis*, 11.01926°N, 85.40997°W, 440 m, 22.i–22.ii.2008, DHJPAR0023534. 1♀, from *Desmia Solis*100 on *Ludwigia foliobracteolata*, 10.90187°N, 85.27495°W, 415 m, 11.ix–2.x.2009, DHJPAR0036702. 1♀, 10.90528°N, 85.27882°W, 405 m, 30.xi–21.xii.2012, DHJPAR0050921. 1♀, 10.94741°N, 85.31501°W, 491 m, 12–30.vii.2010, DHJPAR0041190. 1♀, 2–20.v.2010, DHJPAR0041204. 1♀, 10.96187°N, 85.28045°W, 96 m, 12.x–5.xi.2009, DHJPAR0037901. 1♀, 13–31.viii.2009, DHJPAR0037905. 1♀, 17.viii–1.ix.2009, DHJPAR0037921. 1♀,

17.viii–2.ix.2009, DHJPAR0037904. 1♀, 17.viii–8.ix.2009, DHJPAR0037903. 1♀, from *Desmia Solis*19 on *Psychotria panamensis*, 10.94741°N, 85.31501°W, 491 m, 21.x–5.xi.2009, DHJPAR0037893 (EMUS, HIC).

***Alabagrus jennyphillipsae* Sharkey n. sp.**

Figure 39

DIAGNOSIS. Difficult or impossible to distinguish morphologically from *A. isidrochaconi* and *A. jeanfrancoislandryi*. Gena rounded or with an obtuse angle posteroventrally. Head less elongate than *A. arua*. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Forewing entirely infusate, not clear or yellow at base. Pronotum melanic. Propodeum orange. Terga orange except apical terga melanic. First tergum varying from weakly convex to with a short rounded longitudinal bulge.

NOTES. Specimens of this species were included under *A. albispina* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 7.2 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less, or areolate rugose in ventral half or more. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.




Figure 40 *Alabagrus jimmilleri* n. sp., A–G holotype; I paratype male: A. lateral habitus, B. dorsal head, C. anterior head, D. dorsal habitus, E. wings, F. ventral hind femur, G. propodeum and tergum 1, H. lateral head and mesosoma, I. propodeum and tergum 1.

HOST INFORMATION. *Leucochromodes melusinalis*DHJ01.

ETYMOLOGY. Named in honor of Dr. Jenny Phillips of INBio and the Museo Nacional de Costa Rica, who has long and masterfully taxonomized the ACG Pyraloidea and Tortricoidea inventory, as well as for all of Costa Rica.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Leucochromodes melusinalis*DHJ01 on *Mandevilla hirsuta*, 10.93332°N, 85.25331°W, 135 m, 19.v–1.vi.2009, DHJPAR0035225 (EMUS). PARATYPES: reared: 1♀, *Leucochromodes melusinalis*DHJ01 on *Tassadia obovata*, 10.99697°N, 85.39666°W, 470 m, 30.xi–21.xii.2009, DHJPAR0037923 (EMUS). 1♀, on *Tassadia obovata*, 10.99637°N, 85.40195°W, 510 m, 23.iii–11.iv.2010, DHJPAR0039523 (HIC).

Alabagrus jimmilleri Sharkey n. sp.

Figure 40

DIAGNOSIS. Gena acute posteroventrally. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum varying from weakly convex (most females) to with a rounded longitudinal bulge (most males). Body black and red to black and orange. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Scutellum melanic. Metapleuron and metasoma entirely orange.

NOTES. Specimens of this species were included under *Alabagrus combos* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 5.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.4× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum with more sculpture; metasomal median tergite 1 with sharper median longitudinal bulge.

HOST INFORMATION. *Lygropia tripunctata*DHJ04, *Microthyris anormalis*DHJ02, *Microthyris prolongalis*, *Microthyris prolongalis*DHJ02, *Palpusia* Janzen02.

ETYMOLOGY. Named in honor of Dr. Jim Miller of New York City and the American Museum of Natural History, who has long and masterfully taxonomized the ACG Notodontidae inventory, as well as for all of Costa Rica.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Lygropia tripunctata*DHJ04 on *Ipomoea phillomega*, 10.93332°N, 85.25331°W, 135 m, 24.vi–7.vii.2009, DHJPAR0036373 (EMUS). PARATYPES: reared: 1♀, from *Microthyris anormalis*DHJ02 on *Ipomoea batatas*, 10.86472°N, 85.41531°W, 540 m, 20.vii–9.viii.2012, DHJPAR0049946. 1♂, from *Microthyris prolongalis* on *Ipomoea phillomega*, 10.90661°N, 85.28784°W, 400 m, 25.v–26.vi.2010, DHJPAR0040337. 1♂, 10.90037°N, 85.37254°W, 500 m, 3–18.vi.2012, DHJPAR0049046. 1♂, 10.90093°N, 85.28915°W, 400 m, 4–21.vi.2014, DHJPAR0055343. 1♂, from *M. prolongalis*DHJ02 on *Merremia tuberosa*, 11.01602°N, 85.38053°W, 380 m, 23.vii–9.viii.2012, DHJPAR0049938. 1♂, from *Palpusia* Janzen02 on *Turbina corymbosa*, 11.02865°N, 85.48669°W, 280 m, 15.xi–5.xii.2004, DHJPAR0009372. Nonreared paratypes: 4♀, 2♂, from Limón and Puntarenas Provinces (HIC, INBio).

Alabagrus johnbrowni Sharkey n. sp.

Figure 41

DIAGNOSIS. Gena acute posteroventrally. Mesoscutum yellow. Precoxal sulcus distinct and foveolate about ½ length of mesopleuron. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Forefemur black. Hind coxa with a black patch laterally. Forewing yellow in basal ⅓ gradually becoming infusate apically. Costal vein melanic.

NOTES. Specimens of this species were included under *A. cocto* and *A. yaruro* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 8.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures




Figure 41 *Alabagrus johnbrowni* n. sp., A, B, E holotype; C, D paratypes: A. lateral habitus, color variation, B. wings, C. lateral habitus, color variation, D. lateral habitus, E. dorsal habitus.

ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor short, not longer than metasoma, or longer than metasoma but not longer than body.

Males. Propodeum with more sculpture.

Variation. Body color varies considerably; thorax varying from almost completely melanic to almost completely yellow; forewing may have a less distinct border between the yellow and melanic colors shown in Figure 41D.

HOST INFORMATION. *Desmia* BioLep09, *Desmia* BioLep19, *Desmia* BioLep42, *Desmia tages*, *Lamprosema* Dapkey23, *Phostria* Janzen05, *Piletosoma thialis*, *Piletosoma thialis*DHJ03.

ETYMOLOGY. Named in honor of Dr. John Brown of the Systematic Entomology Laboratory in the U.S. National Museum of Natural History, the taxonomist who has long and masterfully taxonomized the ACG Tortricoidea inventory, as well as for all of the Neotropics.

MATERIAL EXAMINED. HOLOTYPE ♀, from *Piletosoma thialis* on *Doliocarpus multiflorus*, 10.89678°N, 85.27001°W, 420 m, 1–20.vi.2009, DHJPAR0035226 (EMUS). PARATYPES: 1♀, 4♂, from *Desmia* BioLep09 on *Cissus alata*, 10.87741°N, 85.32363°W, 560 m, 11.vii–2.viii.2011, DHJPAR0045060, DHJPAR0045058, DHJPAR0045059, DHJPAR0045057, DHJPAR0045061. 1♀, from *Desmia* BioLep19 on *Hamelia patens*, 10.9305°N, 85.37223°W, 527 m, 22.viii–10.ix.2014, DHJPAR0056357. 1♀, 11.04249°N, 85.40339°W, 390 m, 8–27.viii.2014, DHJPAR0056297. 1♂, from *Desmia* BioLep42

on *Vitis tiliifolia*, 10.89678°N, 85.27001°W, 420 m, 16.vii–7.viii.2014, DHJPAR0055980. 1♀, 10.905280°N, 85.278820°W, 405 m, 22.vii–13.viii.2014, DHJPAR0057743. 1♀, from *Desmia tages* on *Hamelia patens*, 10.87741°N, 85.32363°W, 560 m, 1–21.x.2010, DHJPAR0041591. 1♂, 10.9305°N, 85.37223°W, 527 m, 11.x–1.xi.2011, DHJPAR0045782. 1♂, 11.03234°N, 85.43954°W, 380 m, 30.i–23.ii.2005, DHJPAR0009378. 1♂, from *Lamprosema* Dapkey23 on *Serjania mexicana*, 10.9301°N, 85.25205°W, 109 m, 25.ii–13.iii.2013, DHJPAR0051669. 1♂, from *Phostria* Janzen05 on *Psychotria cyanococca*, 10.90187°N, 85.27495°W, 415 m, 24.viii–9.ix.2009, DHJPAR0036707. 1♀, from *Piletosoma thialis* on *Doliocarpus multiflorus*, 10.89678°N, 85.27001°W, 420 m, 22.ii–15.iii.2013, DHJPAR0051667. 1♀, 28.v–14.vi.2014, DHJPAR0055346. 1♀, 6.xi–13.xii.2013, DHJPAR0054531. 1♂, 9.vi–1.vii.2009, DHJPAR0036335. 1♀, 9.vi–7.vii.2009, DHJPAR0036336. 1♀, 9.vi–9.vii.2009, DHJPAR0035954. 1♂, 1–22.vii.2009, DHJPAR0036375. 1♂, 10.91847°N, 85.30338°W, 320 m, 19.iii–12.iv.2013, DHJPAR0052086. 1♀, 10.940760°N, 85.3177°W, 461 m, 28.ix–14.x.2014, DHJPAR0057418. 1♂, 28.ix–15.x.2014, DHJPAR0057417. 1♂, 10.94741°N, 85.31501°W, 491 m, 11–26.vi.2011, DHJPAR0045029. 1♂, 12.x–11.xi.2011, DHJPAR0046749. 1♀, 12.x–12.xi.2011, DHJPAR0046755. 1♀, 1♂, 12.x–8.xi.2011, DHJPAR0046748, DHJPAR0046754. 1♀, 10.99637°N, 85.40195°W, 510 m, 24.vi–20.vii.2009, DHJPAR0040069. 1♀, from *P. thialis*DHJ03, 10.90424°N, 85.2712°W, 410 m, 21.ii–21.iii.2006, DHJPAR0009364. 1♀, 21.ii–

23.iii.2006, DHJPAR0015517. 1 ♀, 2–29.xi.2005, DHJPAR0015516. 1 ♀, 8–27.vii.2005, DHJPAR0009407. 1 ♀, 11.01926°N, 85.40997°W, 440 m, 22.ii–18.iii.2007, DHJPAR0017277 (EMUS, HIC).

Alabagrus johnburnsi Sharkey n. sp.

Figure 42

DIAGNOSIS. Female: Shape of occiput dorsolaterally rounded. Propodeum completely smooth. Metasoma entirely yellow. Pronotum mesoscutum and scutellum melanic, propodeum melanic but slightly paler than mesonotum. Forewing yellow in basal $\frac{2}{3}$ gradually becoming infusate apically. Costal vein melanic. Hind coxa pale with a large melanic patch laterally. Male: Precoxal sulcus with one or several foveae posteroventrally, with or without a smooth groove extending anteriorly. Mesoscutum melanic. Propodeum weakly areolated. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum with a rounded longitudinal bulge extending more than $\frac{1}{2}$ length of tergum. Body color black and yellow. Third tergum lacking transverse depression, or depression barely indicated.

DESCRIPTION. Body length 8.8 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Some propodea areolate; precoxal sulcus always foveolate.

HOST INFORMATION. Unknown.

ETYMOLOGY. Named in honor of Dr. John M. Burns of the U.S. National Museum of Natural History in the Smithsonian Institution, the taxonomist who has long and masterfully taxonomized all of the ACG Hesperoidea as well as that for North America.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, La Cruz, Estación Pitilla, 9 km S Santa Cecilia, 700 m, vi.1996, C. Moranga, P. Rios, Malaise, L_N_330200_380200, #47562, H10616 (INBio). PARATYPES: 1 ♂, 11 ♀, from the following provinces: Guanacaste, Alajuela, San Jose, Puntarenas, altitudes vary between 560 and 1040 m (HIC, INBio, MUCR, EMUS).

Alabagrus johnbryckii Sharkey n. sp.

Figure 43

DIAGNOSIS. Female: Metapleuron mostly or entirely melanic. Metasoma entirely orange. Ovipositor distinctly longer than body. First tergum weakly convex. Propodeum smooth. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Occiput rounded dorsolaterally. Male: First tergum with well-defined median longitudinal carina. Propodeum with more defined and complete sculpture.

NOTES. Specimens of this species were included under *A. englishi* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 6.3 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several




Figure 42 *Alabagrus johnburnsi* n. sp., A–C holotype; D, E paratype male: A. wings, B. lateral habitus, C. dorsal habitus, D. forewing, E. propodeum and tergum 1.

distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. First tergum with well-defined median longitudinal carina; propodeum with more defined and complete sculpture; third tergum lacking transverse depression, or depression barely indicated.

ETYMOLOGY. Named in honor of Dr. John Obrycki, former chair of the Department of Entomology, University of Kentucky, for his many years of support to M.J.S.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, 3 km SE R. Naranjo, 14–20.viii.1993, F.D. Parker (EMUS). PARATYPES: 1 ♀, Alajuela, 20 km S Upala, 22–30.ix.1991, F. D. Parker (HIC). 1 ♂, Cartago, PN Tapantí, Macizo de la Muerte, Torre circa del Est., 1200 m, 9.vi.2000, L_N_193800_560050, H10967 (INBio).

Alabagrus johnsharkeyi Sharkey n. sp.

Figure 44

DIAGNOSIS. Gena right angled posteroventrally. Median tergite 1 red, tergite 2 red and black, remaining terga black. Hind femur with




Figure 43 *Alabagrus johnobryckii* n. sp., holotype: **A.** lateral habitus, **B.** wings, **C.** dorsal habitus.

weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Median tergite 1 about as wide as long. Hind femur, metapleuron and pronotum all entirely melanic.

NOTES. Specimens of this species were included under *A. semialbus* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 8.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $0.9\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of John Sharkey, youngest brother of M.J.S., and lawyer extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, 3 km SE R. Naranjo, 22–24.vii.1992, F.D. Parker, H17276 (EMUS).

Alabagrus kaciejoae Sharkey n. sp.

Figure 45

DIAGNOSIS. First tergum with two converging longitudinal carinae, more pronounced in males. Propodeum with weak irregular ridges. Pronotum melanic.

NOTES. Specimens of this species were included under *A. parvifaciatus* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 8.9 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; with two converging longitudinal carinae. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. First tergum with more pronounced longitudinal carinae.

ETYMOLOGY. Named in honor of Kacie Athey, former graduate student of M.J.S.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, 700 m, 9 km S Santa Cecilia, set 1989, UTM 330200_380200, GNP Biodiversity Study, H10955 (INBio). PARATYPES: 1 ♀, 2 ♂, Guanacaste and Alajuela (INBio, EMUS, HIC).

Alabagrus karensharkeyae Sharkey n. sp.

Figure 46

DIAGNOSIS. Precoxal sulcus with several large foveae posteroventrally. First tergum narrow basally, widening rapidly at midlength, with well-defined median longitudinal carina extending most of its length. Pronotum melanic. Metapleuron entirely pale. Hind femur mostly or entirely melanic. Forewing mostly infusate with pale patches that do not form discrete bands. Apical metasomal terga melanic.

NOTES. Specimens of this species were included under *A. maya* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 7.1 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.4\times$ longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated, or with deep transverse depression. Ovipositor longer than metasoma but not longer than body.

Males. Similar to female.

HOST INFORMATION. *Microthyris prolongalis*, *Microthyris prolongalis*DHJ02, *Palpusia* Janzen02, *Psara obscuralis*DHJ02.




Figure 44 *Alabagrus johnsharkeyi* n. sp., holotype: A. lateral habitus, B. dorsal habitus, C. lateral head, D. lateral mesosoma, E. propodeum and tergum 1, F. wings.

ETYMOLOGY. Named in honor of my (M.J.S.) beautiful wife, Karen.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Microthyris prolongalis* on *Ipomoea phillomega*, 10.9301°N, 85.25205°W, 109 m,

17.v–3.vi.2010, DHJPAR0041200 (EMUS). PARATYPES: reared: 1♂, from *Microthyris prolongalis* on *Ipomoea batatas*, 10.87766°N, 85.39343°W, 645 m, 2–17.viii.2013, DHJPAR0052904. 1♀, 10.90093°N, 85.28915°W, 400 m, 23.v–10.vi.2006, DHJPAR0010196. 1♀, 10.90661°N, 85.28784°W, 400 m, 11–27.vi.2007, DHJPAR0021170. 1♀, 17.viii–14.ix.2005, DHJPAR0015531. 1♀, 8–29.v.2013, DHJPAR0052079. 1♀, 10.9305°N, 85.37223°W, 527 m, 13–31.vii.2009, DHJPAR0036348. 1♀, 10.93332°N, 85.25331°W, 135 m, 27.viii–19.ix.2009, DHJPAR0036706. 1♀, on *Ipomoea lindenii*, 9–24.viii.2009, DHJPAR0036715. 1♀, 10.9305°N, 85.37223°W, 527 m, 23.viii–10.ix.2012, DHJPAR0050362. 1♂, from *M. prolongalis* DHJ02 on *Ipomoea trifida*, 10.77175°N, 85.434°W, 305 m, 27.v–15.vi.2011, DHJPAR0045002. 1♂, from *Palpusia* Janzen02 on *Ipomoea batatas*, 10.93332°N, 85.25331°W, 135 m, 24.xi–10.xii.2011, DHJPAR0046742. 1♀, 10.93548°N, 85.25314°W, 123 m, 5–19.vii.2014, DHJPAR0055978. 1♀, from *Psara obscuralis* DHJ02 on *Ipomoea phillomega*, 10.9163°N, 85.37869°W, 460 m, 10–31.iii.2014, DHJPAR0055102. Nonreared paratypes: 4♂, 17♀, 0–200 m, Limón, Alajuela, Puntarenas, Guanacaste (EMUS, HIC, INBio, EMUS). MEXICO: Veracruz (CNCI).

Alabagrus kaydodgeae Sharkey n. sp.

Figure 47




Figure 45 *Alabagrus kaciejoe* n. sp., A–D holotype; E paratype male: A. lateral habitus, B. wings, C. dorsal habitus, D. propodeum and tergum 1, E. propodeum and tergum 1.

DIAGNOSIS. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression, usually posteriorly, crenulae always lacking. Head pale except vertex melanic, or melanic except margins of gena yellow, or melanic except margins of gena yellow and yellow patch on face. Mesoscutellum black contrasting with the otherwise yellow mesosoma.

NOTES. Formerly considered as *A. mojos* (Leathers and Sharkey, 2003). Male color variable; almost all have some black, at least near the superior orbits of the eyes.

DESCRIPTION. Body length 6.6 mm. Gena right angled or acute posteroventrally. Precoxal sulcus absent or almost absent, represented at




Figure 46 *Alabagrus karensharkeyae* n. sp., holotype: **A.** lateral habitus, **B.** anterior head, **C.** dorsal head, **D.** dorsal habitus, **E.** wings, **F.** lateral head and mesosoma, **G.** propodeum and tergum 1.

most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or

no more heavily sculptured than dorsal surface. First tergum 1.36× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.




Figure 47 *Alabagrus kaydodgeae* n. sp., **A–C, E–H** paratype females; **D, I, J** paratype males: **A.** lateral habitus, **B.** anterior head, **C.** dorsal habitus, **D.** lateral habitus, **E.** wings, **F.** dorsal head, **G.** lateral head, **H.** propodeum, **I.** wings, **J.** anterior head.




Figure 48 *Alabagrus keithwillmotti* n. sp., A–F, H holotype; G, I paratype male: A. lateral habitus, B. ventral hind femur, C. anterior head, D. dorsal head, E. dorsal habitus, F. wings, G. forewing, H. propodeum and tergum 1, I. propodeum and tergum 1.

Males. Color variable; forewing varies from banded, as in the female, to mostly infusate with a weak pale band at midlength that does not include the stigma, to completely infusate; head usually mostly yellow with vertex and upper occiput melanic.

HOST INFORMATION. *Ategumia* Solis01.

ETYMOLOGY. Named in honor of Ms. Kay Dodge of Santa Cruz, Guanacaste, Costa Rica, in recognition of her patient and very long term efforts to support the growth and evolution of ACG since its beginnings in 1986.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, 11 km SE La Virgen, 10.33°N, 84.07°W, 450–550 m, 17.iv.2003, H1200 (INBio). PARATYPES: reared: 1♂, from *Ategumia* Solis01 on *Psychotria alfaroana*, 11.02367°N, 85.41884°W, 375 m, 31.v–25.vi.2013, DHJPAR0053007. 1♀, 11.019259°N, 85.40997°W, 440 m, host caterpillar, same as above, host plant=*Psychotria grandis*, collected 10.xi.2014, DHJPAR0057442 (EMUS). Nonreared paratypes: 44♀, 39♂, COSTA RICA, Limón, Puntarenas, Alujuela, Colón, San Jose; PANAMA, Colón, Canal Zone. (EMUS, HIC, INBio).

***Alabagrus keithwillmotti* Sharkey n. sp.**

Figure 48

DIAGNOSIS. Gena rounded posteroventrally. Propodeum melanic but slightly paler than remainder of mesosoma. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. First tergum weakly convex. Third tergum lacking transverse depression, or depression barely indicated. Metasoma yellow except apical tergum somewhat melanic.

DESCRIPTION. Body length 9.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures

ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing darker, lacking most pale patches found in the female; propodeum with more sculpture; first metasomal median tergite wider.

HOST INFORMATION. *Asturodes fimbriauralis* DHJ02, *Desmia* Janzen07, *Desmia* Janzen14, *Desmia octomaculalis*, *Herpetogramma* Solis10, *Phostria* Janzen03, *Pilocrocis* Solis20, *spilobiolep01* biolep498.

ETYMOLOGY. Named in honor of Dr. Keith Willmott of the McGuire Center for Lepidoptera and Biodiversity of the University of Florida in recognition of his contributions to the taxonomy and systematics of Neotropical butterflies and their biodiversity.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Pilocrocis* Solis20 on *Saurauia montana*, 10.92918°N, 85.46426°W, 1220 m, 23.x–25.xi.2004, DHJPAR0009389 (EMUS). 1♂, from *Asturodes fimbriauralis* DHJ02 on *Colubrina spinosa*, 10.98816°N, 85.39581°W, 485 m, 4–26.iii.2010, DHJPAR0038809. 1♂, 10.99697°N, 85.39666°W, 470 m, 2–30.vi.2009, DHJPAR0036339. 1♀, 11.01983°N, 85.41342°W, 445 m, 21.vii–19.viii.2003, DHJPAR0015336. 3♂, from *Desmia* Janzen07 on *Psychotria microbotrys*, 10.98758°N, 85.41967°W, 680 m, 23.xi–21.xii.2009, DHJPAR0037909, DHJPAR0037910, DHJPAR0037911. 1♂, 23.xi–23.xii.2009, DHJPAR0037913. 1♂, 23.xi–24.xii.2009, DHJPAR0037907. 1♀, 12.x–7.xi.2009, DHJPAR0037912. 1♂, 27.vi–17.vii.2013, DHJPAR0052730. 1♀, from *Desmia* Janzen14 on *Hoffmannia longipetiolata*, 10.98332°N, 85.43623°W, 900 m, 22.xii.2012–28.i.2013, DHJPAR0050935. 1♀, 3.i–11.ii.2013, DHJPAR0050930. 1♀, 3.i–8.ii.2013, DHJPAR0050868. 1♂, from *Desmia octomaculalis* on *Notopleura*




Figure 49 *Alabagrus leedyeri* n. sp., holotype: A. lateral head, B. dorsal head, C. lateral habitus, D. anterior head, E. dorsal habitus, F. forewing, G. lateral mesosoma, H. scutellum and propodeum.

uliginosa, 10.86472°N, 85.41531°W, 540 m, 13.v–6.vi.2010, DHJPAR0039357. 3♂, from *Herpetogramma Solis10* on *Justicia macrantha*, 10.99637°N, 85.40195°W, 510 m, 7–29.ix.2011, DHJPAR0045789, DHJPAR0045808, DHJPAR0045809. 1♂, from *Phostria* Janzen03 on *Psychotria elata*, 10.98758°N, 85.41967°W, 680 m, 16.ii–12.iii.2010, DHJPAR0038808. 2♀, from *Pilocrocis Solis20* on *Saurauia montana*, 10.92918°N, 85.46426°W, 1220 m, 23.x–19.xi.2004, DHJPAR0009387, DHJPAR0009388. 1♂, 23.x–23.xi.2004, DHJPAR0009386. 3♀, 23.x–25.xi.2004, DHJPAR0009385, DHJPAR0009389, DHJPAR0009390. 1♀, 24.vii–21.viii.2000, 00-SRNP-10070. 1♀, 24.vii–24.viii.2000, 00-SRNP-10067. 1♀, 24.vii–25.viii.2000, 00-SRNP-10080. 1♀, 24.vii–26.viii.2000, 00-SRNP-10087. 1♂, 10.99616°N, 85.45562°W, 560 m, 14.x–6.xi.2004, DHJPAR0007207. 1♀, from *spilobiolep01* biolep498 on *Faramaea stenura*, 10.98758°N, 85.41967°W, 680 m, 9.v–14.vi.2009, DHJPAR0035227 (EMUS, HIC).

***Alabagrus leedyeri* Sharkey n. sp.**

Figure 49

DIAGNOSIS. Gena acute posteroventrally. First tergum longer than wide and weakly convex. Propodeum melanic and areolate, with at least one closed areola. All femora mostly or entirely melanic.

NOTES. Formerly considered as part of *A. tricarinatus* (Leathers and Sharkey, 2003).

DESCRIPTION. Body length 7.1 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly,

or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. In honor of Lee Dyer, entomologist and ecologist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, caterpillar feeding on *Byttneria aculeata*, Heredia, La Selva Biol. Sta., 3 km S Puerto Viejo, 10.43°N, 84.02°W, Coll. 27.iii.1980, eclosed 7.iv.1980, H10613 (HIC). PARATYPE: 1♀, Ala(juela), 20 km S Upala (10.84°N, 85.07°W), 24.i.1991, F.D. Parker (EMUS).

***Alabagrus lindapitkiniae* Sharkey n. sp.**

Figure 50

DIAGNOSIS. Gena right angled or acute posteroventrally. Precoxal sulcus distinct with large foveae extending about $\frac{1}{2}$ length of mesopleuron. Third tergum lacking transverse depression, or depression barely indicated. Metapleuron entirely orange. Midfemur black except pale apex. Hind femur black basally, more so dorsally, and with a black




Figure 50 *Alabagrus lindapitkiniae* n. sp., holotype: **A.** lateral habitus, **B.** wings, **C.** lateral head and mesosoma, **D.** dorsal habitus, **E.** propodeum and tergum 1.

patch apicodorsally. Terga 1–2 orange and about as long as wide, tergum 3 orange anteriorly and black posteriorly, remaining terga black.

DESCRIPTION. Body length 8.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Omiodes* Janzen05.

ETYMOLOGY. Named in honor of Dr. Linda Pitkin of The Natural History Museum, London, in recognition of her very fruitful labors for the parataxonomists, taxonomy, and systematics of Geometridae of ACG and the world.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Omiodes* Janzen05 on *Entada gigas*, 10.9555°N, 85.28381°W, 221 m, 4–26.iv.2013, DHJPAR0052082 (EMUS). PARATYPE: 1♀, *Omiodes* Janzen05 on *Entada gigas*, 10.96187°N, 85.28045°W, 96 m, 1–25.xi.2013, DHJPAR0054497 (EMUS).

Alabagrus longinoi Sharkey n. sp.

Figure 51

DIAGNOSIS. Third tergum lacking transverse depression, or depression barely indicated. Propodeum lacking complete areolae, usually mostly or completely smooth. Ovipositor longer than metasoma but not longer than body. All femora yellow. Pronotum melanic. Metapleuron entirely yellow; hind coxa almost entirely or entirely black. Apical yellow band of forewing starting near apex of stigma and complete to posterior margin.

DESCRIPTION. Body length 6.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.6\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of John (Jack) Longino, myrmecologist extraordinaire, for his support during years of research in Costa Rica.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, 11 km SE La Virgen, 10.33°N, 84.07°W, 450–550 m, 23.iii.2003 (INBio). PARATYPES: 3♀, Heredia, Limón, 220–550 m, ii.2003 and x.1999 (INBio, HIC, EMUS).

Alabagrus maculipes (Cameron, 1887)

Figure 52

Microdus maculipes Cameron, 1887:404 holotype ♂, Guatemala (BMNH).

Microdus trochanteratus Cameron, 1905:50 holotype ♀, Nicaragua (BMNH)

Alabagrus maculipes (Sharkey, 1988:380)

DIAGNOSIS. Gena rounded posteroventrally. Head pale, or pale except vertex melanic. Forewing evenly infuscate or with some pale regions but always lacking distinct bands.

DESCRIPTION. Body length 5.8 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae




Figure 51 *Alabagrus longinói* n. sp., holotype: **A.** lateral habitus, **B.** dorsal habitus, **C.** Lateral head and mesosoma, **D.** wings, **E.** propodeum and tergum 1.

posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide;

varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body. **Males.** Forewing varies from as in the female to completely infuscate. **HOST INFORMATION.** *Desmia ploralis*DHJ02, *Herpetogramma* Janzen04, *Herpetogramma salbialis*, *Herpetogramma semilaniata*, *Massepha grammalis*, *Palpusia* Janzen02, *Rhectocraspeda* Solis05, *Pilocrocis ramentalis*, *Salbia haemorrhoidalis*, *Syngamia florella*.




Figure 52 *Alabagrus maculipes*: **A.** lateral habitus, **B.** dorsal head, **C.** anterior head, **D.** lateral head and mesosoma, **E.** propodeum and tergum 1, **F.** wings.




Figure 53 *Alabagrus malcolmscoblei* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus, D. lateral head, E. mesopleuron, F. propodeum and tergum 1.

DISTRIBUTION. Widespread from Mexico, the Caribbean south through Central America, and the Caribbean countries of South America (Sharkey, 1988).

MATERIAL EXAMINED. Reared specimens: 1 ♀, from *Desmia ploralis* DHJ02 on *Psychotria horizontalis*, 10.76712°N, 85.59663°W, 180 m, 11–22.viii.2009, DHJPAR0036337 (EMUS). 2 ♀, from *Herpetogramma* Janzen04 on *Achyranthes aspera*, 10.7416°N, 85.42734°W, 420 m, 12–30.xi.2010, DHJPAR0041150, DHJPAR0041151. 1 ♀, from *Herpetogramma salbialis* on *Delilia biflora*, 10.93548°N, 85.25314°W, 123 m, 8–18.ix.2014, DHJPAR0056292. 1 ♀, from *Herpetogramma semilaniata* on *Justicia* 14522, 10.77594°N, 85.65799°W, 7 m, 24.ix–9.x.2000, DHJPAR0015349. 1 ♀, on *Justicia carthaginensis*, 10.77079°N, 85.37422°W, 733 m, 17–29.vi.2010, DHJPAR0040329. 1 ♀, from *Massepha grammalis* on *Maranta arundinacea*, 10.85827°N, 85.61089°W, 280 m, host collection 21.vii.1990, DHJPAR0015346. 1 ♂, from *Palpusia* Janzen02 on *Ipomoea trifida*, 10.77079°N, 85.37422°W, 733 m, 28.vi–11.vii.2010, DHJPAR0040212. 1 ♂, 28.vi–12.vii.2010, DHJPAR0040213. 1 ♂, from *Rhectocraspeda* Solis05 on *Solanum jamaicense*, 10.95991°N, 85.28298°W, 160 m, 18–27.xii.2012, DHJPAR0050947. 1 ♀, on *Solanum aturense*, 10.93548°N, 85.25314°W, 123 m, 8–20.ix.2014, DHJPAR0056291. 1 ♀, from *Pilocrocis ramentalis* on *Dyschoriste quadrangularis*, 10.78626°N, 85.55835°W, 105 m, 2–12.ix.1994, 94-SRNP-7176. 1 ♀, 2–14.ix.1994, DHJPAR0015347. 1 ♀, 2–18.ix.1994, DHJPAR0015348. 11 ♀, host collection 2.ix.1994, 94-SRNP-7143, 94-SRNP-7144, 94-SRNP-7145, 94-SRNP-7146, 94-SRNP-7148, 94-SRNP-7160, 94-SRNP-7164, 94-SRNP-7165, 94-SRNP-7169, 94-SRNP-7173, 94-SRNP-7187. 1 ♂, 10.83764°N, 85.61871°W, 295 m, 7–16.vi.2012, DHJPAR0048735. 1 ♀, 7–19.vi.2012, DHJPAR0048736. 1 ♀, from *Salbia haemorrhoidalis* on *Lantana camara*, 10.93332°N, 85.25331°W, 135 m, 23.ix–4.x.2009, DHJPAR0037188. 1 ♀, from *Syngamia florella* on *Spermacoce exilis*,

11.0006°N, 85.438°W, 620 m, 5–26.ii.2009, DHJPAR0036725 (EMUS, HIC).

***Alabagrus malcolmscoblei* Sharkey n. sp.**

Figure 53

DIAGNOSIS. Gena right angled or acute posteroventrally. Propodeum lacking complete areolae, mostly smooth. First tergum weakly convex. Forewing lacking two distinct bands; pale basally to near level of apex of stigma with a darkening near parastigma. Mesoscutum, tegula, and forefemur melanic. Hind coxa orange.

DESCRIPTION. Body length 9.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Phostria latiapicalis*.

ETYMOLOGY. Named in honor of Dr. Malcolm Scoble of The Natural History Museum, London, in recognition of his very fruitful labors for the parataxonomists, taxonomy, and systematics of Geometridae and Hedylidae of ACG and the world.




Figure 54 *Alabagrus marcepsteini* n. sp., holotype: A. dorsal head, B. lateral habitus, C. anterior head, D. dorsal habitus, E. wings, F. lateral mesosoma, G. lateral head, H. propodeum and tergum 1.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phostria latipicalis* on *Chimarrhis parviflora*, 10.9305°N, 85.37223°W, 527 m, 10–29.v.2012, DHJPAR0048718 (EMUS).

Alabagrus marcepsteini Sharkey n. sp.

Figure 54

DIAGNOSIS. Gena right angled or acute posteroventrally. Precoxal sulcus with several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Forewing lacking two distinct bands; pale basally to near level of apex of stigma with a darkening near parastigma. Forefemur and metapleuron mostly or entirely melanic.




Figure 55 *Alabagrus mariaheikkilae* n. sp., holotype: A. lateral habitus, B. dorsal habitus, C. propodeum and tergum 1, D. wings.

Propodeum mostly or entirely orange. Hind coxa mostly orange with a vertical black stripe.

NOTES. Specimens of this species were included under *A. kagaba* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 9.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. crambidJanzen01 Janzen21.

ETYMOLOGY. Named in honor of Dr. Marc Epstein of California Department of Food and Agriculture, Sacramento, California, in recognition of his long labor at taxonomizing the Limacodidae of ACG and all of Costa Rica for their inventories.

MATERIAL EXAMINED. HOLOTYPE ♀, host=crambidJanzen01 Janzen21 on *Elaeagia auriculata*, 10.84886°N, 85.3281°W, 980 m, 17.viii–8.ix.2010, DHJPAR0041184 (EMUS). PARATYPES: reared: 1 ♀, from crambidJanzen01 Janzen21 on *Elaeagia auriculata*, 10.84886°N, 85.3281°W, 980 m, 17.viii–6.ix.2010, DHJPAR0041192 (ovipositor broken at base). Nonreared: 1 ♀, Limón, Valle de la Estrella, Res. Biol. Hitoy Cerere, Sendero Toma de Agua, 140 m, 17.vi–17.vii.2000, LN_643400_184600 (INBio).

Alabagrus mariaheikkilae Sharkey n. sp.

Figure 55

DIAGNOSIS. Gena acute posteroventrally; forewing yellow basally, infuscate in apical ¼ with a sharp margin between colors; metasoma entirely yellowish orange; propodeum with weak areolae.

DESCRIPTION. Body length 5.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly,




Figure 56 *Alabagrus markmetzi* n. sp., A–G holotype; H paratype male: A. anterior head, B. dorsal head, C. lateral habitus, D. dorsal habitus, E. lateral head and mesosoma, F. wings, G. propodeum and tergum 1, H. propodeum and tergum 1.

or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Salbia cassidalis* DHJ02.

ETYMOLOGY. Named in honor of Dr. Maria Heikkilä of the Systematic Entomology Laboratory in the U.S. National Museum of Natural History in recognition of her enthusiastic taxonomizing of ACG Gelechioidea for the inventory.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Salbia cassidalis* DHJ02 on *Lasiacis ruscifolia*, 10.9305°N, 85.37223°W, 527 m, 23.x–21.xi.2014, DHJPAR0057004 (EMUS). PARATYPE: 1 ♀, same data as holotype, except date 23.x–13.xi.2014, DHJPAR0057434 (HIC).

Alabagrus markmetzi Sharkey n. sp.

Figure 56

DIAGNOSIS. Gena rounded posteroventrally. Precoxal sulcus with several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Propodeum areolate. First tergum weakly convex. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Scutellum melanic. Body color black and orange to black and red.

NOTES. Specimens of this species were included under *A. roibasi* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 6.2 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse

punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeal sculpture more pronounced, areolae deeper.

HOST INFORMATION. *Syllepte aechmisalis*.

ETYMOLOGY. Named in honor of Dr. Mark Metz of the Systematic Entomology Laboratory in the U.S. National Museum of Natural History in recognition of his enthusiastic taxonomizing of ACG Gelechioidea for the inventory.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Syllepte aechmisalis* on *Triumfetta lappula*, 11.02991°N, 85.45364°W, 345 m, 12.x–12.xi.2004, DHJPAR0009359 (EMUS). PARATYPES: all from *Syllepte aechmisalis* on *Triumfetta lappula*. 1 ♀, 10.76737°N, 85.43313°W, 325 m, 17.vi–6.vii.2005, DHJPAR0009417. 1 ♀, 10.76901°N, 85.43465°W, 335 m, 10.viii–2.ix.2011, DHJPAR0046737. 1 ♂, 10.78938°N, 85.55098°W, 85 m, 22.viii–13.ix.2006, DHJPAR0015388. 1 ♀, 11.04788°N, 85.45266°W, 280 m, 28.x–21.xi.2004, DHJPAR0009363. 1 ♀, 10.78506°N, 85.6637°W, 10 m, 17–30.vii.1990, DHJPAR0015476. 1 ♀, 11.02364°N, 85.49139°W, 290 m, 26.ix–16.x.2004, DHJPAR0015368 (EMUS, HIC).

Alabagrus masneri Sharkey, 1988

Figure 57

Alabagrus masneri Sharkey, 1988:381–2. Holotype ♀, Panama (EMUS)

DIAGNOSIS. Female: Gena rounded posteroventrally. Propodeum lacking complete areolae, mostly or completely smooth. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Ovipositor about as long as body. Body length 4.01–5.79 mm., shorter than similar species. Head melanic. Mesoscutum yellow. Fore- and midcoxae both partly or entirely melanic. Hind femur mostly or entirely yellow. Male: Darker than females and with a well areolated propodeum and a distinct median carina on tergum 1. Hind coxa partly to entirely melanic. Pronotum pale, or melanic anteriorly and pale posteriorly. Precoxal sulcus with one or several distinct foveae posteroventrally, with




Figure 57 *Alabagrus masneri*, A, B, E female; C, D male: A. lateral habitus, B. dorsal habitus, C. lateral habitus, D. propodeum and tergum 1, E. wings.

or without a smooth groove extending anteriorly, or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron.

DESCRIPTION. Body length 4.9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.3\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Propodeum areolate. First tergum often with a median carina.

DISTRIBUTION. Widespread from southern Mexico to northern South America (Leathers and Sharkey, 2003).

Alabagrus mattottoi Sharkey n. sp.

Figure 58

DIAGNOSIS. Gena rounded posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Propodeum mostly smooth with a few irregular rugae anteromedially. Propleuron yellow. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Fore- and midcoxae entirely yellow.

NOTES. Specimens of this species were included under *A. imitatus* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 6.8 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae

posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.6\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Mr. Mathew Otto, golfer and friend extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, Los Almendros, P.N. Guanacaste A.C., 300 m, 3–22.viii.1993, L_N_334800_369800 (INBio).

Alabagrus miqa Sharkey, 1988

Figure 59

Alabagrus miqa Sharkey, 1988:384–5. Holotype ♀, Mexico (BMNH)

DIAGNOSIS. Tergum 1 wider than long with a distinct median longitudinal carina. Third tergum with deep transverse depression. Ovipositor short, not longer than metasoma. Forewing almost entirely infusate, ignore small clear patches posterad stigma.

DESCRIPTION. Body length 7.3 mm. Gena right angled or acute posteroventrally, or rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum




Figure 58 *Alabagrus mattotloi* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $0.8\times$ longer than wide; with well-defined median longitudinal carina. Third tergum with deep transverse depression. Ovipositor short, not longer than metasoma.

Males. Unknown.

DISTRIBUTION. Southern Mexico to Costa Rica (Leathers and Sharkey, 2003).

Alabagrus nickgrishini Sharkey n. sp.

Figure 60

DIAGNOSIS. Gena right acute posteroventrally. Propodeum completely smooth with a hint of smooth shallow areolation. Forewing lacking two distinct bands; pale basally to near level of apex of stigma with a darkening near parastigma. First tergum weakly convex. Mesoscutum melanic. Tegula yellow. Forefemur black. Hind coxa entirely reddish orange.




Figure 59 *Alabagrus miqa*: A. lateral habitus, B. dorsal habitus, C. wings.

NOTES. No specimens that are parasitoids of *Phostria* have *COI* data, but the morphology and color are good matches.

DESCRIPTION. Body length 7.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

Variation. Some specimens have a hint of a posterior median areola on the propodeum.




Figure 60 *Alabagrus nickgrishini* n. sp., holotype: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. wings, F. lateral head and mesosoma, G. propodeum and tergum 1.

HOST INFORMATION. *Microthyris alvinalis*, *Erilusa leucoplagialis*, *Phostria latiapicalis*, *Pleuroptya* Solis02.

ETYMOLOGY. Named in honor of Dr. Nick Grishin of the Howard Hughes Medical Institute, Dallas, Texas, in recognition of his enthusiastic taxonomizing of Neotropical Hesperidae and deep genetic dissection of ACG complexes.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Microthyris alvinalis* on *Banisteriopsis muricata*, 10.77079°N, 85.37422°W, 733 m, 6–24.viii.2009, DHJPAR0036347 (EMUS). PARATYPES: 1♀, from *Erilusa leucoplagialis* on *Calycophyllum candidissimum*, 10.76261°N, 85.429790°W, 330 m, 29.xii.2014–14.i.2015, DHJPAR0057425. 1♀, from *Microthyris alvinalis* on *Banisteriopsis muricata*, 10.83764°N, 85.61871°W, 295 m, 12–29.viii.2011, DHJPAR0045022. 1♀, 28.vi–12.vii.2000, DHJPAR0015345. 1♀, 11.01087°N, 85.48817°W, 400 m, 25.viii–14.ix.2009, DHJPAR0037892. 2♀, from *Phostria latiapicalis* on *Calycophyllum candidissimum*, 10.83575°N, 85.61253°W, 290 m, 17.xi–4.xii.1990, DHJPAR0015342, DHJPAR0015343. 1♀, 10.83575°N, 85.61253°W, 290 m, 17.xi–4.xii.1990, DHJPAR0015344. 1♀, from *Pleuroptya* Solis02 on *Guettarda macrosperma*, 10.77079°N, 85.37422°W, 733 m, 11–28.ix.2012, DHJPAR0050367. 1♀, 11–30.ix.2012, DHJPAR0050366 (EMUS, HIC).

Alabagrus nicoya Sharkey, 1988

Figure 61

Alabagrus nicoya Sharkey, 1988:388–9. Holotype ♀, Costa Rica (CNCI)

DIAGNOSIS. Hind femur wider than any other species and heavily rugose ventrally. First metasomal tergite weakly convex and wider than long. Metapleuron and median tergites 1–3 orange. Pronotum melanic. First tergum varying from weakly convex to with a rounded longitudinal bulge.

DESCRIPTION. Body length 9.3 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less, or areolate rugose in ventral half or more. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 0.7× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. As in females.

HOST INFORMATION. *Dichogama colotha*, *Dichogama diffusalis*, *Dichogama redtenbacheri*.

DISTRIBUTION. Southern Mexico to northwestern Costa Rica.

MATERIAL EXAMINED. Reared specimens: 1♀, from *Dichogama colotha* on *Capparis flexuosa*, 10.83575°N, 85.61253°W, 290 m, 30.vii–22.viii.2001, 01-SRNP-16138. 1♀, 18.vii–22.viii.2001, 01-SRNP-15782. 1♂, on *Capparis incana*, 10.81703°N, 85.64366°W, 270 m, 18.viii–14.ix.1993, 93-SRNP-4958. 1♂, 18.viii–20.ix.1993, 93-SRNP-4953. 1♀, 3.viii–13.ix.1992, DHJPAR0015396. 1♀, 3.viii–20.ix.1992, 92-SRNP-4222. 1♀, 3.viii–5.ix.1992, DHJPAR0015397. 1♂, 31.vii.1992–12.iii.1993, DHJPAR0015398. 1♀, 5.viii.1992–22.iii.1993, 92-SRNP-4242. 1♂, 5.viii–10.ix.1992, DHJPAR0015394. 1♀, 5.viii–29.ix.1992, DHJPAR0015395. 1♂, 5.viii–5.ix.1992, 92-SRNP-4230. 1♀, 7.viii–7.ix.1992, DHJPAR0015393. 1♂, on *Capparis indica*, 10.78938°N, 85.55098°W, 85 m, 12.viii–10.ix.2007, DHJPAR0022193. 1♂, 10.79342°N, 85.6666°W, 5 m, 10.v–5.vi.1996, 96-SRNP-1278. 1♂, 10.80274°N, 85.67423°W, 10 m, 15.vi–18.vii.1996, 96-SRNP-4969. 1♀, 10.8269°N, 85.60413°W, 240 m, 16.vii–25.viii.2001, 01-SRNP-15698. 1♂, 10.83575°N, 85.61253°W, 290 m, 11.vii–10.viii.1998, 98-SRNP-9501. 1♀,




Figure 61 *Alabagrus nicoya*: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. wings, F. hind leg, G. lateral mesosoma, H. propodeum and tergum 1.

11.vii–14.xii.1998, 98-SRNP-9492. 1♂, 14.viii.2002–2.ii.2003, 02-SRNP-13211. 1♀, 1♂, 14.viii–10.ix.2002, DHJPAR0015399, DHJPAR0015400. 1♀, 14.viii–11.ix.2002, 02-SRNP-13198. 1♂, 14.viii–12.ix.2002, 02-SRNP-13187. 1♀, 14.viii–13.ix.2002, 02-SRNP-13194. 1♀, 14.viii–26.xi.2002, 02-SRNP-13199. 1♀, 16.vi–26.vii.1989, 89-SRNP-204. 2♀, 1♂, 21.v–18.vi.2001, 01-SRNP-12510, 01-SRNP-12512, 01-SRNP-12513. 1♀, 21.v–26.vi.2001, 01-SRNP-12518. 1♂, 30.vii.2001–23.iii.2002, 01-SRNP-16233. 1♀, 30.vii–22.viii.2001, 01-SRNP-16203. 1♂, 30.vii–23.viii.2001, 01-SRNP-16173. 2♀, 1♂, 30.vii–25.viii.2001, 01-SRNP-16151, 01-SRNP-16211, 01-SRNP-16214. 1♂, 30.vii–27.viii.2001, 01-SRNP-16153. 1♀, 1♂, 30.vii–28.viii.2001, 01-SRNP-16254, 01-SRNP-16290. 1♀, 30.vii–29.viii.2001, 01-SRNP-16272. 1♂, 30.vii–30.viii.2001, 01-SRNP-16302. 1♂, 8.vii–10.viii.1979, 79-SRNP-211. 1♀, 10.84389°N, 85.61384°W, 300 m, 19.xii.2006–22.i.2007, DHJPAR0016931. 2♀, 21.v–12.vi.1993, DHJPAR0015392, 93-SRNP-831. 1♀, 7.vii–19.viii.1990, 90-SRNP-1194. 1♀, 9.vii–11.viii.2012, DHJPAR0049650. 1♀, 10.85413°N, 85.61534°W, 280 m, 13.vii–1.viii.1994, 94-SRNP-5606. 1♀, from *Dichogama diffusalis* on *Capparis discolor*, 11.03004°N, 85.52699°W, 280 m, 26.i–23.ii.2001, 01-SRNP-9101. 1♂, from *Dichogama redtenbacheri* on *Capparis frondosa*, 10.78506°N, 85.6637°W, 10 m, 17.vii–10.viii.1990, 90-SRNP-1407.2. 1♂, 17.vii–11.viii.1990, 90-SRNP-1407.3. 1♂, 17.vii–8.viii.1990, 90-SRNP-1407.1. 1♂, 7–27.vii.1990, 90-SRNP-1242.1. 1♀, 10.83381°N, 85.61271°W, 295 m, 12.vii–7.viii.2014, DHJPAR0055977. 1♂, 15.vii–3.viii.2014, DHJPAR0055972. 1♀, 15.vii–7.viii.2014, DHJPAR0055976. 1♀, 10.83575°N, 85.61253°W, 290 m, 12.viii–4.ix.1990, 90-SRNP-1955. 1♀, 12.viii–4.ix.1990, 90-

SRNP-1955. 1♂, on *Morisonia americana*, 10.82224°N, 85.64338°W, 270 m, 11.viii–4.ix.1993, DHJPAR0015391 (EMUS, HIC).

***Alabagrus patsharkeyi* Sharkey n. sp.**

Figure 62

DIAGNOSIS. Gena rounded posteroventrally. Body length more than 7 mm, and ovipositor clearly longer than body. Precoxal sulcus distinct and foveolate about ½ length of mesopleuron. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Propleuron yellow, or melanic and yellow. Hind coxa yellow except extreme apex melanic. Propodeum with weak rugose sculpture medially. Fore- and midcoxae both partly or entirely melanic.

NOTES. Specimens of this species were included under *A. imitatus* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 8.7 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.4× longer than wide; varying from weakly convex to with a rounded longitudinal bulge.




Figure 62 *Alabagrus patsharkeyi* n. sp., holotype: A. ventral hind femur, B. lateral habitus, C. wings, D. dorsal habitus.

Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Patrick Sharkey, second youngest brother of M.J.S. and great fighter.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, ACG, Los Almendros, 3–22.viii.1993, 300 m, LN 334800 369800 (INBio). PARATYPES: 1 ♀, same data as holotype (HIC). 4 ♀, Guanacaste, ~4 km S Cañas, ix–xii, 1990–1991, F. Parker (EMUS, EMUS, HIC). 1 ♀, Limón, Sector Cocori, 30 km N Cariari, 100 m, ii.1995, LN 286000 567500 (INBio).

Alabagrus paulgoldsteini Sharkey n. sp.

Figure 63

DIAGNOSIS. Gena acute posteroventrally. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Third tergum with transverse depression barely indicated. Most of dorsal surface of hind femur melanic, most of ventral surface pale. Terga 1–3 orange, remaining terga melanic. Hind femur melanic dorsally, pale ventrally. Forewing yellow basally, infusate apically, with a melanic patch posterad parastigma. Propodeum orange, contrasting with melanic metapleuron.

DESCRIPTION. Body length 13 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Syllepte amando* DHJ02.




Figure 63 *Alabagrus paulgoldsteini* n. sp., holotype: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. wings, F. propodeum and tergum 1, G. lateral head and mesosoma.




Figure 64 *Alabagrus paulheberti* n. sp., holotype: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. lateral head and mesosoma, F. wings, G. propodeum and tergum 1.

ETYMOLOGY. Named in honor of Dr. Paul Goldstein of the Systematic Entomology Laboratory in the U.S. National Museum of Natural History in recognition of his diligent and enthusiastic taxonomizing of ACG Noctuoidea for the inventory.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Syllepte amando* DHJ02 on *Banara guianensis*, 11.01525°N, 85.39766°W, 415 m, 28.xi.2009–8.i.2010, DHJPAR0037908 (EMUS).

Alabagrus paulheberti Sharkey n. sp.

Figure 64

DIAGNOSIS. Ovipositor short, not longer than metasoma. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Pronotum melanic anteriorly and ventrally, pale posteriorly and dorsally.

DESCRIPTION. Body length 8.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor short, not longer than metasoma.

Males. Unknown.

HOST INFORMATION. *Aponia itzalis*, *Aponia minnithalis*.

ETYMOLOGY. Named in honor of Dr. Paul Hebert of the Centre for Biodiversity Genomics of the University of Guelph in recognition of his essential invention and evolution of DNA barcoding, without which the ACG Lepidoptera and parasitoid inventory would be impossible.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Aponia itzalis* on *Cornutia grandifolia*, 10.76737°N, 85.43313°W, 325 m, 20.vi–9.vii.2010, DHJPAR0040223 (EMUS). PARATYPES: 1 ♀, from *Aponia itzalis* on *Cornutia grandifolia*, 10.85827°N, 85.61089°W, 280 m, 30.vi–26.vii.1989, DHJPAR0015389. 1 ♀, from *Aponia minnithalis*

on *Cornutia grandifolia*, 10.87741°N, 85.32363°W, 560 m, 5.xi–1.xii.2011, DHJPAR0046953. 1 ♀, 5–27.xi.2011, DHJPAR0046746 (EMUS, HIC).

Alabagrus paulsharkeyi Sharkey n. sp.

Figure 65

DIAGNOSIS. Forefemur yellow. Pronotum and scutellum melanic. Metasoma entirely yellow.

NOTES. Specimens of this species were included under *A. latreillei* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 6.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Paul Sharkey, older brother of M.J.S. and surgeon extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, 50–150 m, Est. Biol. La Selva, 10.43°N, 84.02°W, 10.vii.2000 (INBio).

Alabagrus paulthiaucourti Sharkey n. sp.

Figure 66

DIAGNOSIS. Gena rounded posteroventrally. First tergum with well-defined median longitudinal carina in females and with several longitudinal carinae and rugosities in male. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more




Figure 65 *Alabagrus paulsharkeyi* n. sp., holotype: **A.** lateral habitus, **B.** Dorsal habitus, **C.** wings, **D.** Lateral head and mesosoma, **E.** propodeum and tergum 1.

heavily sculptured than dorsal surface. Pronotum melanic. All terga except apical (which is intermediate or variable) orange.

DESCRIPTION. Body length 9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.4\times$ longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely

indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Male with more propodeal sculpture; first tergum with rugosostriate sculpture laterally.

HOST INFORMATION. *Diatraea* Janzen989.

ETYMOLOGY. Named in honor of Dr. Paul Thiaucourt of Paris, France, in recognition of his long and fruitful taxonomizing of the Notodontidae of the world, with special attention to the hundreds found by the ACG inventory.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Diatraea* Janzen989 on *Paspalum orbiculatum*, 10.9301°N, 85.25205°W, 109 m,




Figure 66 *Alabagrus paulthiaucourti* n. sp., **A–D, F, G** holotype; **E** paratype male: **A.** lateral habitus, **B.** anterior head, **C.** dorsal head, **D.** propodeum and tergum 1, **E.** propodeum and tergum 1, **F.** wings, **G.** Lateral head and mesosoma.




Figure 67 *Alabagrus pecki*: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. wings, F. lateral head and mesosoma, G. propodeum and tergum 1.

27.vi–18.vii.2011, DHJPAR0045021 (EMUS). PARATYPE: 1 ♂, same data as holotype, DHJPAR0045061 (EMUS).

Alabagrus pecki Sharkey, 1988
Figure 67

Alabagrus pecki Sharkey, 1988:395. Holotype ♀, Mexico (CNCI)

DIAGNOSIS. Females: Gena rounded posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Propodeum smooth with areolate pattern barely indicated. Propleuron and mesoscutum yellow. Forewing yellow in basal $\frac{3}{4}$, infusate distally. Fore- and midfemora mostly or entirely melanic. Margins of mesopleuron without a greenish tinge, concolorous with remainder of pleuron. Males: Propodeum with more sculpture, transverse and longitudinal rugae/carinae present. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Propleuron yellow, or melanic and yellow. Forefemur melanic.

DESCRIPTION. Body length 8.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron smooth. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or

depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Propodeum with more sculpture, weak transverse and longitudinal rugae/carinae present; forewing almost entirely infusate; first median tergite may have a rounded median longitudinal carina.

DISTRIBUTION. Southern Mexico (Palenque) to Costa Rica, where it is widespread at lower altitudes.

Alabagrus quickei Sharkey n. sp.
Figure 68

DIAGNOSIS. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Forefemur melanic. Forewing yellow basally, infusate apically with a large infusate patch below parastigma. Scutellum melanic. Metapleuron entirely pale. Apical metasomal terga melanic.

NOTES. Specimens of this species were included under *A. paruyana* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 9.8 mm. Gena rounded or with an obtuse angle posteroventrally. Occiput dorsolaterally rounded. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron smooth. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge.




Figure 68 *Alabagrus quickei* n. sp., holotype: **A.** lateral habitus, **B.** wings, **C.** dorsal habitus.

Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Donald Quicke, hymenopterist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, P.N. Corcovado, 1–100 m, i–iii.1992, LS 270500 508300 (INBio).

Alabagrus ramyamanjunathae Sharkey n. sp.

Figure 69

DIAGNOSIS. Gena right angled or acute posteroventrally. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Forewing yellow in basal $\frac{2}{3}$, infusate apically with infusate patch posterior to parastigma, costal vein melanic. Hind femur entirely melanic, sometimes with a hint of reddish color ventrally but much darker than *A. paulgoldsteini*. Propodeum orange contrasting with melanic metapleuron. Third tergum lacking transverse depression, or depression barely indicated. Apical metasomal terga melanic.

DESCRIPTION. Body length 8.9 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length

of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Phostria temira*.

ETYMOLOGY. Named in honor of Ms. Ramaya Manjunath of the Centre for Biodiversity Genomics (CBG), University of Guelph, Canada, in recognition of her long and patient curation of the ACG Lepidoptera inventory data insertion to the BOLD web site for DNA barcoding for the CBG.

MATERIAL EXAMINED. HOLOTYPE ♀, from *Phostria temira* on *Ipomoea batatas*, 10.93332°N, 85.25331°W, 135 m, 28.viii–13.ix.2009, DHJPAR0036701 (EMUS).

Alabagrus reddypallii Sharkey n. sp.

Figure 70

DIAGNOSIS. Gena rounded posteroventrally. Propleuron melanic. Pronotum melanic anteriorly, pale posteriorly. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Propodeum mostly smooth with some rugae anteromedially and laterally.

NOTES. Specimens of this species were treated as part of *A. pecki* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 6.3 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Reddy Palli, present chair of the Department of Entomology, University of Kentucky, for his leadership and support.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, 16 km SSE La Virgen, 10.27°N, 84.08°W, 1150 m, 21.iii.–09.iv.2001 (INBio). PARATYPE: 1 ♀, Mexico, Yucatan, Uxmal ruins, 5 m, 2.viii.1983 (HIC).

Alabagrus roibasi Sharkey, 1988

Figure 71

Alabagrus roibasi Sharkey, 1988:397–8. Holotype ♀, Costa Rica (EMUS)

DIAGNOSIS. Gena rounded or with an obtuse angle posteroventrally. Metapleuron entirely pale. Forewing with yellowish pigmentation basally that reaches second submarginal cell. Mesoscutum melanic. Hind coxa with a large melanic patch laterally. Metasoma mostly orange, apical two terga melanic.




Figure 69 *Alabagrus ramyamanjunathae*, holotype: **A.** lateral habitus, **B.** ventral hind femur, **C.** anterior head, **D.** dorsal head, **E.** dorsal habitus, **F.** wings, **G.** lateral head and mesosoma, **H.** propodeum and tergum 1.

DESCRIPTION. Body length 7.3 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at

least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface, or heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum with more pronounced sculpture; first median tergite wider.

HOST INFORMATION. *Syllepte aechmisalis*, *Microthyris prolongalis*, *Phaenodropsis cernalis*, *Phostria oajacalis*.

DISTRIBUTION. Sharkey (1988) listed the distribution of this species as widespread in Mexico and ranging to Guanacaste Costa Rica; however the COI data presented here suggests that he, and consequently Leathers and Sharkey (2003), confounded multiple species within the concept of *A. roibasi*. Here, we treat the former concept of *A. roibasi* as three species, including *A. brianharrisi* and *A. markmetzi*. The holotype of *A. roibasi* is from Guanacaste and matches well morphologically with the specimens listed below. It is likely that many, or perhaps all, of the specimens that are found north of Guanacaste belong to other species.

MATERIAL EXAMINED. Reared specimens: 1♀, from *Syllepte aechmisalis* on *Triumfetta lappula*, 10.83764°N, 85.61871°W, 295 m, host collection 10.vii.1993, 93-SRNP-3578. 1♀, 10.85827°N, 85.61089°W, 280 m, host collection 3.x.2000, 00-SRNP-18739. 1♀, from *Microthyris prolongalis* on *Ipomoea alba*, 10.77175°N, 85.434°W, 305 m, 28.ix–16.x.2011, DHJPAR0046736. 1♀, from *Phaenodropsis cernalis* on *Triplaris melaenodendron*, 10.78506°N, 85.6637°W, 10 m, 7–22.vii.1990, 90-SRNP-1247A. 1♀, 10.78938°N, 85.55098°W, 85 m, host collection 26.viii.2000, 00-SRNP-17699. 1♀, from *Phostria oajacalis* on *Merremia umbellata*, 11.02681°N, 85.49547°W, 290 m, 4.viii–5.ix.2009, DHJPAR0037882. 1♂, 4–21.viii.2009, DHJPAR0036690. 1♀, 1♂, 4–28.viii.2009, DHJPAR0036691, DHJPAR0036693. 1♀, on *Turbina corymbosa*, 11.02865°N,




Figure 70 *Alabagrus reddypallii* n. sp., holotype: **A.** lateral habitus, **B.** lateral head, **C.** wings, **D.** dorsal mesosoma, **E.** dorsal metasoma.




Figure 71 *Alabagrus roibasi*, A–G female; H male: A. lateral habitus, B. wings, C. dorsal head, D. dorsal habitus, E. anterior head, F. lateral head and mesosoma, G. propodeum and tergum 1, H. propodeum and tergum 1.

85.48669°W, 280 m, 16–27.xi.2004, DHJPAR0009358. 1♀, 11.03376°N, 85.47715°W, 290 m, 15.vi–2.vii.2009, DHJPAR0036709 (EMUS, HIC).

Alabagrus rudolfmeieri Sharkey n. sp.

Figure 72

DIAGNOSIS. Precoxal sulcus absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Pronotum melanic anteriorly, pale posteriorly. Forewing almost entirely infusate, ignore small clear patches posterad stigma. Hind femur entirely melanic.

DESCRIPTION. Body length 9.3 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Rudolf Meier, dipterist extraordinaire, for his unselfish mentoring of M.J.S.'s graduate student Sarah Meierotto.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, Buenos Aires, Estación Altamira, Senero a Casa Coca, 1700 m, 16.vii–16.viii.2000, L S 574400 331750 (INBio).

Alabagrus sarahmeierottoae Sharkey n. sp.

Figure 73

DIAGNOSIS. Females: Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Propodeum with complete areolation. First tergum with well-defined

median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body. Body orange and melanic. Pronotum melanic. Metasoma entirely orange. Propodeum sculptured with irregular rugae. Metapleuron entirely orange. First tergum weakly convex. Males: Gena rounded or with an obtuse angle posteroventrally. Pronotum melanic.

NOTES. Specimens of this species, including the holotype, were included under *A. englishi* in Leathers and Sharkey (2003).




Figure 72 *Alabagrus rudolfmeieri* n. sp., holotype: A. dorsal mesosoma, B. dorsal metasoma, C. lateral habitus, D. wings.




Figure 73 *Alabagrus sarahmeierottoae* n. sp., A–D holotype; E paratype male: A. lateral habitus, B. wings, C. dorsal habitus, D. lateral head and mesosoma, E. propodeum and tergum 1.

DESCRIPTION. Body length 7.61 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.4× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Propodeum areolate; median tergite 1 with sharp median carina.

ETYMOLOGY. Named in honor of Sarah Meierotto, graduate student extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, P.N. Guanacaste, Sta. Cecilia, 22.x–2.xi.1992, L_N 830200 880200 (INBio). PARATYPES: 1♂, Cartago, P.N. Tapantí, Macizo de la Muerte, Torre circa del Est., 1200 m, 9.vi.2000, L_N 193800 560050 (INBio). 1♀, Limón, Sector Corcori, 30 km N Cariari, 100 m, ii.1995, L_N 286000 567500 (HIC).

Alabagrus sarahsharkeyae Sharkey n. sp.

Figure 74

DIAGNOSIS. Precoxal sulcus almost absent, represented by small, shallow depression posteriorly, crenulae always lacking. Propodeum mostly smooth with median longitudinal and transverse carinae indicated. Metasoma entirely pale. Propleuron melanic. Pronotum pale. Ovipositor longer than body.

NOTES. Specimens of this species were included under *A. englishi* in Leathers and Sharkey (2003).




Figure 74 *Alabagrus sarahsharkeyae* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

DESCRIPTION. Body length 7 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.3× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Sarah Sharkey, youngest sister of M.J.S. and chef extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, P.N. Guanacaste, Sta. Cecilia, 3–9.ii.1992, L_N 330200 380200 (INBio). PARATYPE: 1 ♀, same data as holotype, except 700 m, 1994 (HIC).

Alabagrus sarapiqui Leathers and Sharkey, 2003

Figure 75

Alabagrus sarapiqui Leathers and Sharkey, 2003:44–45. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Gena acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. Body black and yellow. Pronotum melanic. Metasoma entirely yellow.

DESCRIPTION. Body length 6.5 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Male with more pronounced propodeal sculpture.

DISTRIBUTION. Costa Rica.

Alabagrus scottmilleri Sharkey n. sp.

Figure 76

DIAGNOSIS. Females: Head melanic. Propleuron yellow. Propodeum areolate. Midfemur yellow. Hind coxa with a black patch laterally or near apex. Forewing banded from base: yellow, infusate, yellow, infusate, second yellow band complete to hind margin of wing. Midfemur entirely pale. Ovipositor longer than metasoma and shorter than body. Males: Median tergite 1 entirely melanic. Forefemur pale.

DESCRIPTION. Body length 5.8 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing mostly infusate; propodeum and metasomal terga almost entirely melanic; midfemur and hind leg almost entirely black.

HOST INFORMATION. *Neurophyseta* Janzen222, *Neurophyseta* BioLep219, *Neurophyseta camptogrammalis* DHJ01.

ETYMOLOGY. Named in honor of Dr. Scott Miller of the U.S. National Museum of Natural History and Smithsonian Institution, the taxonomist who has long and masterfully taxonomized all of the ACG Dalceridae and supported the movement of ACG Lepidoptera into the Smithsonian safe harbor.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Neurophyseta* Janzen222 on *Lomariopsis vestita*, 10.90093°N, 85.28915°W, 400 m, 17.ii–4.iii.2009, DHJPAR0030469 (EMUS). PARATYPES: reared: 1 ♀, from *Neurophyseta* BioLep219 on *Alsophila firma*, 10.90661°N,




Figure 75 *Alabagrus sarapiqui*, paratype: A. hind leg, B. wings, C. lateral habitus, D. dorsal habitus, E. propodeum.

85.28784°W, 400 m, 24.xii.2009–15.i.2010, DHJPAR0037942. 1 ♂, from *Neurophyseta camptogrammalis* DHJ01, 10.89928°N, 85.27486°W, 420 m, 21.iii–5.iv.2013, DHJPAR0052088 (EMUS, HIC).

Alabagrus scottshawi Sharkey n. sp.

Figure 77

DIAGNOSIS. Gena acute posteroventrally. First tergum weakly convex, almost flat. Forefemur yellow apically and basally, melanic centrally, or mostly yellow but melanic distally. Mesoscutum melanic. Terga 1–3 yellow, remaining terga black.

NOTES. Specimens of this species were included under *A. pachamama* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 9.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron smooth, or with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1.2\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. Scott Shaw, braconologist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Heredia, La Selva Biol. Sta., 50–150 m, 10.43°N, 84.02°W, 7.vii.2000 (INBio). PARATYPES: 7 ♀, Heredia, La Selva Biol. Sta., 50–150 m, ii, iv, vi, vii, ix (INBio,



Figure 76 *Alabagrus scottmilleri* n. sp., A–D, F–H holotype; E paratype male: A. lateral habitus, B. anterior head, C. dorsal head, D. dorsal habitus, E. lateral habitus, F. wings, G. lateral head and mesosoma, H. propodeum and tergum 1.




Figure 77 *Alabagrus scottshawi* n. sp., holotype: A. lateral habitus, B. dorsal habitus, C. lateral head and mesosoma, D. propodeum and tergum 1, E. wings.




Figure 78 *Alabagrus semihesperheidei* n. sp., holotype: **A.** lateral habitus, **B.** dorsal habitus, **C.** wings, **D.** lateral head, **E.** lateral mesosoma, **F.** propodeum and tergum 1.

HIC, EMUS). 2 ♀, PANAMA, Darien, P.N. Darien, Est. Rancho Frio, 80 m, 16.xi.2000–17.i.2001 and 21.iii.–4.iv.2000 (EMUS).

***Alabagrus semihesperheidei* Sharkey n. sp.**

Figure 78

DIAGNOSIS. Gena rounded or with an obtuse angle posteroventrally. Metapleuron entirely pale. Median tergites 1 and 2 partly or entirely red; tergite 3 and following tergites melanic.

DESCRIPTION. Body length 8.3 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

ETYMOLOGY. Named for its similarity to *A. hesperheidei*.

MATERIAL EXAMINED. HOLOTYPE ♀, Puntarenas, Península de Osa, Rancho Quemado, 200 m, vii.1991, L_S 292500 511000 (INBio).

***Alabagrus stigma* (Brullé, 1846)**

Figure 79

Agathis stigma Brullé, 1846:501. Holotype ♀, Brazil (MNHN)

Microdus stigmaterus Cresson, 1865:65. Holotype ♀, Cuba (ANSP)

Microdus diatraeae Turner, 1918:82. Holotype ♀, Guyana (BMNH)

Alabagrus citreistigma Enderlein, 1920:203. Holotype ♀, Brazil (ZMPA)

Microdus crossi Brèthes, 1927:163. Holotype ♀, Argentina (depository?)

Microdus sacchari Myers, 1931:274. Holotype ♀, Guyana (BMNH)

Alabagrus stigma (Sharkey, 1988:401–402)

DIAGNOSIS. Hind femur heavily sculptured ventrally with aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. Ovipositor longer than metasoma but not distinctly longer than body, or distinctly longer than body. First tegite with a median carinae and lateral rugosities in female; male with two longitudinal carinae and more lateral rugae. Forewing pattern variable but stigma and area near stigma always yellow.

DESCRIPTION. Body length 9.1 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron entirely densely punctate or aciculate. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth, or punctate in anterior half, areolate posteriorly, with or without areolae. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 1.5× longer than wide; with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body, or distinctly longer than body.

Males. Much more pronounced propodeal sculpture; median tergite 1 with two longitudinal carinae and entire surface rugosostriate.

HOST INFORMATION. *Diatraea canella*, *Diatraea*, *Diatraea lineolata*, *Diatraea saccharalis*.

DISTRIBUTION. Widespread from the southeastern USA (e.g., Florida, Louisiana) through most of the lowland, warm Neotropics.

***Alabagrus stiremani* Sharkey n. sp.**

Figure 80

DIAGNOSIS. Precoxal sulcus absent. Propodeum smooth, melanic. Metasomal terga 1–4 yellow, remaining terga black.




Figure 79 *Alabagrus stigma*, A–F female; H male: A. lateral habitus, B. dorsal habitus, C. anterior head, D. lateral head and mesosoma, E. dorsal head, F. propodeum and tergum 1, G. wings, H. propodeum and tergum 1.

NOTES. Specimens of this species were included under *A. paruyana* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 10.5 mm. Gena rounded or with an obtuse angle posteroventrally. Occiput dorsolaterally rounded. Precoxal sulcus absent or almost absent, represented at most by small, shallow depression posteriorly, crenulae always lacking. Margin between metepisternum and metepimeron with very small weak crenulae. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum $1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Dr. John Stireman, dipterist extraordinaire.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, Estación Pitilla, 9 km S Santa Cecilia, 700 m, iii.1995, L_N 329500 380450 (INBio). PARATYPE: 1 ♀, Guanacaste, Estación Pitilla, 9 km S Santa Cecilia, 700 m, vi.1994, L_N 330200 380200 (HIC).

Alabagrus tanyadapkeyae Sharkey n. sp.

Figure 81

DIAGNOSIS. Terga melanic except rarely first tergum with some red color. Precoxal sulcus with large foveae extending about $\frac{1}{2}$ length of mesopleuron. Forewing infusate in basal $\frac{2}{3}$, white/opaque apically. Mesosoma including legs entirely black.

DESCRIPTION. Body length 7.5 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate $\frac{1}{2}$

or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins, or areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum $1.1\times$ longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Forewing entirely infusate; propodeum and median tergite 1 red.

HOST INFORMATION. *Phostria cyrisalis*.

ETYMOLOGY. Named in honor of Ms. Tanya Dapkey of the University of Pennsylvania, Philadelphia, the information and specimen manager who has long maintained the curatorial organization and databasing process for the ACG Lepidoptera inventory.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Phostria cyrisalis* on *Goethalsia meiantha*, 11.01602°N, 85.38053°W, 380 m, 17.vii–21.viii.2010, DHJPAR0040218 (EMUS). PARATYPES: reared: 1 ♀, from *Phostria cyrisalis* on *Goethalsia meiantha*, 10.93010°N, 85.252050°W, 109 m, 25.vi–15.vii.2014, DHJPAR0055970. 1 ♀, 17.vii–23.viii.2010, DHJPAR0040225. 1 ♂, on *Trichospermum galeottii*, 10.95644°N, 85.266110°W, 121 m, 30.ix–23.x.2014, DHJPAR0057429 (EMUS, HIC). Nonreared: 1 ♀, Heredia, 50–150 m, Est. Biol. La Selva, 10.43°N, 84.02°W, v.2000 (INBio).

Alabagrus tommyersi Sharkey n. sp.

Figure 82

DIAGNOSIS. Third tergum with deep transverse depression. First tergum weakly convex. Forewing with two yellow bands, one complete




Figure 80 *Alabagrus stiremani* n. sp., holotype: **A.** lateral habitus, **B.** wings, **C.** dorsal habitus.

basal yellow band and a weaker, incomplete, yellow, triangular band directly below (posterior to) stigma. Precoxal sulcus with one or several distinct foveae posteroventrally.

DESCRIPTION. Body length 11.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola, or lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 0.8× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum with deep transverse depression. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum areolate.

HOST INFORMATION. *Syllepte belialis*.

ETYMOLOGY. In honor of Tom Myers for the generous use of his diverse property for research by the Sharkey lab.

MATERIAL EXAMINED. HOLOTYPE ♀, Guanacaste, 3 km N de Nacaome, 100 m, P.N. Barra Honda, 5–27.vii.1992, L_N 239000 386000 (HIC). PARATYPES: reared: 1 ♀, from *Syllepte belialis* on

Casearia corymbosa, 10.84389°N, 85.61384°W, 300 m, 26.ix–26.x.2004, DHJPAR0015402. 1 ♀, 4–25.vi.1982, 82-SRNP-162. 1 ♂, 10.85827°N, 85.61089°W, 280 m, 25.vii–27.viii.1992, DHJPAR0015401. Nonreared paratypes: 3 ♀, Guanacaste, vi–viii (EMUS, HIC, INBio).

Alabagrus varius (Enderlein, 1920)

Figure 83

Astiria varia Enderlein, 1920:207. Holotype ♀, Mexico (ZMPA)

Alabagrus varius (Sharkey, 1988:410)

DIAGNOSIS. Occiput with squared protuberance dorsolaterally. Mesosoma mostly pale anteriorly and posteriorly, melanic at midlength.

DESCRIPTION. Body length 10 mm. Gena right angled or acute posteroventrally, or rounded or with an obtuse angle posteroventrally. Clypeus with lateral protrusions. Occiput dorsolaterally with squared protuberances. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

DISTRIBUTION. Known from southern Mexico and Guanacaste (Leathers and Sharkey, 2003).

Alabagrus victoriapookae Sharkey n. sp.

Figure 84

DIAGNOSIS. Occiput with squared protuberances dorsolaterally. Mesosoma melanic except propodeum pale.

DESCRIPTION. Body length 9.5 mm. Gena rounded or with an obtuse angle posteroventrally. Clypeus with lateral protrusions. Occiput dorsolaterally with squared protuberances. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly, or distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron smooth. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum lacking complete areolae, usually mostly or completely smooth. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor distinctly longer than body.

Males. Unknown.

ETYMOLOGY. Named in honor of Victoria Pook, scientist extraordinaire and former student of M.J.S.

MATERIAL EXAMINED. HOLOTYPE ♀, Alajuela, A.C.A. San Carlos, P.N. Volcán Arenal, Sector Colada, 600 m, 25.vi–26.vii.1999, L_N 269900 456750 (INBio).

Alabagrus voto Sharkey, 1988

Figure 85

Alabagrus voto Sharkey, 1988:411. Holotype ♀, Costa Rica (EMUS)

DIAGNOSIS. Head yellow. Forewing with two distinct yellow bands. Fore- and midlegs entirely yellow.




Figure 81 *Alabagrus tanyadapkeyae* n. sp., A–G holotype; H paratype female: A. lateral habitus, B. dorsal habitus, C. wings, D. anterior head, E. lateral mesosoma, F. lateral head, G. propodeum and tergum 1, H. tergum 1.




Figure 82 *Alabagrus tommyersi* n. sp., A–E, G, H holotype; F paratype male: A. lateral habitus, B. dorsal head, C. lateral head, D. anterior head, E. dorsal habitus, F. propodeum and tergum 1, G. wings, H. propodeum and tergum 1.




Figure 83 *Alabagrus varius*: **A.** lateral habitus, **B.** ovipositor (on same scale as A), **C.** anterior head, **D.** lateral head, **E.** lateral mesosoma, **F.** wings, **G.** propodeum and tergum 1, **H.** dorsal habitus.

DESCRIPTION. Body length 7.1 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth;

sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression,




Figure 84 *Alabagrus victoriapookae* n. sp., holotype: **A.** lateral habitus, **B.** ovipositor (on same scale as A), **C.** anterior head, **D.** wings, **E.** lateral head, **F.** lateral mesosoma, **G.** dorsal head, **H.** propodeum and tergum 1, **I.** dorsal habitus.




Figure 85 *Alabagrus voto*, A–D, G–I female; E, F male: A. lateral habitus, B. dorsal habitus, C. anterior head, D. dorsal head, E. propodeum, F. tergum 1, G. wings, H. lateral head and mesosoma, I. propodeum and tergum 1.




Figure 86 *Alabagrus watsoni*, A, C–G female; B male: A. lateral habitus, B. propodeum and terga 1–3, C. dorsal head, D. dorsal habitus, E. anterior head, F. wings, G. lateral head and mesosoma.

or depression barely indicated. Ovipositor short, not longer than metasoma, or longer than metasoma but not longer than body.

Males. Propodeum with more sculpture; first median tergite with distinct median carina; body color may be slightly darker; first median tergite wider.

DISTRIBUTION. Costa Rica: Guanacaste and Puntarenas (Leathers and Sharkey, 2003).

Alabagrus watsoni Leathers and Sharkey, 2003

Figure 86

Alabagrus watsoni Leathers and Sharkey, 2003:51–51. Holotype ♀, Costa Rica (INBio)

DIAGNOSIS. Gena acute posterolaterally. Precoxal sulcus with one or two large foveae posteroventrally. Terga 1–2 orange, tergum 3 orange and melanic, remaining terga melanic. Median tergite 3 pale anteriorly, melanic posteriorly. Forewing almost entirely infusate, or mostly infusate, but mostly clear basad 1cu-a. Metapleuron mostly orange.

DESCRIPTION. Body length 6.8 mm. Gena right angled or acute posteroventrally. Precoxal sulcus with one or several distinct foveae posteroventrally, with or without a smooth groove extending anteriorly. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron areolate rugose in ventral quarter or less. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 0.9× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Propodeum with slightly more sculpture; first median tergite with sharper median carina.

HOST INFORMATION. *Omiodes fulvicauda*, *Omiodes humeralis*, *Omiodes insolitalis*, *Omiodes humeralis*DHJ02, *Omiodes humeralis*DHJ07, *Lamprosema* Dapkey23.

DISTRIBUTION. Costa Rica.

MATERIAL EXAMINED. 1♀, from *Lamprosema* Dapkey23 on *Serjania rhombea*, 10.87766°N, 85.393430°W, 645 m, 29.iii–14.iv.2014, DHJPAR0055347. 1♀, from *Omiodes fulvicauda* on *Dioclea malacocarpa*, 10.90661°N, 85.28784°W, 400 m, 14–31.v.2010, DHJPAR0039356. 1♂, from *Omiodes humeralis* on *Inga chocoensis*, 10.89678°N, 85.27001°W, 420 m, 13.xi–9.xii.2013, DHJPAR0054496. 1♂, 10.9163°N, 85.37869°W, 460 m, 22.ix–13.x.2005, DHJPAR0015519. 2♀, 11.01926°N, 85.40997°W, 440 m, host collection 9.xi.2007, 07-SRNP-33841, 07-SRNP-33842. 1♀, on *Inga oerstediana*, 10.90037°N, 85.37254°W, 500 m, 4–28.x.2007, DHJPAR0023531. 1♂, 10.90661°N, 85.28784°W, 400 m, 13–28.xii.2004, DHJPAR0015526. 1♀, 29.ii–14.iii.2012, DHJPAR0048722. 1♀, 10.91847°N, 85.30338°W, 320 m, 23.ii–16.iii.2004, DHJPAR0015341. 1♀, 23.x–4.xi.2009, DHJPAR0037915. 1♂, 10.93332°N, 85.25331°W, 135 m, 3–25.i.2010, DHJPAR0037937. 1♀, 10.95642°N, 85.26617°W, 123 m, 8–25.ix.2010, DHJPAR0041201. 1♀, 10.9867°N, 85.38503°W, 440 m, 23.xi–10.xii.2011, DHJPAR0046747. 1♀, 23.xi–12.xii.2011, DHJPAR0046740. 1♀, 23.xi–15.xii.2011, DHJPAR0046743. 1♀, 11.0006°N, 85.438°W, 620 m, 11–28.i.2008, 08-SRNP-20276. 1♂, 11.01373°N, 85.42531°W, 525 m, 14.xi–2.xii.2007, DHJPAR0023290. 2♀, 14.xi–3.xii.2007, DHJPAR0023299, DHJPAR0023300. 1♂, 14.xi–7.xii.2007, DHJPAR0023298. 1♀, 18.xii.2007–8.i.2008, DHJPAR0023286. 1♀, 7.xi–2.xii.2007, DHJPAR0023297. 1♀, 7–26.xi.2007, DHJPAR0023285. 1♂, 11.01618°N, 85.35902°W, 340 m, 10–30.x.2007, DHJPAR0022189. 1♂, 6.xii.2007–10.i.2008, DHJPAR0023293. 1♀, 6–24.xii.2007, 07-




Figure 87 *Alabagrus yuanmaofangi* n. sp., holotype: A. lateral habitus, B. wings, C. dorsal habitus.

SRNP-66142. 1♀, 11.01825°N, 85.37199°W, 360 m, 25.ix–8.x.2010, DHJPAR0041202. 1♀, 11.01926°N, 85.40997°W, 440 m, 1.xi–15.xii.2007, DHJPAR0028282. 1♀, 8–29.xii.2007, DHJPAR0023292. 1♂, 9.xi–2.xii.2007, DHJPAR0028283. 1♀, 9.xi–7.xii.2007, DHJPAR0028285. 1♀, host collection 1.xi.2007, 07-SRNP-33778. 1♀, host collection 26.xii.2007, 07-SRNP-34425. 3♀, host collection 7.xii.2007, 07-SRNP-34085, 07-SRNP-34088, 07-SRNP-34089. 1♀, host collection 9.xi.2007, 07-SRNP-33856. 1♂, 11.03306°N, 85.42876°W, 400 m, 15.i–3.ii.2008, DHJPAR0028293. 1♀, 17.i–10.ii.2008, DHJPAR0028308. 1♀, 21.i–11.ii.2008, DHJPAR0023535. 1♀, 27.xi–11.xii.2007, DHJPAR0028286. 2♀, 27.xi–14.xii.2007, DHJPAR0023287, DHJPAR0023289. 1♂, 27.xi–20.xii.2007, DHJPAR0028291. 1♀, 27.xi–22.xii.2007, DHJPAR0028290. 1♀, 7–25.xii.2007, DHJPAR0023288. 1♀, 11.04249°N, 85.40339°W, 390 m, 22.xi–20.xii.2007, DHJPAR0028284. 1♀, on *Inga samanensis*, 10.89666°N, 85.29003°W, 400 m, 17.xii.2014–7.i.2015, DHJPAR0057000. 1♀, 10.93332°N, 85.25331°W, 135 m, host collection 2.ii.2010, 10-SRNP-75204. 1♂, 11.01373°N, 85.42531°W, 525 m, 12–25.ii.2007, DHJPAR0021196. 1♀, 11.01926°N, 85.40997°W, 440 m, host collection 14.i.2004, 04-SRNP-30212. 1♀, on *Inga vera*,




Figure 88 *Alabagrus yuchinkengae* n. sp., holotype male: A. lateral habitus, B. dorsal habitus, C. wings, D. lateral head and mesosoma, E. propodeum and tergum 1.

10.94041°N, 85.68043°W, 275 m, 23.vi–7.vii.1993, DHJPAR0015390. 1♀, from *Omiodes humeralis*DHJ02 on *Inga oerstediana*, 10.93332°N, 85.25331°W, 135 m, 11–22.i.2010, DHJPAR0037938. 1♀, 11–25.i.2010, DHJPAR0037936. 1♀, from *Omiodes humeralis*DHJ07 on *Inga oerstediana*, 10.93319°N, 85.25335°W, 95 m, 8.i–8.ii.2008, DHJPAR0023530. 1♀, from *Omiodes insolutalis* on *Mucuna holtonii*, 10.90376°N, 85.30274°W, 340 m, 10–26.iii.2011, DHJPAR0042827. 1♀, 5.ii.2011, 11-SRNP-40659. 1♀, 10.90528°N, 85.27882°W, 405 m, 2–16.vi.2011, DHJPAR0045018. 1♀, 10.91847°N, 85.30338°W, 320 m, 24.iii–20.iv.2011, DHJPAR0042825 (EMUS, HIC). Also known from La Selva Biol. Station, Heredia (Leathers and Sharkey, 2003).

***Alabagrus yuanmaofangi* Sharkey n. sp.**

Figure 87

DIAGNOSIS. Gena right angled or acute posteroventrally. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum with a rounded longitudinal bulge. Ovipositor longer than metasoma but shorter than body. Mesoscutum pale. Fore- and midfemora mostly or entirely melanic. Hind coxa entirely pale.

NOTES. Specimens of this species were included under *A. cocto* and *A. yaruro* in Leathers and Sharkey (2003).

DESCRIPTION. Body length 7.4 mm. Gena right angled or acute posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur with weak sparse punctures ventrally, little or no more heavily sculptured than dorsal surface. First tergum 1.2× longer than wide; varying from weakly convex to with a rounded longitudinal bulge. Third tergum lacking transverse depression,

or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Similar to females.

HOST INFORMATION. *Desmia* Janzen09, *Desmia* biolep02, *Desmia* biolep06, *Desmia* ploralisDHJ03, *Trichaea pilicornis*.

ETYMOLOGY. Named in honor of Mr. Yuan-Mao Fang of Jiangsu, China (24 Aug. 1912–2 Dec. 1986), father to Ming Fang of Hawaii, who is a major supporter of the ACG forest in which this wasp lives.

MATERIAL EXAMINED. HOLOTYPE ♀, host=*Desmia* Janzen09 on *Psychotria panamensis*, 10.94076°N, 85.3177°W, 461 m, 22.ix–11.x.2009, DHJPAR0037899 (EMUS). PARATYPES: 1♂, from *Desmia* biolep02 on *Hoffmannia valerii*, 10.89928°N, 85.27486°W, 420 m, 9–28.vii.2009, DHJPAR0036334. 1♀, from *Desmia* BioLep06, 14.i–8.ii.2010, DHJPAR0037941. 1♂, from *Desmia* Janzen09 on *Psychotria officinalis*, 10.94741°N, 85.31501°W, 491 m, 3–21.xi.2009, DHJPAR0037894. 1♀, from *Desmia* ploralisDHJ03 on *Psychotria chagrensis*, 4–24.x.2013, DHJPAR0053600. 1♀, from *Trichaea pilicornis* on *Psychotria chagrensis*, 10.8962°N, 85.27769°W, 430 m, 23.iii–12.iv.2011, DHJPAR0042834 (EMUS, HIC).

***Alabagrus yuchinkengae* Sharkey n. sp.**

Figure 88

DIAGNOSIS. Gena rounded or with an obtuse angle posteroventrally. Median tergites 1–2 red, tergite 3 pale anteriorly, melanic posteriorly, remaining tergites melanic.

DESCRIPTION. Body length 12.1 mm. Gena rounded or with an obtuse angle posteroventrally. Precoxal sulcus distinct and foveolate ½ or more length of mesopleuron. Margin between metepisternum and metepimeron distinctly and deeply crenulate. Metapleuron mostly or entirely smooth; sculpture, if present, restricted to extreme margins. Propodeum areolate, with at least one closed areola. Hind femur heavily sculptured ventrally with large aciculations or rugosities, distinctly more heavily sculptured than dorsal surface. First tergum 0.9× longer than

wide; varying from weakly convex to with a rounded longitudinal bulge, or with well-defined median longitudinal carina. Third tergum lacking transverse depression, or depression barely indicated. Ovipositor longer than metasoma but not longer than body.

Males. Unknown.

HOST INFORMATION. *Microthyris prolongalis*.

ETYMOLOGY. Named in honor of Ms. Yuchin Keng of Kengchuan, Henan, China (16 Feb. 1911–12 Jul. 1987), mother to Ming Fang of Hawaii, who is a major supporter of the ACG forest in which this wasp lives.

MATERIAL EXAMINED. HOLOTYPE ♂, from *Microthyris prolongalis* on *Ipomoea trifida*, 10.77074°N, 85.42874°W, 365 m, 24.vi–18.vii.2011, DHJPAR0045112 (EMUS).

ACKNOWLEDGMENTS

Thank you to Drs. Kees van Actenberg and Yves Braet for reviewing the manuscript. We gratefully acknowledge the unflagging support of the team of ACG parataxonomists (D.J. and W.H.), who found and reared the specimens used in this study, and the team of biodiversity managers who protect and manage the ACG forests that host these wasps and their caterpillar hosts. The study has been supported by U.S. National Science Foundation grants BSR 9024770 and DEB 9306296, 9400829, 9705072, 0072730, 0515699 and grants from the Wege Foundation, International Conservation Fund of Canada, Jessie B. Cox Charitable Trust, Blue Moon Fund, Guanacaste Dry Forest Conservation Fund, Área de Conservación Guanacaste, Permian Global, and University of Pennsylvania (D.J. and W.H.). This study has been supported by the Government of Canada through its ongoing support of Genome Canada, the Biodiversity Institute of Ontario, and the Ontario Genomics Institute (2008-0GI-ICI-03; M.A.S.) and by a Discovery Grant from Natural Sciences and Engineering Research Council of Canada (M.A.S.). Funding was also provided by Hatch projects KY008041 and KY008065 (to M.J.S.). The information reported in this paper (17-08-113) is part of a project of the Kentucky Agricultural Experiment Station and is published with the approval of the Director.

LITERATURE CITED

- Brèthes, J. 1927. Parasites and hyperparasites of *Diatraea saccharalis* in Tucumán sugar-cane. *Bulletin of Entomological Research* 18:205–207.
- Brullé, M.A. 1846. Tome Quatrième. Des Hyménoptères. Les Ichneumonides. pp. 56–521 In: Lepeletier de Saint-Fargeau A. “Histoire Naturelles des Insectes.” Paris. 680 pp.
- Cameron, P. 1887. Hymenoptera. Pp. 471–472 In: Godman F.D. & Salvin O. *Biologia Centrali-Americana*; or, Contributions to the knowledge of the fauna and flora of Mexico and Central America. Zoology. 471–472.
- Cameron, P. 1905. New Hymenoptera, mostly from Nicaragua. *Invertebrata Pacifica* 1:46–69.
- Cresson, E.T. 1865. On the Hymenoptera of Cuba. *Proceedings of the Entomological Society of Philadelphia* 4:1–200.
- Drummond, A.J., B. Ashton, M. Cheung, J. Heled, M. Kearse, R. Moir, S. Stones-Havas, T. Thierer, and A. Wilson. 2009. Geneious version 6.1.5. <http://www.geneious.com> [accessed 1 April 2015].
- Enderlein, G. 1920. Zur Kenntnis aussereuropäischer Braconiden. *Archiv für Naturgeschichte* 84(A)11(1918):51–224.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39:783–791.
- Gries, C., E.E. Gilbert, and N.M. Franz. 2014. Symbiota—A virtual platform for creating voucher-based biodiversity information communities. *Biodiversity Information Journal* 2014:e1114.
- Hebert, P.D.N., E.H. Penton, J.M. Burns, D.H. Janzen, and W. Hallwachs. 2004. Ten species in one: DNA barcoding reveals cryptic species in the Neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America* 101:14812–14817.
- Huelsenbeck, J.P., and B. Rannala. 2004. Frequentist properties of Bayesian posterior probabilities of phylogenetic trees under simple and complex substitution models. *Systematic Biology* 53:904–913.
- Huelsenbeck, J.P., and F. Ronquist. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755.
- Janzen, D.H., W. Hallwachs, P. Blandin, J.M. Burns, J. Cadiou, I. Chacon, T. Dapkey, A.R. Deans, M.E. Epstein, B. Espinoza, J.G. Franclemont, W.A. Haber, M. Hajibabaei, J.P.W. Hall, P.D.N. Hebert, I.D. Gauld, D.J. Harvey, A. Hausmann, I. Kitching, D. Lafontaine, J. Landry, C. Lemaire, J.Y. Miller, J.S. Miller, L. Miller, S.E. Miller, J. Montero, E. Munroe, S. Rab Green, S. Ratnasingham, J.E. Rawlins, R.K. Robbins, J.J. Rodriguez, R. Rougerie, M.J. Sharkey, M.A. Smith, M.A. Solis, J.B. Sullivan, P. Thiaucourt, D.B. Wahl, S.J. Weller, J.B. Whitfield, K.R. Willmott, D.M. Wood, N.E. Woodley, and J.J. Wilson. 2009. Integration of DNA barcoding into an ongoing inventory of complex tropical biodiversity. *Molecular Ecology Resources* 9(Supplement 1):1–26.
- Katoh, K., K. Kuma, H. Toh, and T. Miyata. 2006. MAFFT version 5: Improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33:511–518.
- Leathers, J., and M.J. Sharkey. 2003. Taxonomy and life history of Costa Rican *Alabagrus* (Hymenoptera: Braconidae), with a key to world species. *Contributions in Science* 497:1–82.
- Maddison, W.P., and D.R. Maddison. 2005. MacClade: Analysis of phylogeny and character evolution. Version 4.08a. <http://www.macclade.org/> [accessed 1 April 2015].
- Marshall, D.C., C. Simon, and T.R. Buckley. 2006. Accurate branch length estimation in partitioned Bayesian analyses requires accommodation of among-partition rate variation and attention to branch length priors. *Systematic Biology* 55:992–1003.
- Myers, J.G. 1931. Descriptions and records of parasitic Hymenoptera from British Guiana and the West Indies. *Bulletin of Entomological Research* 22(2):267–277.
- Ratnasingham, S., and P.D.N. Hebert. 2007. The Barcode of Life Data System. *Molecular Ecology Notes* 7(3):355–364.
- Rodriguez, F., J.L. Oliver, A. Marin, and J.R. Medina. 1990. The general stochastic model of nucleotide substitution. *Journal of Theoretical Biology* 142:485–501.
- Ronquist, F., and J.P. Huelsenbeck. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
- Sharkey, M.J. 1988. A taxonomic revision of *Alabagrus* (Hymenoptera: Braconidae). *Bulletin of the British Museum (Natural History)* 57:1–137.
- Sharkey, M.J., and E.G. Chapman. 2017. Phylogeny of the Agathidinae (Hymenoptera: Braconidae) with a revised tribal classification and the description of a new genus. *Proceedings of the Entomological Society of Washington* 119(sp1):823–842.
- Sharkey, M.J., and R.A. Wharton. 1997. Morphology and terminology. In *Manual of the New World genera of Braconidae (Hymenoptera)*, eds. R.A. Wharton, P.M. Marsh, and M.J. Sharkey, 19–38. Washington, D.C.: International Society of Hymenopterists 1.
- Swofford, D.L. 2003. PAUP* Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sunderland, Massachusetts: Sinauer Associates.
- Turner, R.E. 1918. On Braconidae parasitic on *Diatraea saccharalis* in Demerara. *Bulletin of Entomological Research* 9:81–82.
- Yoder, M.J., I. Mikó, K.C. Seltmann, M.A. Bertone, and A.R. Deans. 2010. A gross anatomy ontology for Hymenoptera. *PLoS ONE* 5:e15991.
- Zwickl, D.J. 2006. *Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion*. PhD dissertation. Austin: The University of Texas. <https://repositories.lib.utexas.edu/handle/2152/2666> [accessed May 17, 2016], 115 pp.

Submitted for publication August 23, 2017; accepted February 20, 2018.

Appendix 1 Specimens in the molecular analysis including specimen number, host, geographic area, and BOLD and GenBank accession numbers. Specimens with unconventional host species names (e.g., *Hyalorista exuvialis*DHJ02) are as of yet undescribed.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Pharpa</i> sp. (outgroup)	H1158			Colombia: Meta	ATRMK257-11
<i>Alabagrus almasolisae</i>	DHJPAP0028269	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF031-09
<i>Alabagrus almasolisae</i>	DHJPAP0028270	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF032-09
<i>Alabagrus almasolisae</i>	DHJPAP0028271	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF033-09
<i>Alabagrus almasolisae</i>	DHJPAP0028272	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF034-09
<i>Alabagrus almasolisae</i>	DHJPAP0028273	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF035-09
<i>Alabagrus almasolisae</i>	DHJPAP0030384	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYB1123-09
<i>Alabagrus almasolisae</i>	DHJPAP0037924	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYC4669-10
<i>Alabagrus almasolisae</i>	DHJPAP0037931	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYC4676-10
<i>Alabagrus almasolisae</i>	DHJPAP0037932	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYC4677-10
<i>Alabagrus almasolisae</i>	DHJPAP0037933	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYC4678-10
<i>Alabagrus almasolisae</i>	DHJPAP0037935	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYC4680-10
<i>Alabagrus almasolisae</i>	DHJPAP0041205	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF1111-11
<i>Alabagrus almasolisae</i>	DHJPAP0041578	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYF1484-11
<i>Alabagrus almasolisae</i>	DHJPAP0042816	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYH574-11
<i>Alabagrus almasolisae</i>	DHJPAP0042817	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYH575-11
<i>Alabagrus almasolisae</i>	DHJPAP0042818	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYH576-11
<i>Alabagrus almasolisae</i>	DHJPAP0053625	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYM2979-13
<i>Alabagrus almasolisae</i>	DHJPAP0054493	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYD3658-14
<i>Alabagrus almasolisae</i>	DHJPAP0054507	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYD3672-14
<i>Alabagrus almasolisae</i>	DHJPAP0054508	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYD3673-14
<i>Alabagrus almasolisae</i>	DHJPAP0055101	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	ASHYH1648-14
<i>Alabagrus almasolisae</i>	DHJPAP0058543	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	MHMYN8143-16
<i>Alabagrus almasolisae</i>	DHJPAP0058544	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	MHMYN8144-16
<i>Alabagrus almasolisae</i>	DHJPAP0058545	Crambidae	<i>Ceratocilia sixolalis</i>	Costa Rica: Alajuela	MHMYN8145-16
<i>Alabagrus andresfreitasi</i>	DHJPAP0036714	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ASHYE1625-09
<i>Alabagrus andresfreitasi</i>	DHJPAP0037880	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ASHYC4625-10
<i>Alabagrus andresfreitasi</i>	DHJPAP0037959	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ASHYC4704-10
<i>Alabagrus andresfreitasi</i>	DHJPAP0050365	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ACGAZ1679-12
<i>Alabagrus andywarreni</i>	DHJPAP0015471	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ASAG157-07
<i>Alabagrus andywarreni</i>	DHJPAP0015472	Crambidae	<i>Phostria truncatalis</i>	Costa Rica: Guanacaste	ASAG158-07
<i>Alabagrus andywarreni</i>	DHJPAP0037906	Crambidae	<i>Phostria metalobalis</i>	Costa Rica: Guanacaste	ASHYC4651-10
<i>Alabagrus andywarreni</i>	DHJPAP0041199	Crambidae	<i>Phostria metalobalis</i>	Costa Rica: Guanacaste	ASHYF1105-11
<i>Alabagrus andywarreni</i>	DHJPAP0044988	Crambidae	<i>Phostria metalobalis</i>	Costa Rica: Guanacaste	ACGAZ209-11
<i>Alabagrus arawak</i>	H1196			Costa Rica: Cartago	MF361679
<i>Alabagrus arawak</i>	H1201			Costa Rica: Heredia	MF361683
<i>Alabagrus arawak</i>	H12092			Costa Rica: Heredia	MF361699
<i>Alabagrus arawak</i>	H12186			Costa Rica: Heredia	MF361703
<i>Alabagrus arua</i>	DHJPAP0015518	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASAG204-07
<i>Alabagrus arua</i>	DHJPAP0021202	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASBC1014-07
<i>Alabagrus arua</i>	DHJPAP0030487	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYB1230-09
<i>Alabagrus arua</i>	DHJPAP0035224	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYE1046-09
<i>Alabagrus arua</i>	DHJPAP0035296	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYE1118-09
<i>Alabagrus arua</i>	DHJPAP0035297	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYE1119-09
<i>Alabagrus arua</i>	DHJPAP0041194	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYF1100-11
<i>Alabagrus arua</i>	DHJPAP0049054	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ACGBA2596-12
<i>Alabagrus arua</i>	DHJPAP0051379	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ACGBA3971-13
<i>Alabagrus axelhausmanni</i>	DHJPAP0042832, H6745	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ASHYH590-11
<i>Alabagrus bernardoespinozai</i>	DHJPAP0036338	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYD1529-09
<i>Alabagrus bobpoolei</i>	DHJPAP0036694	Crambidae	<i>Eulepte</i> Janzen06	Costa Rica: Guanacaste	ASHYE1605-09
<i>Alabagrus bobpoolei</i>	DHJPAP0037887	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Guanacaste	ASHYC4632-10
<i>Alabagrus bobpoolei</i>	DHJPAP0038924	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Guanacaste	ASHYD2497-10
<i>Alabagrus bobpoolei</i>	DHJPAP0041169	Crambidae	<i>Eulepte</i> Janzen12	Costa Rica: Guanacaste	ASHYF1084-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042805	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYH563-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042811	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Guanacaste	ASHYH569-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042813	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYH571-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042814	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYH572-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042815	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYH573-11
<i>Alabagrus bobpoolei</i>	DHJPAP0042828	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYH586-11
<i>Alabagrus bobpoolei</i>	DHJPAP0045785	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ACGAZ974-11
<i>Alabagrus bobpoolei</i>	DHJPAP0048725	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Guanacaste	ACGBA2267-12
<i>Alabagrus bobpoolei</i>	DHJPAP0048727	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Guanacaste	ACGBA2269-12
<i>Alabagrus bobpoolei</i>	DHJPAP0050364	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ACGAZ1678-12
<i>Alabagrus bobpoolei</i>	DHJPAP0051380	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ACGBA3972-13
<i>Alabagrus bobpoolei</i>	DHJPAP0051382	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ACGBA3974-13
<i>Alabagrus bobpoolei</i>	DHJPAP0054495	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYD3660-14
<i>Alabagrus bobpoolei</i>	DHJPAP0054514	Crambidae	<i>Eulepte</i> Solis15	Costa Rica: Alajuela	ASHYD3679-14

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus bobrobbinsi</i>	DHJPAR0037881	Crambidae	<i>Microthyris anormalis</i> DHJ02	Costa Rica: Guanacaste	ASHYC4626-10
<i>Alabagrus bobrobbinsi</i>	DHJPAR0045007	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ228-11
<i>Alabagrus bobrobbinsi</i>	DHJPAR0045008	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ229-11
<i>Alabagrus bobrobbinsi</i>	DHJPAR0045017	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ238-11
<i>Alabagrus bobrobbinsi</i>	DHJPAR0046733	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGBA906-12
<i>Alabagrus bobrobbinsi</i>	DHJPAR0049657	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Alajuela	ASHYB2451-12
<i>Alabagrus brianharrisi</i>	DHJPAR0041187	Crambidae	<i>Phaedropsis cernalis</i>	Costa Rica: Guanacaste	ASHYF1093-11
<i>Alabagrus brianharrisi</i>	DHJPAR0046738	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Guanacaste	ACGBA911-12
<i>Alabagrus cara</i>	BIOUG24991-C07			Costa Rica: Guanacaste	GMACR173-15
<i>Alabagrus cara</i>	BIOUG25233-D08			Costa Rica: Guanacaste	GMACW602-15
<i>Alabagrus cara</i>	DHJPAR0035232	Crambidae	<i>Desmia ufeus</i>	Costa Rica: Alajuela	ASHYE1054-09
<i>Alabagrus cara</i>	DHJPAR0041579, H7675	Crambidae	<i>Diacme</i> BioLep02	Costa Rica: Alajuela	ASHYF1485-11
<i>Alabagrus cara</i>	DHJPAR0052681	Crambidae	<i>Desmia benealis</i> DHJ03	Costa Rica: Alajuela	ASHYM2035-13
<i>Alabagrus cara</i>	DHJPAR0058550	Crambidae	<i>Desmia</i> BioLep22DHJ02	Costa Rica: Alajuela	MHMYN8150-16
<i>Alabagrus cara</i>	H1202			Costa Rica: Heredia	MF361684
<i>Alabagrus cara</i>	H12109			Costa Rica: Heredia	MF361701
<i>Alabagrus cocto</i>	DHJPAR0016927	Crambidae	<i>Conchylodes grammaphora</i>	Costa Rica: Alajuela	ASBR900-07
<i>Alabagrus cocto</i>	DHJPAR0021151	Crambidae	<i>Conchylodes arcifera</i>	Costa Rica: Guanacaste	ASBC963-07
<i>Alabagrus cocto</i>	DHJPAR0030383	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYB1122-09
<i>Alabagrus cocto</i>	DHJPAR0040345, H8044	Crambidae	<i>Syllepte</i> Solis22	Costa Rica: Guanacaste	ASHYE2481-11
<i>Alabagrus cocto</i>	DHJPAR0041594	Crambidae	<i>Syllepte</i> Solis22	Costa Rica: Guanacaste	ASHYF1500-11
<i>Alabagrus cocto</i>	DHJPAR0042356	Crambidae	<i>Syllepte</i> Solis22	Costa Rica: Alajuela	ASHYH120-11
<i>Alabagrus cocto</i>	DHJPAR0045784	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ACGAZ973-11
<i>Alabagrus cocto</i>	DHJPAR0048715	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGBA2257-12
<i>Alabagrus cocto</i>	DHJPAR0050361	Crambidae	<i>Syllepte</i> Solis22	Costa Rica: Alajuela	ACGAZ1675-12
<i>Alabagrus cocto</i>	DHJPAR0053598	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ASHYM2952-13
<i>Alabagrus cocto</i>	DHJPAR0053599	Crambidae	<i>Phostria vitrifera</i>	Costa Rica: Alajuela	ASHYM2953-13
<i>Alabagrus cocto</i>	DHJPAR0053601	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ASHYM2955-13
<i>Alabagrus cocto</i>	DHJPAR0053603	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Alajuela	ASHYM2957-13
<i>Alabagrus combos</i>	DHJPAR0009361	Crambidae	<i>Patania</i> Solis03	Costa Rica: Guanacaste	ASBR628-06
<i>Alabagrus combos</i>	DHJPAR0015530	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ASAG216-07
<i>Alabagrus combos</i>	DHJPAR0017276	Crambidae	<i>Conchylodes platinalis</i>	Costa Rica: Guanacaste	ASBD381-07
<i>Alabagrus combos</i>	DHJPAR0030609	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ASHYB1350-09
<i>Alabagrus combos</i>	DHJPAR0035295	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ASHYE1117-09
<i>Alabagrus combos</i>	DHJPAR0035529	Crambidae	<i>Pycnarmon</i> BioLep66	Costa Rica: Guanacaste	ASHYE1351-09
<i>Alabagrus combos</i>	DHJPAR0036341	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ASHYD1532-09
<i>Alabagrus combos</i>	DHJPAR0036376	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ASHYD1567-09
<i>Alabagrus combos</i>	DHJPAR0036377	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ASHYD1568-09
<i>Alabagrus combos</i>	DHJPAR0037877	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4622-10
<i>Alabagrus combos</i>	DHJPAR0037878	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4623-10
<i>Alabagrus combos</i>	DHJPAR0037879	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4624-10
<i>Alabagrus combos</i>	DHJPAR0037884	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4629-10
<i>Alabagrus combos</i>	DHJPAR0037891	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4636-10
<i>Alabagrus combos</i>	DHJPAR0037896	Crambidae	<i>Ategumia matutinalis</i> DHJ03	Costa Rica: Alajuela	ASHYC4641-10
<i>Alabagrus combos</i>	DHJPAR0037897	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	ASHYC4642-10
<i>Alabagrus combos</i>	DHJPAR0037898	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	ASHYC4643-10
<i>Alabagrus combos</i>	DHJPAR0037914	Crambidae	<i>Ategumia matutinalis</i> DHJ03	Costa Rica: Guanacaste	ASHYC4659-10
<i>Alabagrus combos</i>	DHJPAR0037929	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYC4674-10
<i>Alabagrus combos</i>	DHJPAR0037955	Crambidae	<i>Ategumia matutinalis</i> DHJ03	Costa Rica: Alajuela	ASHYC4700-10
<i>Alabagrus combos</i>	DHJPAR0037958	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYC4703-10
<i>Alabagrus combos</i>	DHJPAR0038807	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Alajuela	ASHYD2380-10
<i>Alabagrus combos</i>	DHJPAR0038812	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Alajuela	ASHYD2385-10
<i>Alabagrus combos</i>	DHJPAR0041185	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYF1091-11
<i>Alabagrus combos</i>	DHJPAR0041186	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Alajuela	ASHYF1092-11
<i>Alabagrus combos</i>	DHJPAR0041562, H7958	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYF1468-11
<i>Alabagrus combos</i>	DHJPAR0041583, H7981	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYF1489-11
<i>Alabagrus combos</i>	DHJPAR0041596, H7948	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYF1502-11
<i>Alabagrus combos</i>	DHJPAR0041598, H7823	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYF1504-11
<i>Alabagrus combos</i>	DHJPAR0042819	Crambidae	<i>Ategumia lotanalis</i> DHJ08	Costa Rica: Alajuela	ASHYH577-11
<i>Alabagrus combos</i>	DHJPAR0042826	Crambidae	<i>spilo</i> Janzen01 Janzen14DHJ02	Costa Rica: Alajuela	ASHYH584-11
<i>Alabagrus combos</i>	DHJPAR0046734	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Alajuela	ACGBA907-12
<i>Alabagrus combos</i>	DHJPAR0046735	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ACGBA908-12
<i>Alabagrus combos</i>	DHJPAR0046739	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Alajuela	ACGBA912-12
<i>Alabagrus combos</i>	DHJPAR0049475, H16752	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ACGBA3017-12
<i>Alabagrus combos</i>	DHJPAR0050368	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGAZ1682-12
<i>Alabagrus combos</i>	DHJPAR0050372	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ACGAZ1686-12
<i>Alabagrus combos</i>	DHJPAR0051350, H16026	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ACGBA3942-13
<i>Alabagrus combos</i>	DHJPAR0051354, H15917	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ACGBA3946-13

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus combos</i>	DHJPAR0051383, H16066	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ACGBA3975-13
<i>Alabagrus combos</i>	DHJPAR0051384, H16067	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ACGBA3976-13
<i>Alabagrus combos</i>	DHJPAR0053607	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYM2961-13
<i>Alabagrus combos</i>	DHJPAR0053635	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYM2989-13
<i>Alabagrus combos</i>	DHJPAR0054498	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	ASHYD3663-14
<i>Alabagrus combos</i>	DHJPAR0054506	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Alajuela	ASHYD3671-14
<i>Alabagrus combos</i>	DHJPAR0055974	Crambidae	<i>Ategumia matutinalis</i>	Costa Rica: Alajuela	ASHYH2711-14
<i>Alabagrus combos</i>	DHJPAR0055975	Crambidae	<i>Ategumia matutinalis</i>	Costa Rica: Alajuela	ASHYH2712-14
<i>Alabagrus combos</i>	DHJPAR0056358, H15821	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	MHMYC2438-15
<i>Alabagrus combos</i>	DHJPAR0057427	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	ACGBA5337-15
<i>Alabagrus combos</i>	DHJPAR0057437	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGBA5347-15
<i>Alabagrus combos</i>	DHJPAR0057438	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGBA5348-15
<i>Alabagrus combos</i>	DHJPAR0058538	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Alajuela	MHMYN8138-16
<i>Alabagrus cuna</i>	DHJPAR0042807	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYH565-11
<i>Alabagrus cuna</i>	DHJPAR0043025, H8836	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYH783-11
<i>Alabagrus cuna</i>	DHJPAR0043033, H4852	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYH791-11
<i>Alabagrus cuna</i>	DHJPAR0049045	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ACGBA2587-12
<i>Alabagrus cuna</i>	DHJPAR0049047	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ACGBA2589-12
<i>Alabagrus cuna</i>	DHJPAR0054521, H15804	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYD3686-14
<i>Alabagrus cuna</i>	DHJPAR0054525, H15916	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYD3690-14
<i>Alabagrus cuna</i>	DHJPAR0054535, H15930	Crambidae	<i>Neurophyseta</i> BioLep226	Costa Rica: Alajuela	ASHYD3700-14
<i>Alabagrus donharveyi</i>	DHJPAR0021163	Crambidae	<i>Coelorrhyncidia pandaralis</i>	Costa Rica: Alajuela	ASBC975-07
<i>Alabagrus donharveyi</i>	DHJPAR0021165	Crambidae	<i>Coelorrhyncidia pandaralis</i>	Costa Rica: Alajuela	ASBC977-07
<i>Alabagrus donharveyi</i>	DHJPAR0036688	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Alajuela	ASHYE1599-09
<i>Alabagrus donharveyi</i>	DHJPAR0036711	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Alajuela	ASHYE1622-09
<i>Alabagrus donharveyi</i>	DHJPAR0037930	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Alajuela	ASHYC4675-10
<i>Alabagrus donharveyi</i>	DHJPAR0045810	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Alajuela	ACGAZ999-11
<i>Alabagrus donharveyi</i>	DHJPAR0048734	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Guanacaste	ACGBA2276-12
<i>Alabagrus donharveyi</i>	DHJPAR0054512	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Guanacaste	ASHYD3677-14
<i>Alabagrus donharveyi</i>	DHJPAR0054513	Crambidae	<i>Coenostolopsis apicalis</i>	Costa Rica: Guanacaste	ASHYD3678-14
<i>Alabagrus donlafontainei</i>	DHJPAR0038923	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ASHYD2496-10
<i>Alabagrus donlafontainei</i>	DHJPAR0038927	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ASHYD2500-10
<i>Alabagrus donlafontainei</i>	DHJPAR0038928	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ASHYD2501-10
<i>Alabagrus donlafontainei</i>	DHJPAR0038929	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ASHYD2502-10
<i>Alabagrus donlafontainei</i>	DHJPAR0050137, H16759	Crambidae	musoBioLep01 BioLep448	Costa Rica: Guanacaste	ACGAZ1451-12
<i>Alabagrus donnai</i>	H1199			Costa Rica: Heredia	MF361681
<i>Alabagrus donnai</i>	H12192			Costa Rica: Heredia	MF361704
<i>Alabagrus ekchuah</i>	DHJPAR0015403	Crambidae	<i>Syllepte amando</i> DHJ02	Costa Rica: Guanacaste	ASAG089-07
<i>Alabagrus ekchuah</i>	DHJPAR0015404	Crambidae	<i>Pantographa suffusalis</i>	Costa Rica: Guanacaste	ASAG090-07
<i>Alabagrus ekchuah</i>	DHJPAR0015405	Crambidae	<i>Pantographa suffusalis</i>	Costa Rica: Guanacaste	ASAG091-07
<i>Alabagrus ekchuah</i>	DHJPAR0021168	Crambidae	<i>Syllepte amando</i> DHJ02	Costa Rica: Guanacaste	ASBC980-07
<i>Alabagrus englishi</i>	H12110			Costa Rica: Heredia	MF361702
<i>Alabagrus englishi</i>	INB0003765129			Costa Rica: San Jose	ASINH998-12
<i>Alabagrus englishi</i>	INBIOCRI002005193			Costa Rica: Guanacaste	ASINH1007-12
<i>Alabagrus fernandodiasi</i>	DHJPAR0009377	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Guanacaste	ASBR644-06
<i>Alabagrus fernandodiasi</i>	DHJPAR0017273	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Guanacaste	ASBD378-07
<i>Alabagrus fernandodiasi</i>	DHJPAR0021198	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Guanacaste	ASBC1010-07
<i>Alabagrus fernandodiasi</i>	DHJPAR0037883	Crambidae	<i>Rhectocraspeda</i> Janzen347	Costa Rica: Guanacaste	ASHYC4628-10
<i>Alabagrus fernandodiasi</i>	DHJPAR0037890	Crambidae	<i>Rhectocraspeda</i> Janzen347	Costa Rica: Guanacaste	ASHYC4635-10
<i>Alabagrus fernandodiasi</i>	DHJPAR0040215	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Guanacaste	ASHYE2382-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0041163	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Guanacaste	ASHYF1078-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0042804	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Guanacaste	ASHYH562-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0044991	Crambidae	spiloBioLep01 BioLep403	Costa Rica: Guanacaste	ACGAZ212-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0045799	Crambidae	spiloBioLep01 BioLep402	Costa Rica: Guanacaste	ACGAZ988-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0045800	Crambidae	spiloBioLep01 BioLep402	Costa Rica: Guanacaste	ACGAZ989-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0045801	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Guanacaste	ACGAZ990-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0045804	Crambidae	spiloBioLep01 BioLep403	Costa Rica: Guanacaste	ACGAZ993-11
<i>Alabagrus fernandodiasi</i>	DHJPAR0046750	Crambidae	<i>Rhectocraspeda periusalis</i>	Costa Rica: Alajuela	ACGBA923-12
<i>Alabagrus fernandodiasi</i>	DHJPAR0048731	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ACGBA2273-12
<i>Alabagrus fernandodiasi</i>	DHJPAR0048732	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ACGBA2274-12
<i>Alabagrus fernandodiasi</i>	DHJPAR0050360	Crambidae	<i>Desmia benealis</i> DHJ02	Costa Rica: Alajuela	ACGAZ1674-12
<i>Alabagrus fernandodiasi</i>	DHJPAR0052080	Crambidae	spiloBioLep01 BioLep414	Costa Rica: Alajuela	ASHYH1192-13
<i>Alabagrus fernandodiasi</i>	DHJPAR0052081	Crambidae	spiloBioLep01 BioLep414	Costa Rica: Alajuela	ASHYH1193-13
<i>Alabagrus fernandodiasi</i>	DHJPAR0052085	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYH1197-13
<i>Alabagrus fernandodiasi</i>	DHJPAR0052089	Crambidae	spiloBioLep01 BioLep311	Costa Rica: Alajuela	ASHYH1201-13
<i>Alabagrus fernandodiasi</i>	DHJPAR0052090	Crambidae	<i>Desmia</i> Janzen19	Costa Rica: Alajuela	ASHYH1202-13
<i>Alabagrus fernandodiasi</i>	DHJPAR0054488	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYD3653-14
<i>Alabagrus fernandodiasi</i>	DHJPAR0054490	Crambidae	spiloBioLep01 BioLep243	Costa Rica: Alajuela	ASHYD3655-14

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus fernandodiasi</i>	DHJPARG0055088	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYH1635-14
<i>Alabagrus fernandodiasi</i>	DHJPARG0055098	Crambidae	<i>Rhctocraspeda periusalis</i>	Costa Rica: Guanacaste	ASHYH1645-14
<i>Alabagrus fernandodiasi</i>	DHJPARG0055099	Crambidae	<i>Rhctocraspeda periusalis</i>	Costa Rica: Guanacaste	ASHYH1646-14
<i>Alabagrus fernandodiasi</i>	DHJPARG0055236, H15945	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYH1783-14
<i>Alabagrus fernandodiasi</i>	DHJPARG0055818	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYH2555-14
<i>Alabagrus fernandodiasi</i>	DHJPARG0056299	Crambidae	<i>Rhctocraspeda</i> Solis05	Costa Rica: Guanacaste	MHMYC2379-15
<i>Alabagrus fernandodiasi</i>	DHJPARG0057002, H15957	Crambidae	spiloBioLep01 BioLep415	Costa Rica: Alajuela	ACGBA4912-15
<i>Alabagrus genemonroei</i>	DHJPARG0015510	Crambidae	<i>Lamprosema</i> Dapkey23	Costa Rica: Alajuela	ASAG196-07
<i>Alabagrus iankitchingi</i>	DHJPARG0045001	Crambidae	<i>Phaedropsis</i> Solis350DHJ02	Costa Rica: Guanacaste	ACGAZ222-11
<i>Alabagrus iankitchingi</i>	DHJPARG0054545, H15841	Crambidae	<i>Phaedropsis</i> Solis350DHJ02	Costa Rica: Guanacaste	ASHYD3710-14
<i>Alabagrus imitatus</i>	H10807			USA: Missouri	MF361690
<i>Alabagrus imitatus</i>	H10850			USA: Missouri	MF361691
<i>Alabagrus imitatus</i>	H10874			USA: Missouri	MF361692
<i>Alabagrus imitatus</i>	H10876			USA: Missouri	MF361693
<i>Alabagrus imitatus</i>	H10882			USA: Missouri	MF361694
<i>Alabagrus imitatus</i>	H10888			USA: Missouri	MF361695
<i>Alabagrus isidrochaconi</i>	DHJPARG0009405	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASBR672-06
<i>Alabagrus isidrochaconi</i>	DHJPARG0015520	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASAG206-07
<i>Alabagrus isidrochaconi</i>	DHJPARG0015523	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASAG209-07
<i>Alabagrus isidrochaconi</i>	DHJPARG0015525	Crambidae	<i>Eulepte</i> Janzen03	Costa Rica: Guanacaste	ASAG211-07
<i>Alabagrus isidrochaconi</i>	DHJPARG0052674	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2028-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0052675	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2029-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0052676	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2030-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0052683	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2037-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0052684	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2038-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053608	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2962-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053609	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2963-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053624	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2978-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053626	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2980-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053627	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2981-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053629	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2983-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053630	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2984-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053631	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2985-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053633	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2987-13
<i>Alabagrus isidrochaconi</i>	DHJPARG0053634	Crambidae	<i>Glyphodes sibillalis</i> DHJ01	Costa Rica: Guanacaste	ASHYM2988-13
<i>Alabagrus ixtilton</i>	H7357			Mexico: Jalisco	ATRMK377-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0037886	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Guanacaste	ASHYC4631-10
<i>Alabagrus jackiemillerae</i>	DHJPARG0037895	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ASHYC4640-10
<i>Alabagrus jackiemillerae</i>	DHJPARG0038805	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYD2378-10
<i>Alabagrus jackiemillerae</i>	DHJPARG0041165	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Guanacaste	ASHYF1080-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0045798	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGAZ987-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0045802	Crambidae	spiloBioLep01 BioLep403	Costa Rica: Guanacaste	ACGAZ991-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0045803	Crambidae	spiloBioLep01 BioLep403	Costa Rica: Guanacaste	ACGAZ992-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0045805	Crambidae	spiloBioLep01 BioLep403	Costa Rica: Guanacaste	ACGAZ994-11
<i>Alabagrus jackiemillerae</i>	DHJPARG0046753	Crambidae	<i>Pilocrocis purpurascens</i>	Costa Rica: Alajuela	ACGBA926-12
<i>Alabagrus jackiemillerae</i>	DHJPARG0048729	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ACGBA2271-12
<i>Alabagrus jackiemillerae</i>	DHJPARG0051668	Crambidae	<i>Phostria</i> Janzen05	Costa Rica: Alajuela	ACGBA4260-13
<i>Alabagrus jackiemillerae</i>	DHJPARG0052078	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYH1190-13
<i>Alabagrus jackiemillerae</i>	DHJPARG0052682	Crambidae	<i>Patania</i> Solis03	Costa Rica: Alajuela	ASHYM2036-13
<i>Alabagrus jackiemillerae</i>	DHJPARG0053593	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYM2947-13
<i>Alabagrus jackiemillerae</i>	DHJPARG0053596	Crambidae	<i>Herpetogramma</i> Solis11	Costa Rica: Alajuela	ASHYM2950-13
<i>Alabagrus jackiemillerae</i>	DHJPARG0054491	Crambidae	<i>Rhctocraspeda</i> Janzen347	Costa Rica: Alajuela	ASHYD3656-14
<i>Alabagrus jackiemillerae</i>	DHJPARG0054492	Crambidae	<i>Rhctocraspeda</i> Janzen347	Costa Rica: Alajuela	ASHYD3657-14
<i>Alabagrus jackiemillerae</i>	DHJPARG0057439	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ACGBA5349-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057440	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ACGBA5350-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057441	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ACGBA5351-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057445	Crambidae	<i>Rhctocraspeda</i> Solis05	Costa Rica: Alajuela	ACGBA5355-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057446	Crambidae	<i>Phostria</i> Janzen05	Costa Rica: Alajuela	ACGBA5356-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057448	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Alajuela	ACGBA5358-15
<i>Alabagrus jackiemillerae</i>	DHJPARG0057450	Crambidae	<i>Rhctocraspeda</i> Solis05	Costa Rica: Alajuela	ACGBA5360-15
<i>Alabagrus janzeni</i>	DHJPARG0009348	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBR615-06
<i>Alabagrus janzeni</i>	DHJPARG0009354	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR621-06
<i>Alabagrus janzeni</i>	DHJPARG0009355	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR622-06
<i>Alabagrus janzeni</i>	DHJPARG0009356	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR623-06
<i>Alabagrus janzeni</i>	DHJPARG0009362	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR629-06
<i>Alabagrus janzeni</i>	DHJPARG0009374	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR641-06
<i>Alabagrus janzeni</i>	DHJPARG0009375	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR642-06
<i>Alabagrus janzeni</i>	DHJPARG0009376	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR643-06

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus janzeni</i>	DHJPARG0009400	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR667-06
<i>Alabagrus janzeni</i>	DHJPARG0009403	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR670-06
<i>Alabagrus janzeni</i>	DHJPARG0009433	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBR700-06
<i>Alabagrus janzeni</i>	DHJPARG0009434	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBR701-06
<i>Alabagrus janzeni</i>	DHJPARG0010083	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBC667-06
<i>Alabagrus janzeni</i>	DHJPARG0010509	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBC668-06
<i>Alabagrus janzeni</i>	DHJPARG0010510	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBC669-06
<i>Alabagrus janzeni</i>	DHJPARG0015384	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG070-07
<i>Alabagrus janzeni</i>	DHJPARG0015493	Crambidae	<i>Eulepte Solis15</i>	Costa Rica: Guanacaste	ASAG179-07
<i>Alabagrus janzeni</i>	DHJPARG0015494	Crambidae	<i>Eulepte Solis15</i>	Costa Rica: Guanacaste	ASAG180-07
<i>Alabagrus janzeni</i>	DHJPARG0015495	Crambidae	<i>Eulepte Solis15</i>	Costa Rica: Guanacaste	ASAG181-07
<i>Alabagrus janzeni</i>	DHJPARG0015496	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG182-07
<i>Alabagrus janzeni</i>	DHJPARG0015497	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG183-07
<i>Alabagrus janzeni</i>	DHJPARG0015498	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG184-07
<i>Alabagrus janzeni</i>	DHJPARG0015499	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG185-07
<i>Alabagrus janzeni</i>	DHJPARG0015511	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASAG197-07
<i>Alabagrus janzeni</i>	DHJPARG0015512	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASAG198-07
<i>Alabagrus janzeni</i>	DHJPARG0015513	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASAG199-07
<i>Alabagrus janzeni</i>	DHJPARG0015515	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASAG201-07
<i>Alabagrus janzeni</i>	DHJPARG0016416	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASTAP445-06
<i>Alabagrus janzeni</i>	DHJPARG0016417	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASTAP446-06
<i>Alabagrus janzeni</i>	DHJPARG0016428	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASTAP457-06
<i>Alabagrus janzeni</i>	DHJPARG0016429	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASTAP458-06
<i>Alabagrus janzeni</i>	DHJPARG0016430	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASTAP459-06
<i>Alabagrus janzeni</i>	DHJPARG0016431	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASTAP460-06
<i>Alabagrus janzeni</i>	DHJPARG0016432	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASTAP461-06
<i>Alabagrus janzeni</i>	DHJPARG0016926	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBR899-07
<i>Alabagrus janzeni</i>	DHJPARG0016928	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBR901-07
<i>Alabagrus janzeni</i>	DHJPARG0016930	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR903-07
<i>Alabagrus janzeni</i>	DHJPARG0021149	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBC961-07
<i>Alabagrus janzeni</i>	DHJPARG0021208	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASBC1020-07
<i>Alabagrus janzeni</i>	DHJPARG0021211	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBC1023-07
<i>Alabagrus janzeni</i>	DHJPARG0029179	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE586-09
<i>Alabagrus janzeni</i>	DHJPARG0036363	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYD1554-09
<i>Alabagrus janzeni</i>	DHJPARG0036365	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYD1556-09
<i>Alabagrus janzeni</i>	DHJPARG0036366	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYD1557-09
<i>Alabagrus janzeni</i>	DHJPARG0036697	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE1608-09
<i>Alabagrus janzeni</i>	DHJPARG0036698	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE1609-09
<i>Alabagrus janzeni</i>	DHJPARG0036699	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE1610-09
<i>Alabagrus janzeni</i>	DHJPARG0036712	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE1623-09
<i>Alabagrus janzeni</i>	DHJPARG0036716	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYE1627-09
<i>Alabagrus janzeni</i>	DHJPARG0036718	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYE1629-09
<i>Alabagrus janzeni</i>	DHJPARG0037190	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYE2024-10
<i>Alabagrus janzeni</i>	DHJPARG0037902	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYC4647-10
<i>Alabagrus janzeni</i>	DHJPARG0037922	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYC4667-10
<i>Alabagrus janzeni</i>	DHJPARG0041166	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYF1081-11
<i>Alabagrus janzeni</i>	DHJPARG0041167	Crambidae	<i>Eulepte Janzen02</i>	Costa Rica: Guanacaste	ASHYF1082-11
<i>Alabagrus janzeni</i>	DHJPARG0041195	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYF1101-11
<i>Alabagrus janzeni</i>	DHJPARG0041197	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYF1103-11
<i>Alabagrus janzeni</i>	DHJPARG0041198	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYF1104-11
<i>Alabagrus janzeni</i>	DHJPARG0041576, H7674	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYF1482-11
<i>Alabagrus janzeni</i>	DHJPARG0044992	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ213-11
<i>Alabagrus janzeni</i>	DHJPARG0044993	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ214-11
<i>Alabagrus janzeni</i>	DHJPARG0044994	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ215-11
<i>Alabagrus janzeni</i>	DHJPARG0044995	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ216-11
<i>Alabagrus janzeni</i>	DHJPARG0044996	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ217-11
<i>Alabagrus janzeni</i>	DHJPARG0044997	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ACGAZ218-11
<i>Alabagrus janzeni</i>	DHJPARG0044998	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ACGAZ219-11
<i>Alabagrus janzeni</i>	DHJPARG0048724	Crambidae	<i>Eulepte Solis15</i>	Costa Rica: Guanacaste	ACGBA2266-12
<i>Alabagrus janzeni</i>	DHJPARG0049940	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ACGAZ1254-12
<i>Alabagrus janzeni</i>	DHJPARG0050363	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ACGAZ1677-12
<i>Alabagrus janzeni</i>	DHJPARG0050371	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ACGAZ1685-12
<i>Alabagrus janzeni</i>	DHJPARG0052083	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1195-13
<i>Alabagrus janzeni</i>	DHJPARG0053587	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYM2941-13
<i>Alabagrus janzeni</i>	DHJPARG0053588	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYM2942-13
<i>Alabagrus janzeni</i>	DHJPARG0053589	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYM2943-13
<i>Alabagrus janzeni</i>	DHJPARG0053590	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYM2944-13
<i>Alabagrus janzeni</i>	DHJPARG0053622	Crambidae	<i>Eulepte Janzen06</i>	Costa Rica: Guanacaste	ASHYM2976-13

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus janzeni</i>	DHJPAR0053623	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASHYM2977-13
<i>Alabagrus janzeni</i>	DHJPAR0054487	Crambidae	<i>Eulepte</i> Janzen06	Costa Rica: Guanacaste	ASHYD3652-14
<i>Alabagrus janzeni</i>	DHJPAR0055089	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1636-14
<i>Alabagrus janzeni</i>	DHJPAR0055090	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1637-14
<i>Alabagrus janzeni</i>	DHJPAR0055091	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1638-14
<i>Alabagrus janzeni</i>	DHJPAR0055092	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1639-14
<i>Alabagrus janzeni</i>	DHJPAR0055093	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1640-14
<i>Alabagrus janzeni</i>	DHJPAR0055094	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1641-14
<i>Alabagrus janzeni</i>	DHJPAR0055095	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1642-14
<i>Alabagrus janzeni</i>	DHJPAR0055096	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1643-14
<i>Alabagrus janzeni</i>	DHJPAR0055097	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	ASHYH1644-14
<i>Alabagrus janzeni</i>	DHJPAR0058537	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	MHMYN8137-16
<i>Alabagrus janzeni</i>	DHJPAR0058539	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	MHMYN8139-16
<i>Alabagrus janzeni</i>	DHJPAR0058542	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Alajuela	MHMYN8142-16
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0009353	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASBR620-06
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0009357	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASBR624-06
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0009373	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASBR640-06
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0009406	Crambidae	<i>Conchylodes arcifera</i>	Costa Rica: Guanacaste	ASBR673-06
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015316	Crambidae	<i>Eulepte concordalis</i>	Costa Rica: Guanacaste	ASAG002-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015328	Crambidae	<i>Phaedropsis cernalis</i>	Costa Rica: Guanacaste	ASAG014-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015329	Crambidae	<i>Herpetogramma semilaniata</i>	Costa Rica: Guanacaste	ACGBA6164-16
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015330	Crambidae	<i>Herpetogramma semilaniata</i>	Costa Rica: Guanacaste	ASAG016-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015387	Crambidae	<i>Ategumia lotanalis</i> DHJ07	Costa Rica: Guanacaste	ASAG073-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015521	Crambidae	<i>Conchylodes arcifera</i>	Costa Rica: Guanacaste	ASAG207-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0015522	Crambidae	<i>Microthyris anormalis</i>	Costa Rica: Guanacaste	ASAG208-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0021157	Crambidae	<i>Conchylodes platinalis</i>	Costa Rica: Guanacaste	ASBC969-07
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0030388	Crambidae	<i>Ategumia lotanalis</i>	Costa Rica: Guanacaste	ASHYB1127-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036349	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASHYD1540-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036353	Crambidae	<i>Conchylodes platinalis</i>	Costa Rica: Guanacaste	ASHYD1544-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036356	Crambidae	<i>Ategumia lotanalis</i> DHJ09	Costa Rica: Alajuela	ASHYD1547-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036374	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYD1565-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036683	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYE1594-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036685	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYE1596-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036696	Crambidae	<i>Patania Solis</i> 03	Costa Rica: Guanacaste	ASHYE1607-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036700	Crambidae	<i>Phostria samealis</i>	Costa Rica: Alajuela	ASHYE1611-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036703	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYE1614-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0036717	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYE1628-09
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037187	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASHYE2021-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037900	Crambidae	<i>Hyalorista exuvialis</i> DHJ02	Costa Rica: Alajuela	ASHYC4645-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037917	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYC4662-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037948	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYC4693-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037950	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYC4695-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037951	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASAG283-16
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037953	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ASHYC4698-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0037956	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASHYC4701-10
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0041196	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ASHYF1102-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0041603, H7902	Crambidae	<i>Herpetogramma Solis</i> 11	Costa Rica: Guanacaste	ASHYF1509-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0042824	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ASHYH582-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0044999	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ACGAZ220-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045000	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ACGAZ221-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045003	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ACGAZ224-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045004	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ225-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045005	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ226-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045006	Crambidae	<i>Herpetogramma phaeopteralis</i>	Costa Rica: Guanacaste	ACGAZ227-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045010	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ231-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045011	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ232-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045012	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ233-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045013	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ234-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045014	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ235-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045015	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Guanacaste	ACGAZ236-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045783	Crambidae	<i>Herpetogramma Solis</i> 11	Costa Rica: Guanacaste	ACGAZ972-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0045786	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ACGAZ975-11
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0046741	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ACGBA914-12
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0049939	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Guanacaste	ACGAZ1253-12
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0050065	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ACGAZ1379-12
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0050370	Crambidae	<i>Patania Solis</i> 03	Costa Rica: Alajuela	ACGAZ1684-12
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0050932	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ACGBA3524-13
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0052689	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYM2043-13

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0052693	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYM2047-13
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0053613	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ASHYM2967-13
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0055979	Crambidae	<i>Rhectocraspeda</i> Solis05	Costa Rica: Alajuela	ASHYH2716-14
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0055989, H15845	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYH2726-14
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0055990, H16019	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYH2727-14
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0056295	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYH2552-14
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0056296	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	MHMYC2376-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0056298	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	MHMYC2378-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0056994, H15959	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGBA4904-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0056999, H15937	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGBA4909-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0057430	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ACGBA5340-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0057431	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ACGBA5341-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0057738, H16751	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYB2672-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0057745, H16735	Crambidae	<i>Eulepte alialis</i>	Costa Rica: Alajuela	ASHYB2679-15
<i>Alabagrus jeanfrancoislandryi</i>	DHJPAR0059747	Crambidae	<i>Pilocrocis ramentalis</i>	Costa Rica: Guanacaste	ACGBA6165-16
<i>Alabagrus jeanfrancoislandryi</i>	H7628			Mexico: Yucatan	ATRMK411-11
<i>Alabagrus jeanfrancoislandryi</i>	H7630			Mexico: Yucatan	ATRMK412-11
<i>Alabagrus jeanfrancoislandryi</i>	H7631			Mexico: Yucatan	ATRMK413-11
<i>Alabagrus jeanfrancoislandryi</i>	H831			Mexico: Yucatan	ATRMK246-11
<i>Alabagrus jeanfrancoislandryi</i>	H873			Mexico: Yucatan	ATRMK247-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009416	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR683-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009424	Crambidae	<i>Desmia</i> Janzen09	Costa Rica: Alajuela	ASBR691-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009425	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR692-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009427	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR694-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009428	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR695-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009429	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR696-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0009430	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASBR697-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0010513	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Guanacaste	ASBC672-06
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0015507	Crambidae	<i>Desmia ploralis</i> DHJ02	Costa Rica: Guanacaste	ASAG193-07
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0015508	Crambidae	<i>Desmia</i> Janzen09	Costa Rica: Alajuela	ASAG194-07
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0015509	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Guanacaste	ASAG195-07
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0023534	Crambidae	<i>Desmia ploralis</i> DHJ06	Costa Rica: Guanacaste	ASHYM286-08
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0028030	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYE267-08
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0028030	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYE267-08
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0036687	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYE1598-09
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0036702	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYE1613-09
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0036704	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYE1615-09
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0036710	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYE1621-09
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037893	Crambidae	<i>Desmia</i> Solis19	Costa Rica: Guanacaste	ASHYC4638-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037901	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYC4646-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037903	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYC4648-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037904	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYC4649-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037905	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYC4650-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037918	Crambidae	<i>Desmia ploralis</i> DHJ04	Costa Rica: Alajuela	ASHYC4663-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037921	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYC4666-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037934	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYC4679-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0037954	Crambidae	<i>Desmia ploralis</i> DHJ04	Costa Rica: Alajuela	ASHYC4699-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0038804	Crambidae	<i>Desmia ploralis</i> DHJ04	Costa Rica: Alajuela	ASHYD2377-10
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0041190	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYF1096-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0041193	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYF1099-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0041204	Crambidae	<i>Desmia</i> Solis100	Costa Rica: Alajuela	ASHYF1110-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0041206	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYF1112-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0044985	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGAZ206-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0044987	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGAZ208-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0044989	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGAZ210-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0045796	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Guanacaste	ACGAZ985-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0045797	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Guanacaste	ACGAZ986-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0045806	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Alajuela	ACGAZ995-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0045807	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Guanacaste	ACGAZ996-11
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0046751	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Guanacaste	ACGBA924-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0046752	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Guanacaste	ACGBA925-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0048716	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGBA2258-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0048717	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGBA2259-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0048730	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ACGBA2272-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0049656	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYB2450-12
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0051346, H15927	Crambidae	<i>Desmia ploralis</i> DHJ01	Costa Rica: Guanacaste	ACGBA3938-13
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0052084	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYH1196-13
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0052690	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYM2044-13

Appendix 1 Continued.

Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0052691	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYM2045-13
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0052695	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYM2049-13
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0053592	Crambidae	<i>Desmia octomaculalis</i>	Costa Rica: Alajuela	ASHYM2946-13
<i>Alabagrus jeanmariecadoui</i>	DHJPAR0055969	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYH2706-14
<i>Alabagrus jennyphillipsae</i>	DHJPAR0035225	Crambidae	<i>Leucochromodes melusinalis</i> DHJ01	Costa Rica: Alajuela	ASHYE1047-09
<i>Alabagrus jennyphillipsae</i>	DHJPAR0037923	Crambidae	<i>Leucochromodes melusinalis</i> DHJ01	Costa Rica: Guanacaste	ASHYC4668-10
<i>Alabagrus jennyphillipsae</i>	DHJPAR0039523, H8033	Crambidae	<i>Leucochromodes melusinalis</i> DHJ01	Costa Rica: Guanacaste	ASHYB1811-10
<i>Alabagrus jimmilleri</i>	DHJPAR0009372	Crambidae	<i>Palpusia</i> Janzen02	Costa Rica: Guanacaste	ASBR639-06
<i>Alabagrus jimmilleri</i>	DHJPAR0036373	Crambidae	<i>Lygropia tripunctata</i> DHJ04	Costa Rica: Alajuela	ASHYD1564-09
<i>Alabagrus jimmilleri</i>	DHJPAR0040337, H8022	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYE2473-11
<i>Alabagrus jimmilleri</i>	DHJPAR0049046	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ACGBA2588-12
<i>Alabagrus jimmilleri</i>	DHJPAR0049938	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ1252-12
<i>Alabagrus jimmilleri</i>	DHJPAR0049946	Crambidae	<i>Microthyris anormalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ1260-12
<i>Alabagrus johnbrowni</i>	DHJPAR0009364	Crambidae	<i>Pileosoma thialis</i> DHJ03	Costa Rica: Alajuela	ASBR631-06
<i>Alabagrus johnbrowni</i>	DHJPAR0009378	Crambidae	<i>Desmia tages</i>	Costa Rica: Guanacaste	ASBR645-06
<i>Alabagrus johnbrowni</i>	DHJPAR0009407	Crambidae	<i>Pileosoma thialis</i> DHJ03	Costa Rica: Alajuela	ASBR674-06
<i>Alabagrus johnbrowni</i>	DHJPAR0015516	Crambidae	<i>Pileosoma thialis</i> DHJ03	Costa Rica: Alajuela	ASAG202-07
<i>Alabagrus johnbrowni</i>	DHJPAR0015517	Crambidae	<i>Pileosoma thialis</i> DHJ03	Costa Rica: Alajuela	ASAG203-07
<i>Alabagrus johnbrowni</i>	DHJPAR0017277	Crambidae	<i>Pileosoma thialis</i> DHJ03	Costa Rica: Guanacaste	ASBD382-07
<i>Alabagrus johnbrowni</i>	DHJPAR0035226	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYE1048-09
<i>Alabagrus johnbrowni</i>	DHJPAR0035954	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYD1335-09
<i>Alabagrus johnbrowni</i>	DHJPAR0036335	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYD1526-09
<i>Alabagrus johnbrowni</i>	DHJPAR0036336	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYD1527-09
<i>Alabagrus johnbrowni</i>	DHJPAR0036375	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYD1566-09
<i>Alabagrus johnbrowni</i>	DHJPAR0036707	Crambidae	<i>Phostria</i> Janzen05	Costa Rica: Alajuela	ASHYE1618-09
<i>Alabagrus johnbrowni</i>	DHJPAR0040069, H6492	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Guanacaste	ASHYE2244-11
<i>Alabagrus johnbrowni</i>	DHJPAR0041591, H7910	Crambidae	<i>Desmia tages</i>	Costa Rica: Alajuela	ASHYF1497-11
<i>Alabagrus johnbrowni</i>	DHJPAR0045029	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ACGAZ250-11
<i>Alabagrus johnbrowni</i>	DHJPAR0045058	Crambidae	<i>Desmia</i> BioLep09	Costa Rica: Alajuela	ACGAZ279-11
<i>Alabagrus johnbrowni</i>	DHJPAR0045059	Crambidae	<i>Desmia</i> BioLep09	Costa Rica: Alajuela	ACGAZ280-11
<i>Alabagrus johnbrowni</i>	DHJPAR0045782	Crambidae	<i>Desmia tages</i>	Costa Rica: Alajuela	ACGAZ971-11
<i>Alabagrus johnbrowni</i>	DHJPAR0046748	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ACGBA921-12
<i>Alabagrus johnbrowni</i>	DHJPAR0046749	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ACGBA922-12
<i>Alabagrus johnbrowni</i>	DHJPAR0046754	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ACGBA927-12
<i>Alabagrus johnbrowni</i>	DHJPAR0046755	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ACGBA928-12
<i>Alabagrus johnbrowni</i>	DHJPAR0051669	Crambidae	<i>Lamprosema</i> Dapkey23	Costa Rica: Alajuela	ACGBA4261-13
<i>Alabagrus johnbrowni</i>	DHJPAR0052086	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYH1198-13
<i>Alabagrus johnbrowni</i>	DHJPAR0054531, H15910	Crambidae	<i>Pileosoma thialis</i>	Costa Rica: Alajuela	ASHYD3696-14
<i>Alabagrus johnbrowni</i>	DHJPAR0055980	Crambidae	<i>Desmia</i> BioLep42	Costa Rica: Alajuela	ASHYH2717-14
<i>Alabagrus johnbrowni</i>	DHJPAR0056297	Crambidae	<i>Desmia</i> BioLep19	Costa Rica: Guanacaste	MHMYC2377-15
<i>Alabagrus johnbrowni</i>	DHJPAR0056357, H15823	Crambidae	<i>Desmia</i> BioLep19	Costa Rica: Alajuela	MHMYC2437-15
<i>Alabagrus johnbrowni</i>	DHJPAR0057743, H16755	Crambidae	<i>Desmia</i> BioLep42	Costa Rica: Alajuela	ASHYB2677-15
<i>Alabagrus karensharkeyae</i>	DHJPAR0010196	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASBC477-06
<i>Alabagrus karensharkeyae</i>	DHJPAR0021170	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASBC982-07
<i>Alabagrus karensharkeyae</i>	DHJPAR0036348	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYD1539-09
<i>Alabagrus karensharkeyae</i>	DHJPAR0036706	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYE1617-09
<i>Alabagrus karensharkeyae</i>	DHJPAR0036715	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYE1626-09
<i>Alabagrus karensharkeyae</i>	DHJPAR0041200	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYF1106-11
<i>Alabagrus karensharkeyae</i>	DHJPAR0045002	Crambidae	<i>Microthyris prolongalis</i> DHJ02	Costa Rica: Guanacaste	ACGAZ223-11
<i>Alabagrus karensharkeyae</i>	DHJPAR0046742	Crambidae	<i>Palpusia</i> Janzen02	Costa Rica: Alajuela	ACGBA915-12
<i>Alabagrus karensharkeyae</i>	DHJPAR0052904	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Alajuela	ASHYM2258-13
<i>Alabagrus karensharkeyae</i>	DHJPAR0055978	Crambidae	<i>Palpusia</i> Janzen02	Costa Rica: Alajuela	ASHYH2715-14
<i>Alabagrus kaydodgeae</i>	DHJPAR0053007, H16737	Crambidae	<i>Ategumia</i> Solis01	Costa Rica: Guanacaste	ASHYM2361-13
<i>Alabagrus kaydodgeae</i>	H1200			Costa Rica: Heredia	MF361682
<i>Alabagrus kaydodgeae</i>	INBIOCRI000737989			Costa Rica: Limón	ASINH1036-12
<i>Alabagrus keithwillmotti</i>	DHJPAR0007207	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASTAV449-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009385	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR652-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009386	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR653-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009387	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR654-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009388	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR655-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009389	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR656-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0009390	Crambidae	<i>Pilocrocis</i> Solis20	Costa Rica: Guanacaste	ASBR657-06
<i>Alabagrus keithwillmotti</i>	DHJPAR0035227	Crambidae	spilobiol01 biol0498	Costa Rica: Guanacaste	ASHYE1049-09
<i>Alabagrus keithwillmotti</i>	DHJPAR0036339	Crambidae	<i>Asturodes fimbriauralis</i> DHJ02	Costa Rica: Guanacaste	ASHYD1530-09
<i>Alabagrus keithwillmotti</i>	DHJPAR0037907	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4652-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0037909	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4654-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0037910	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4655-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0037911	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4656-10

Appendix 1 Continued.


Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus keithwillmotti</i>	DHJPAR0037912	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4657-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0037913	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYC4658-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0038808	Crambidae	<i>Phostria</i> Janzen03	Costa Rica: Guanacaste	ASHYD2381-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0038809	Crambidae	<i>Asturodes fimbriauralis</i> DHJ02	Costa Rica: Guanacaste	ASHYD2382-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0039357	Crambidae	<i>Desmia octomaculalis</i>	Costa Rica: Guanacaste	ASHYB1645-10
<i>Alabagrus keithwillmotti</i>	DHJPAR0045789	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Guanacaste	ACGAZ978-11
<i>Alabagrus keithwillmotti</i>	DHJPAR0045808	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Guanacaste	ACGAZ997-11
<i>Alabagrus keithwillmotti</i>	DHJPAR0045809	Crambidae	<i>Herpetogramma</i> Solis10	Costa Rica: Guanacaste	ACGAZ998-11
<i>Alabagrus keithwillmotti</i>	DHJPAR0050868	Crambidae	<i>Desmia</i> Janzen14	Costa Rica: Guanacaste	ACGBA3460-13
<i>Alabagrus keithwillmotti</i>	DHJPAR0052730	Crambidae	<i>Desmia</i> Janzen07	Costa Rica: Guanacaste	ASHYM2084-13
<i>Alabagrus lindapitkinae</i>	DHJPAR0052082	Crambidae	<i>Omiodes</i> Janzen05	Costa Rica: Alajuela	ASHYH1194-13
<i>Alabagrus lindapitkinae</i>	DHJPAR0054497	Crambidae	<i>Omiodes</i> Janzen05	Costa Rica: Alajuela	ASHYD3662-14
<i>Alabagrus maculipes</i>	DHJPAR0036337	Crambidae	<i>Desmia ploralis</i> DHJ02	Costa Rica: Guanacaste	ASHYD1528-09
<i>Alabagrus maculipes</i>	DHJPAR0036725	Crambidae	<i>Syngamia florella</i>	Costa Rica: Guanacaste	ASHYE1636-09
<i>Alabagrus maculipes</i>	DHJPAR0037188	Crambidae	<i>Salbia haemorrhoidalis</i>	Costa Rica: Alajuela	ASHYE2022-10
<i>Alabagrus maculipes</i>	DHJPAR0040212	Crambidae	<i>Palpusia</i> Janzen02	Costa Rica: Guanacaste	ASHYE2379-11
<i>Alabagrus maculipes</i>	DHJPAR0040213	Crambidae	<i>Palpusia</i> Janzen02	Costa Rica: Guanacaste	ASHYE2380-11
<i>Alabagrus maculipes</i>	DHJPAR0040329, H8023	Crambidae	<i>Herpetogramma semilaniata</i>	Costa Rica: Guanacaste	ASHYE2465-11
<i>Alabagrus maculipes</i>	DHJPAR0041150	Crambidae	<i>Herpetogramma</i> Janzen04	Costa Rica: Guanacaste	ASHYF1065-11
<i>Alabagrus maculipes</i>	DHJPAR0041151	Crambidae	<i>Herpetogramma</i> Janzen04	Costa Rica: Guanacaste	ASHYF1066-11
<i>Alabagrus maculipes</i>	DHJPAR0048735	Crambidae	<i>Pilocrocis ramentalis</i>	Costa Rica: Guanacaste	ACGBA2277-12
<i>Alabagrus maculipes</i>	DHJPAR0048736	Crambidae	<i>Pilocrocis ramentalis</i>	Costa Rica: Guanacaste	ACGBA2278-12
<i>Alabagrus maculipes</i>	DHJPAR0056292	Crambidae	<i>Herpetogramma salbialis</i>	Costa Rica: Alajuela	ASHYH2549-14
<i>Alabagrus maculipes</i>	H14909			Mexico: Sonora	MF361709
<i>Alabagrus maculipes</i>	H15106			Mexico: Sonora	MF361713
<i>Alabagrus maculipes</i>	H6020			Mexico: Jalisco	ATRMK370-11
<i>Alabagrus malcolmscoblei</i>	DHJPAR0048718	Crambidae	<i>Phostria latiapicalis</i>	Costa Rica: Alajuela	ACGBA2260-12
<i>Alabagrus marcepesteini</i>	DHJPAR0041184	Crambidae	crambidJanzen01 Janzen21	Costa Rica: Alajuela	ASHYF1090-11
<i>Alabagrus marcepesteini</i>	DHJPAR0041192	Crambidae	crambidJanzen01 Janzen21	Costa Rica: Alajuela	ASHYF1098-11
<i>Alabagrus mariaheikkilae</i>	DHJPAR0057004, H15960	Crambidae	<i>Salbia cassidalis</i> DHJ02	Costa Rica: Alajuela	ACGBA6199-16
<i>Alabagrus markmetzi</i>	DHJPAR0009359	Crambidae	<i>Syllepte aechmisalis</i>	Costa Rica: Guanacaste	ASBR626-06
<i>Alabagrus markmetzi</i>	DHJPAR0009363	Crambidae	<i>Syllepte aechmisalis</i>	Costa Rica: Guanacaste	ASBR630-06
<i>Alabagrus markmetzi</i>	DHJPAR0009417	Crambidae	<i>Syllepte aechmisalis</i>	Costa Rica: Guanacaste	ASBR684-06
<i>Alabagrus markmetzi</i>	DHJPAR0015388	Crambidae	<i>Syllepte aechmisalis</i>	Costa Rica: Guanacaste	ASAG074-07
<i>Alabagrus markmetzi</i>	DHJPAR0046737	Crambidae	<i>Syllepte aechmisalis</i>	Costa Rica: Guanacaste	ACGBA910-12
<i>Alabagrus masneri</i>	H1197			Costa Rica: Cartago	MF361680
<i>Alabagrus nickgrishini</i>	DHJPAR0036347	Crambidae	<i>Microthyris alvinalis</i>	Costa Rica: Guanacaste	ASHYD1538-09
<i>Alabagrus nickgrishini</i>	DHJPAR0037892	Crambidae	<i>Microthyris alvinalis</i>	Costa Rica: Guanacaste	ASHYC4637-10
<i>Alabagrus nickgrishini</i>	DHJPAR0045022	Crambidae	<i>Microthyris alvinalis</i>	Costa Rica: Guanacaste	ACGAZ243-11
<i>Alabagrus nickgrishini</i>	DHJPAR0050366	Crambidae	<i>Patania</i> Solis02	Costa Rica: Guanacaste	ACGAZ1680-12
<i>Alabagrus nickgrishini</i>	DHJPAR0050367	Crambidae	<i>Patania</i> Solis02	Costa Rica: Guanacaste	ACGAZ1681-12
<i>Alabagrus nickgrishini</i>	DHJPAR0057425	Crambidae	<i>Erilusa leucoplagialis</i>	Costa Rica: Guanacaste	ACGBA5335-15
<i>Alabagrus nicoya</i>	DHJPAR0015399	Crambidae	<i>Dichogama colotha</i>	Costa Rica: Guanacaste	ASAG085-07
<i>Alabagrus nicoya</i>	DHJPAR0015400	Crambidae	<i>Dichogama colotha</i>	Costa Rica: Guanacaste	ASAG086-07
<i>Alabagrus nicoya</i>	DHJPAR0016931	Crambidae	<i>Dichogama colotha</i>	Costa Rica: Guanacaste	ASBR904-07
<i>Alabagrus nicoya</i>	DHJPAR0022193	Crambidae	<i>Dichogama colotha</i>	Costa Rica: Guanacaste	ASTAT1331-07
<i>Alabagrus nicoya</i>	DHJPAR0049650	Crambidae	<i>Dichogama colotha</i>	Costa Rica: Guanacaste	ASHYB2444-12
<i>Alabagrus nicoya</i>	DHJPAR0055972	Crambidae	<i>Dichogama redtenbacheri</i>	Costa Rica: Guanacaste	ASHYH2709-14
<i>Alabagrus nicoya</i>	DHJPAR0055976	Crambidae	<i>Dichogama redtenbacheri</i>	Costa Rica: Guanacaste	ASHYH2713-14
<i>Alabagrus nicoya</i>	DHJPAR0055977	Crambidae	<i>Dichogama redtenbacheri</i>	Costa Rica: Guanacaste	ASHYH2714-14
<i>Alabagrus paulgoldsteini</i>	DHJPAR0037908	Crambidae	<i>Syllepte amando</i> DHJ02	Costa Rica: Guanacaste	ASHYC4653-10
<i>Alabagrus paulheberti</i>	DHJPAR0040223	Crambidae	<i>Aponia itzalis</i>	Costa Rica: Guanacaste	ASHYE2390-11
<i>Alabagrus paulheberti</i>	DHJPAR0046746	Crambidae	<i>Aponia minnithalis</i>	Costa Rica: Alajuela	ACGBA919-12
<i>Alabagrus paulheberti</i>	DHJPAR0046953	Crambidae	<i>Aponia minnithalis</i>	Costa Rica: Alajuela	ACGBA1099-12
<i>Alabagrus paulthiaucourti</i>	DHJPAR0045016	Crambidae	<i>Diatraea</i> Janzen989	Costa Rica: Alajuela	ACGAZ237-11
<i>Alabagrus paulthiaucourti</i>	DHJPAR0045021	Crambidae	<i>Diatraea</i> Janzen989	Costa Rica: Alajuela	ACGAZ242-11
<i>Alabagrus ramyamanjunathae</i>	DHJPAR0036701	Crambidae	<i>Phostria temira</i>	Costa Rica: Alajuela	ASHYE1612-09
<i>Alabagrus roibasi</i>	DHJPAR0009358	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASBR625-06
<i>Alabagrus roibasi</i>	DHJPAR0036690	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASHYE1601-09
<i>Alabagrus roibasi</i>	DHJPAR0036691	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASHYE1602-09
<i>Alabagrus roibasi</i>	DHJPAR0036693	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASHYE1604-09
<i>Alabagrus roibasi</i>	DHJPAR0036709	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASHYE1620-09
<i>Alabagrus roibasi</i>	DHJPAR0037882	Crambidae	<i>Phostria oajacalis</i>	Costa Rica: Guanacaste	ASHYC4627-10
<i>Alabagrus roibasi</i>	DHJPAR0046736	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Guanacaste	ACGBA909-12
<i>Alabagrus sanctus</i>	H1339			USA: Kentucky	MF098379
<i>Alabagrus scottmilleri</i>	DHJPAR0030469	Crambidae	<i>Neurophyseta</i> Janzen222	Costa Rica: Alajuela	ASHYB1212-09
<i>Alabagrus scottmilleri</i>	DHJPAR0037942	Crambidae	<i>Neurophyseta</i> BioLep219	Costa Rica: Alajuela	ASHYC4687-10
<i>Alabagrus scottmilleri</i>	DHJPAR0052088	Crambidae	<i>Neurophyseta camptogrammalis</i> DHJ01	Costa Rica: Alajuela	ASHYH1200-13

Appendix 1 Continued.


Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus</i> sp. 1	H7277			Mexico: Yucatan	ATRMK428-11
<i>Alabagrus</i> sp. 1	H874			Mexico: Yucatan	ATRMK248-11
<i>Alabagrus</i> sp. 2	BIOUG24991-C10			Costa Rica: Guanacaste	GMACR176-15
<i>Alabagrus</i> sp. 2	BIOUG25233-E04			Costa Rica: Guanacaste	GMACX053-15
<i>Alabagrus</i> sp. 2	BIOUG27868-B02			Costa Rica: Guanacaste	GMAAT148-16
<i>Alabagrus</i> sp. 2	BIOUG27980-H02			Costa Rica: Guanacaste	GMABS001-16
<i>Alabagrus</i> sp. 2	BIOUG27999-A11			Costa Rica: Guanacaste	GMAAF505-16
<i>Alabagrus</i> sp. 2	BIOUG28111-D02			Costa Rica: Guanacaste	GMADE028-16
<i>Alabagrus</i> sp. 2	BIOUG28111-D03			Costa Rica: Guanacaste	GMADE029-16
<i>Alabagrus</i> sp. 2	BIOUG28160-D10			Costa Rica: Guanacaste	GMADW094-16
<i>Alabagrus</i> sp. 2	H7577			Honduras: Atlantida	ATRMK391-11
<i>Alabagrus</i> sp. 3	H806			Mexico: Yucatan	ATRMK243-11
<i>Alabagrus</i> sp. 4	H11326			USA: Virginia	MF361698
<i>Alabagrus</i> sp. 5	H14623			Mexico: Sonora	MF361705
<i>Alabagrus</i> sp. 6	H12099			Costa Rica: Heredia	MF361700
<i>Alabagrus tanyadapkeyae</i>	DHJPAR0040218	Crambidae	<i>Phostria cyrisalis</i>	Costa Rica: Guanacaste	ASHYE2385-11
<i>Alabagrus tanyadapkeyae</i>	DHJPAR0040225	Crambidae	<i>Phostria cyrisalis</i>	Costa Rica: Guanacaste	ASHYE2392-11
<i>Alabagrus tanyadapkeyae</i>	DHJPAR0055970	Crambidae	<i>Phostria cyrisalis</i>	Costa Rica: Alajuela	ASHYH2707-14
<i>Alabagrus tanyadapkeyae</i>	DHJPAR0057429	Crambidae	<i>Phostria cyrisalis</i>	Costa Rica: Alajuela	ACGBA6201-16
<i>Alabagrus texanus</i>	H11120			USA: Kentucky	MF361696
<i>Alabagrus texanus</i>	H11310			USA: Virginia	MF361697
<i>Alabagrus texanus</i>	H1343			USA: Kentucky	MF361685
<i>Alabagrus texanus</i>	H2735			USA: Kentucky	MF361686
<i>Alabagrus texanus</i>	H2736			USA: Kentucky	MF361687
<i>Alabagrus texanus</i>	H4418			USA: Kentucky	MF361688
<i>Alabagrus texanus</i>	H7301			USA: Kentucky	MF361689
<i>Alabagrus texanus</i>	H7539			USA: Kentucky	MF098381
<i>Alabagrus texanus</i>	H964			USA: Kentucky	MF361678
<i>Alabagrus texanus</i>	H7342			USA: Kentucky	ATRMK454-11
<i>Alabagrus watsoni</i>	DHJPAR0015519	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASAG205-07
<i>Alabagrus watsoni</i>	DHJPAR0015526	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASAG212-07
<i>Alabagrus watsoni</i>	DHJPAR0021196	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASBC1008-07
<i>Alabagrus watsoni</i>	DHJPAR0022189	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASTAT1327-07
<i>Alabagrus watsoni</i>	DHJPAR0023285	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM037-08
<i>Alabagrus watsoni</i>	DHJPAR0023286	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM038-08
<i>Alabagrus watsoni</i>	DHJPAR0023287	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM039-08
<i>Alabagrus watsoni</i>	DHJPAR0023288	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM040-08
<i>Alabagrus watsoni</i>	DHJPAR0023289	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM041-08
<i>Alabagrus watsoni</i>	DHJPAR0023290	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM042-08
<i>Alabagrus watsoni</i>	DHJPAR0023292	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM044-08
<i>Alabagrus watsoni</i>	DHJPAR0023293	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYM045-08
<i>Alabagrus watsoni</i>	DHJPAR0023297	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM049-08
<i>Alabagrus watsoni</i>	DHJPAR0023298	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM050-08
<i>Alabagrus watsoni</i>	DHJPAR0023299	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM051-08
<i>Alabagrus watsoni</i>	DHJPAR0023300	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM052-08
<i>Alabagrus watsoni</i>	DHJPAR0023530	Crambidae	<i>Omiodes humeralis</i> DHJ07	Costa Rica: Alajuela	ASHYM282-08
<i>Alabagrus watsoni</i>	DHJPAR0023531	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYM283-08
<i>Alabagrus watsoni</i>	DHJPAR0023535	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYM287-08
<i>Alabagrus watsoni</i>	DHJPAR0028282	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF044-09
<i>Alabagrus watsoni</i>	DHJPAR0028283	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF045-09
<i>Alabagrus watsoni</i>	DHJPAR0028284	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF046-09
<i>Alabagrus watsoni</i>	DHJPAR0028285	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF047-09
<i>Alabagrus watsoni</i>	DHJPAR0028286	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF048-09
<i>Alabagrus watsoni</i>	DHJPAR0028291	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF053-09
<i>Alabagrus watsoni</i>	DHJPAR0028308	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Guanacaste	ASHYF070-09
<i>Alabagrus watsoni</i>	DHJPAR0037915	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYC4660-10
<i>Alabagrus watsoni</i>	DHJPAR0037936	Crambidae	<i>Omiodes humeralis</i> DHJ02	Costa Rica: Alajuela	ASHYC4681-10
<i>Alabagrus watsoni</i>	DHJPAR0037937	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYC4682-10
<i>Alabagrus watsoni</i>	DHJPAR0039356	Crambidae	<i>Omiodes fulvicauda</i>	Costa Rica: Alajuela	ASAG284-16
<i>Alabagrus watsoni</i>	DHJPAR0041201	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYF1107-11
<i>Alabagrus watsoni</i>	DHJPAR0041202	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYF1108-11
<i>Alabagrus watsoni</i>	DHJPAR0042825	Crambidae	<i>Omiodes insolotalis</i>	Costa Rica: Alajuela	ASHYH583-11
<i>Alabagrus watsoni</i>	DHJPAR0042827	Crambidae	<i>Omiodes insolotalis</i>	Costa Rica: Alajuela	ASHYH585-11
<i>Alabagrus watsoni</i>	DHJPAR0054496	Crambidae	<i>Omiodes humeralis</i>	Costa Rica: Alajuela	ASHYD3661-14
<i>Alabagrus yuanmaofangi</i>	DHJPAR0036334	Crambidae	<i>Desmia</i> biolep02	Costa Rica: Alajuela	ASHYD1525-09
<i>Alabagrus yuanmaofangi</i>	DHJPAR0037894	Crambidae	<i>Desmia</i> Janzen09	Costa Rica: Alajuela	ASHYC4639-10
<i>Alabagrus yuanmaofangi</i>	DHJPAR0037899	Crambidae	<i>Desmia</i> Janzen09	Costa Rica: Alajuela	ASHYC4644-10
<i>Alabagrus yuanmaofangi</i>	DHJPAR0037941	Crambidae	<i>Desmia</i> BioLep06	Costa Rica: Alajuela	ASHYC4686-10

Appendix 1 Continued.


Taxon	Specimen no.	Host family	Host species	Country: region	Accession no.
<i>Alabagrus yuanmaofangi</i>	DHJPAR0042834, H6804	Crambidae	<i>Trichaea pilicornis</i>	Costa Rica: Alajuela	ASHYH592-11
<i>Alabagrus yuanmaofangi</i>	DHJPAR0053600	Crambidae	<i>Desmia ploralis</i> DHJ03	Costa Rica: Alajuela	ASHYM2954-13
<i>Alabagrus yuchinkengae</i>	DHJPAR0045112	Crambidae	<i>Microthyris prolongalis</i>	Costa Rica: Guanacaste	ACGAZ333-11




Supplemental Figure 1 Tree of highest posterior probability from a 10 million generation Bayesian analysis of 681 COI sequences.




Supplemental Figure 1 Continued.




Supplemental Figure 1 Continued.



Supplemental Figure 1 Continued.




Supplemental Figure 1 Continued.




Supplemental Figure 1 Continued.


- 0.001 substitutions/site




Supplemental Figure 1 Continued.




Supplemental Figure 2 Majority-rule consensus tree from a 10 million generation Bayesian analysis of 681 COI sequences.




Supplemental Figure 2 Continued.




Supplemental Figure 2 Continued.




Supplemental Figure 2 Continued.




Supplemental Figure 2 Continued.




Supplemental Figure 2 Continued.




Supplemental Figure 2 Continued.




Supplemental Figure 3 Tree of highest log-likelihood from a 50-replicate ML analysis of 681 COI sequences.




Supplemental Figure 3 Continued.




Supplemental Figure 3 Continued.




Supplemental Figure 3 Continued.




Supplemental Figure 3 Continued.




Supplemental Figure 3 Continued.




Supplemental Figure 3 Continued.




Supplemental Figure 4 Majority-rule consensus tree from a 200-replicate ML bootstrap analysis of 681 COI sequences.



Supplemental Figure 4 Continued.




Supplemental Figure 4 Continued.




Supplemental Figure 4 Continued.

68	Alabagrus	jeanfrancoislandryi	DHJP	0041196	
	Alabagrus	jeanfrancoislandryi	DHJP	0041603	
	Alabagrus	jeanfrancoislandryi	DHJP	0042824	
	Alabagrus	jeanfrancoislandryi	DHJP	0044999	
	Alabagrus	jeanfrancoislandryi	DHJP	0045000	
	Alabagrus	jeanfrancoislandryi	DHJP	0045003	
	Alabagrus	jeanfrancoislandryi	DHJP	0045004	
	Alabagrus	jeanfrancoislandryi	DHJP	0045005	
	Alabagrus	jeanfrancoislandryi	DHJP	0045006	
	Alabagrus	jeanfrancoislandryi	DHJP	0045010	
	Alabagrus	jeanfrancoislandryi	DHJP	0045011	
	Alabagrus	jeanfrancoislandryi	DHJP	0045012	
	Alabagrus	jeanfrancoislandryi	DHJP	0045013	
	Alabagrus	jeanfrancoislandryi	DHJP	0045014	
	Alabagrus	jeanfrancoislandryi	DHJP	0045015	
	Alabagrus	jeanfrancoislandryi	DHJP	0045783	
	Alabagrus	jeanfrancoislandryi	DHJP	0045786	
	Alabagrus	jeanfrancoislandryi	DHJP	0046741	
	Alabagrus	jeanfrancoislandryi	DHJP	0049939	
	Alabagrus	jeanfrancoislandryi	DHJP	0050065	
	Alabagrus	jeanfrancoislandryi	DHJP	0050370	
	Alabagrus	jeanfrancoislandryi	DHJP	0050932	
	Alabagrus	jeanfrancoislandryi	DHJP	0052689	
	Alabagrus	jeanfrancoislandryi	DHJP	0052693	
	Alabagrus	jeanfrancoislandryi	DHJP	0053613	
	Alabagrus	jeanfrancoislandryi	DHJP	0056295	
	Alabagrus	jeanfrancoislandryi	DHJP	0055979	
	Alabagrus	jeanfrancoislandryi	DHJP	0055989	
	Alabagrus	jeanfrancoislandryi	DHJP	0055990	
	Alabagrus	jeanfrancoislandryi	DHJP	0056296	
	Alabagrus	jeanfrancoislandryi	DHJP	0056298	
	Alabagrus	jeanfrancoislandryi	DHJP	0056994	
	Alabagrus	jeanfrancoislandryi	DHJP	0056999	
	Alabagrus	jeanfrancoislandryi	DHJP	0057430	
	Alabagrus	jeanfrancoislandryi	DHJP	0057431	
	Alabagrus	jeanfrancoislandryi	DHJP	0057738	
	Alabagrus	jeanfrancoislandryi	DHJP	0057745	
	Alabagrus	jeanfrancoislandryi	DHJP	0059747	
	51	Alabagrus	jeanfrancoislandryi	H831	
		Alabagrus	jeanfrancoislandryi	H873	
		Alabagrus	jeanfrancoislandryi	H7628	
		Alabagrus	jeanfrancoislandryi	H7630	
		Alabagrus	jeanfrancoislandryi	H7631	
		Alabagrus	janzeni	DHJP	0009348
		Alabagrus	janzeni	DHJP	0009354
		Alabagrus	janzeni	DHJP	0009355
		Alabagrus	janzeni	DHJP	0009356
		Alabagrus	janzeni	DHJP	0009362
		Alabagrus	janzeni	DHJP	0009374
Alabagrus		janzeni	DHJP	0009375	
Alabagrus		janzeni	DHJP	0009376	
Alabagrus		janzeni	DHJP	0009400	
Alabagrus		janzeni	DHJP	0009403	
Alabagrus		janzeni	DHJP	0009433	
Alabagrus		janzeni	DHJP	0009434	
Alabagrus		janzeni	DHJP	0010083	
Alabagrus		janzeni	DHJP	0010509	
Alabagrus		janzeni	DHJP	0010510	
Alabagrus		janzeni	DHJP	0016416	
Alabagrus		janzeni	DHJP	0016417	
Alabagrus		janzeni	DHJP	0016428	
Alabagrus		janzeni	DHJP	0016429	
Alabagrus		janzeni	DHJP	0016430	
Alabagrus		janzeni	DHJP	0016431	
Alabagrus		janzeni	DHJP	0016432	
Alabagrus		janzeni	DHJP	0015384	
Alabagrus		janzeni	DHJP	0015493	
Alabagrus		janzeni	DHJP	0015494	
Alabagrus		janzeni	DHJP	0015495	
Alabagrus		janzeni	DHJP	0015496	
Alabagrus		janzeni	DHJP	0015497	
Alabagrus		janzeni	DHJP	0015498	
Alabagrus		janzeni	DHJP	0015499	
Alabagrus		janzeni	DHJP	0015511	
Alabagrus		janzeni	DHJP	0015512	
Alabagrus		janzeni	DHJP	0015513	
Alabagrus		janzeni	DHJP	0015515	
Alabagrus		janzeni	DHJP	0016926	
Alabagrus		janzeni	DHJP	0016928	
Alabagrus		janzeni	DHJP	0016930	
Alabagrus		janzeni	DHJP	0021149	
Alabagrus		janzeni	DHJP	0021208	
Alabagrus		janzeni	DHJP	0021211	
Alabagrus		janzeni	DHJP	0029179	
Alabagrus		janzeni	DHJP	0036363	
Alabagrus		janzeni	DHJP	0036365	
Alabagrus		janzeni	DHJP	0036366	
90	Alabagrus	janzeni	DHJP	0036697	
	Alabagrus	janzeni	DHJP	0036698	
	Alabagrus	janzeni	DHJP	0036699	
	Alabagrus	janzeni	DHJP	0036712	
	Alabagrus	janzeni	DHJP	0036716	
	Alabagrus	janzeni	DHJP	0036718	
	Alabagrus	janzeni	DHJP	0037190	
	Alabagrus	janzeni	DHJP	0037902	
	Alabagrus	janzeni	DHJP	0037922	
	Alabagrus	janzeni	DHJP	0041166	
	Alabagrus	janzeni	DHJP	0041167	
	Alabagrus	janzeni	DHJP	0041168	

Supplemental Figure 4 Continued.



Supplemental Figure 4 Continued.



Supplemental Figure 4 Continued.